



at Maxima, E

$$f'(x) \Big|_E = 0$$


$$f''(x) \Big|_E < 0$$



at minima E ,

$$f'(x) \Big|_E = 0$$

$$f''(x) \Big|_E > 0$$



① Vander Wall's Equation for Real Gas :

$$\left(P + \frac{a}{V^2} \right) (V - b) = RT$$

This equation is for 1 mole of real gas.

If moles are n and given volume is V_0

$$\Rightarrow \text{Volume occupied by 1 mole} = \left(\frac{V_0}{n} \right)$$

\Rightarrow Equation for 1 mole becomes same:

$$\left[P + \frac{a}{\left(\frac{V_0}{n} \right)^2} \right] \left[\frac{V_0}{n} - b \right] = RT$$

② Similarly $PV = RT$ is equation for 1 mole of ideal gas. If n moles and V_0 volume is given \Rightarrow Volume occupied by 1 mole of ideal gas $= \left(\frac{V_0}{n} \right)$. Pressure and Temp. remain same

throughout

$$\Rightarrow P \left(\frac{V_0}{n} \right) = RT$$

③ Avogadro's Number = 6×10^{23} molecules /mole $\oplus n = \frac{1}{\sqrt{2\pi d^2 n}}$

④ Helium has 2 isotopes (${}_{2}^{3}\text{He}$ and ${}_{2}^{4}\text{He}$) which behave quite differently at low temperatures. This is because ${}_{2}^{3}\text{He}$ is a fermion ($s = \frac{1}{2}$) while ${}_{2}^{4}\text{He}$ is a boson ($s = 1$)

★ NTP : Normal Temperature & Pressure (20°C , 1 atm)

★ STP : Standard Temp. & Pressure (0°C , 1 atm)

SATP : Standard Ambient Temp. & Pressure (25°C , 1 atm)

★ 31.4°C : Critical Temp. for CO_2 $\oplus b = 4 V_{(\text{molecules})}^{\text{ave}}$

26/12/2011

Section B

(Q5) 3 bits + 3 bits

Units of viscosity

CGS

η : POISE

SI

η : POISEUILLE (PL)

(Q6) Elec & Magn.

$\frac{\eta}{\rho}$: STOKE

where $\eta = \frac{F}{A(\frac{dy}{dz})}$, ρ : density

(Q7) EM Theory + (1 bit at max of Thermal Physics)

(Q8) Thermodynamics

15 LECTURES

$\Rightarrow 60 + 30 + 10 = 100$ marks of Thermo

(a) Thermodynamics

① Basic Concepts : Terminology, Laws of thermodynamics, Work, different thermodynamic Processes.

② Heat Engines : Otto & Diesel Engines

③ Entropy

④ Thermodynamic Relations : Maxwell Relations

→ Clapeyron Eqn.

→ Joule Kelvin

→ Adiabatic demagnetization

⑤ VanderWaals Eqn

⑥ Specific Heat of Solids

⑦ Misc. Gibbs Phase Rule / Transport Phenomenon

(b) Statistical Physics



- ① Maxwell - Boltzmann Statistics (Concept of negative temperature)
- ② Fermi Dirac Statistics (Black Body Radiation)
- ③ Bose Einstein Statistics (Specific Heat of Gases)

14 - 15 classes

[B.Sc Honours Course of Thermodynamics]



- ④ Vapour: usually applied to gas in equilibrium with its liquid (i.e. saturated vapour) or to a gas at a temperature below T_c . Its properties are similar to a gas.

- ⑤ For ideal gas



- ⑥ T_b : Boyle's Temperature, temp. at which Boyle's law holds for fairly wide range of Pressure.

- ⑦ Triple Point of water : $0.01^\circ C$, 4.58 mm of mercury
It's used to define Kelvin scale of Temperature.



Thermodynamics (1)

26/12/2011

Heat is a form of energy. Heat can be converted into Work and Work can be converted into Heat. This interconversion is the domain of thermodynamics.

3 techniques to study these interactions

1) Thermometry

- ✓ direct measurement of parameters and getting the relations, thereby.
- ✓ we are not concerned about constituents of system.
i.e Not concerned about individual particles' behaviour
- ✓ we are interested in Macro Variables.


2) Kinetic Theory of Gases

- ✓ trying to track down behaviour of individual molecules.
- ✓ here also we can account for macro variable using integrated effect of individual molecules.

3) Statistical Thermodynamics:

- ✓ in between the previous 2 approaches
- ✓ we measure things at equilibrium: the most probable state of matter.
- ✓ we study aggregate behaviour or expectation values.

Thermodynamic System



"A definite quantity of masses (i.e can exchange mass and energy) with a distinct boundary."

Depending upon exchange of mass & energy b/w system and surrounding, we have :

- 1) Open system : Both energy & mass can be exchanged
- 2) Closed : Energy interaction allowed
- 3) Isolated : No interaction allowed.

Temperature and other thermodynamic variables

How hot or cold the system : Temperature

State of any system can be defined in terms of thermodynamic variables. They are of 2 types:

- 1) Intensive : independent of mass eg. Specific Heat Capacity
- 2) Extensive : dependend on mass eg. Heat Capacity

Also, we have variables as :

- 1) Macro : can be perceived by sense : aggregate behaviour
- 2) Micro : behaviour of individual particles

These variables are measured by the 3 techniques mentioned before.

Equation of state is nothing but relationship between thermodynamic variables.

$$\text{eg. } f(P, V, T) = 0$$

eg for an ideal gas : $PV = nRT$

$$\begin{aligned} \text{Boyle's law} \quad P \propto \frac{1}{V} \\ \text{Charles' law} \quad V \propto T \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{from Thermometry}$$

$$\begin{aligned} P &= \frac{1}{3} N m \frac{\bar{v}^2}{V} \\ \frac{1}{2} m \bar{v}^2 &= \frac{3}{2} kT \\ \Rightarrow PV &= nRT \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{from kinetic Theory}$$

or described by

If thermodynamic system is dependent upon 3 variables
⇒ only 2 are independent. Hence state is known only via 2 variables.

All the thermodynamic variables are measured at equilibrium.

Thermodynamic Equilibrium

It means system is in : thermal equilibrium
mechanical equilibrium
chemical equilibrium

① Thermal Equilibrium : NO temperature gradient inside the system i.e. no heat flow within the system. Thermal Equilibrium defined by a particular temperature.

Concept of Temperature comes from 0th law of temperature.


② Mechanical Eq^b : NO unbalanced forces i.e. Pressure throughout the system is same.

③ Chemical Eq^b : There is no chemical concentration in any part of system. Volume is measured at chemical eq^b.

If there are 2 eq^b, 3rd eq^b. will automatically follow.

Laws of Thermodynamics

1st law : Statement of Conservation of Energy.



$$\boxed{\Delta Q = \Delta U + \Delta W}$$

Heat given to system : +ve
Work done by system : +ve


$$\Rightarrow \Delta U = \Delta Q - \Delta W \quad \begin{matrix} \uparrow & \uparrow \\ \Delta (\text{internal energy}) & \text{heat given to system} \end{matrix}$$

Work done by system

We know

$$\begin{aligned}\Delta W &= F \cdot dx \\ &= P A dx \\ &= P dV\end{aligned}$$

the sum of total kinetic and potential energies of all molecules constitutes the internal energy of the system.



For solid $\Delta W = 0$

$$\Rightarrow \Delta Q \propto \Delta T$$

$\Delta Q \propto$ mass

$$\Delta Q = \underset{\text{heat capacity } (s)}{\cancel{(m s)}} \underset{\text{specific heat } (s) \text{ or } (c)}{\cancel{\Delta T}}$$

$$s = \frac{1}{M} \left(\frac{\Delta Q}{\Delta T} \right)$$

Heat Capacity :-

Extensive Variable

Specific Heat Capacity
intensive variable

Solids have only 1 specific heat capacity since their volume is assumed to be const.

Thermodynamic Process

Going from 1 thermodynamic state of equilibrium to other equilibrium state.

We know $(n-1)$ independent variables, 1 dependent variables. Usually the latter variable is kept const. during any thermodynamic process.

Process is named after that variable which is kept const.

eg.	Isochoric	Adiabatic
	Isobaric	Isothermal

e.g. Volume is kept const. for determining specific heat of solids.

$$C_V = \left(\frac{dQ}{dT} \right)_V \quad \text{i.e. } dQ = C_V dT$$

Similarly we can define

$$C_P = \left(\frac{dQ}{dT} \right)_P \quad \text{i.e. } dQ = C_P dT$$

$C_P = C_V$ in case of solids

$C_P \neq C_V$ in case of liquids and gases.

all: Internal Energy or Intrinsic Energy

Characteristics of molecules

$$dU = dU_{K.E.} + dU_{P.E.}$$

↑
due to
motion of molecules

↑
due to interaction
between molecules

If there is no interaction between particles $dU_{p.e.} = 0$
 eg. Ideal Gas have non-interacting particles.

$$dU_{\text{ideal gas}} = dU_{k.e.}$$

$$dU_{k.e.} = f(T)$$

for ideal gas : $U = U(T)$: for non interacting molecules

$$\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = 0 \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{Joule's law}$$

$$\Rightarrow \left(\frac{\partial U}{\partial P}\right)_T = 0$$

Inter Molecular Interactions are characterized by volume

$$\Rightarrow U = U(T, V) : \text{for interacting}$$

[we could have taken (T, P) also, but we love $f(T, V)$] molecules

$$dU = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

U is a state function i.e. dU is dependent upon initial and final equilibrium positions and independent of the process taken.

$$dQ = dU + P dV$$

$$C_V dT = dU + P \cdot 0$$

\Rightarrow

$$dU = C_V dT$$

① We have calculated $dU = C_V dT$ for $dV = 0$
 If we assume dU as $f(T)$ only, then we can say for same dT (as U is a state function)
 $U = C_V dT$ (always)

$= dU$ But if we assume $U = f(T)$, as is the case normally, then we need to write
 $dU = \left(\frac{\partial U}{\partial V}\right) dV + \left(\frac{\partial U}{\partial T}\right) dT$
 & cannot write in general $dU = C_V dT$

$$dQ = dU + dW$$

$$\Rightarrow C_p dT = C_v dT + P dV \quad (\text{at const. Pressure})$$

$$\Rightarrow C_p = C_v + P \left(\frac{dV}{dT} \right)_P$$

In case of solids $\left(\frac{dV}{dT} \right)_P = 0 \Rightarrow C_p = C_v$

For ideal gases $PV = RT \Rightarrow \left(\frac{dV}{dT} \right)_P = \left(\frac{R}{P} \right)$

$$\Rightarrow C_p = C_v + R$$

0th law of thermodynamics

If 2 systems A and B are in thermal equilibrium and 2 systems B and C are in thermal equilibrium, this implies A and C are in thermal equilibrium.

\Rightarrow At ~~some~~ thermal equilibrium, some thermodynamic property is const.

We define that property as Temperature. T

$$\Rightarrow \begin{aligned} &\text{if } T_A = T_B && (13, 9) \Leftrightarrow \text{coprime} \\ &\text{and } T_B = T_C && (9, 26) \Leftrightarrow \text{coprime} \\ &\Rightarrow T_A = T_C && \text{but } (13, 26) \not\leftrightarrow \text{coprime} \end{aligned}$$

★ Note that if volume change = 0, then $dU = C_v dT$ always. But if volume change $\neq 0$, then

$$dU = C_v dT + \left(\frac{\partial U}{\partial V} \right)_T \cdot dV \quad \text{in general case}$$

For ideal gas, Joule's law states that $\left(\frac{\partial U}{\partial V} \right)_T = 0 \Rightarrow dU = C_v dT$

2nd law of thermodynamics.

100% of ~~heat~~ conversion of heat into work is not possible.

[Heat \rightarrow Work Conversion] via Heat Engine.


[Work \rightarrow Heat] via Refrigerator.

$$\eta = \frac{\text{Output}}{\text{Input}} = \frac{W}{Q}$$

$$\eta < 1$$

Kelvin Planck Statement of 2nd law.

"It is impossible to construct a device whose sole job is to take heat from source at higher temperature and perform equivalent amount of work w/o producing any other changes into parts of system."



$$\eta = \frac{W}{Q_1}$$

$$Q_1 = Q_2 + W \Rightarrow \eta = \left(\frac{Q_1 - Q_2}{Q_1} \right) = \left[1 - \left(\frac{Q_2}{Q_1} \right) \right]$$


For $\eta = 100\% \rightarrow Q_2 = 0$ which is contradicted by Kelvin Planck statement.

~~Perpetual Machine of 1st kind~~: Heat Engine

~~Perpetual Machine of 2nd kind~~: Refrigerator

"Carnot - Ostwald Statement"

"It is impossible to construct a device whose sole job is to get heat from a lower temperature, ~~and release the heat at~~ ~~higher~~ and release the heat at ~~higher~~ temperature."



Ostwald Statement

"It is impossible to have a perpetual machine of 2nd kind."

3rd law of thermodynamics

(Corollary to)
2nd law

- Absolute Zero is unattainable in a finite number of steps.

OR

- Entropy of a system cannot be reduced to zero in a finite number of operations.

OR

- Entropy of a perfect crystal at absolute zero is exactly equal to zero.

$$\eta < 1$$

$$1 - \frac{T_2}{T_1} < 1$$

$$\Rightarrow T_2 > 0 \text{ always}$$

Thermodynamic Processes for Ideal Gas (Only 1 available)
microstate

For an ideal gas : $PV = RT$ (for 1 mole)

Isothermal

$$PV = \text{const.} = RT$$

$$\rightarrow dU = C_V dT = 0 \quad (\text{ideal gas; otherwise } \left(\frac{\partial U}{\partial V}\right)_T dV \text{ term also})$$

$$\rightarrow dW = PdV = \left(\frac{RT}{V}\right) dV = RT \ln\left(\frac{V_2}{V_1}\right) = RT \ln\left(\frac{P_1}{P_2}\right)$$

$V_2 > V_1 \Rightarrow$ Gas expanding \Rightarrow +ve work by system

$V_1 > V_2 \Rightarrow$ Gas compressing \Rightarrow -ve work by system
by external force

$$\rightarrow dQ = dW = RT \ln\left(\frac{V_2}{V_1}\right)$$

Isochoric

$$PV = RT \Rightarrow \frac{P}{T} = \frac{R}{V} = \text{const.}$$

$$dW = PdV = 0$$

$$dU = C_V dT$$

$$dQ = dU = C_V dT$$

Iobaric

$$dW = PdV = P(V_2 - V_1)$$

$$dQ = C_P dT$$

$$dU = dQ - dW$$

Adiabatic

$$dQ = 0$$

$$dW = -dU$$

$$dU = P dV$$

$$dU = -P dV$$

Perpetual Motion Machine of 1st type produces work w/o the input of energy. It thus violates the 1st law of thermodynamics.

Perpetual Motion Machine of 2nd type spontaneously converts thermal energy into mechanical work. When thermal energy is equivalent to work done, this does not violate the law of conservation of energy. However, it violates the 2nd law of thermodynamics.

- We know η of a heat engine will always be less than 1. This means that to get some useful work done, some energy is always lost. The lost energy can add to thermal pollution in our surrounding leading to global warming.... We say so because usually source is not atmosphere but sink is more often than not, the atmosphere.

Thermodynamics (2)

27/12/2011

For adiabatic, we can derive 1 more equation for an ideal gas :-

$$dQ = 0$$

$$\Rightarrow dU + PdV = 0 \Rightarrow C_V dT + PdV = 0$$

$(dU \xrightarrow{\text{constant}} C_V dT)$
means you are talkin about Ideal Gas

$$\Rightarrow C_V dT + \frac{RT}{V} dV = 0$$

(we can replace P in real gas)
also

$$\Rightarrow C_V \left(\frac{dT}{T} \right) + R \left(\frac{dV}{V} \right) = 0$$

P + a

$$\Rightarrow \left(\frac{dT}{T} \right) + \left(\frac{R}{C_V} \right) \left(\frac{dV}{V} \right) = 0$$

$$\frac{C_P}{C_V} = \gamma \quad (\text{dependent on atomicity of gases})$$

Its called gamma

$$C_P - C_V = R$$

$$\Rightarrow (\gamma - 1) C_V = R$$

$$\Rightarrow \frac{R}{C_V} = (\gamma - 1)$$

$$\boxed{\gamma > 1}$$

$$\Rightarrow \left(\frac{dT}{T} \right) + (\gamma - 1) \left(\frac{dV}{V} \right) = 0$$

$$\Rightarrow \int \frac{dT}{T} + (\gamma - 1) \int \frac{dV}{V} = \ln(c) \quad (\text{say})$$

$$\Rightarrow \ln T + (\gamma - 1) \ln V = \ln(c)$$

$$\Rightarrow \boxed{TV^{\gamma-1} = c}$$

For ideal gas

$$\frac{PV}{R} \cdot V^{\gamma-1} = c \Rightarrow$$

$$\boxed{PV^\gamma = \text{const}}$$

isobaric

isothermal

adiabatic

isochoric



$PV^n = \text{const}$: depending on n

it can represent any process


(0, 1, ∞, any other)

$$\star \Delta PV^n + P n V^{n-1} dV = 0 \Rightarrow dV = -\frac{V}{nP} dP \quad \text{if } dV=0 \Rightarrow n \rightarrow \infty \Rightarrow \text{isochoric}$$

for
 $n=\infty$

Indicator diagram (or PV diagram)

Indicator diagram can be b/w (P, T) or (T, V) also.
But area under PV curve gives Work. Hence preferred more.



If we are able to measure P and V at every point i.e. a continuous curve, this implies process goes through infinite quasi-static processes. st. equilibrium is maintained at all points. Quasi-static Process are carried very slowly. At all points condition of equilibrium holds.

✓ Also area under curve gives Work Done.

✓ Also the direction of curve indicates expansion / contraction.

slopes for different processes

For an isothermal process,

$$PV = \text{const}$$

$$\Rightarrow PdV + VdP = 0$$

$$\Rightarrow \left(\frac{dP}{dV}\right)_T = -\left(\frac{P}{V}\right)$$

For an adiabatic process,

$$PV^r = \text{const.}$$

$$\Rightarrow (r-1)PV^{r-1}dV + V^r dP = 0$$

$$\Rightarrow \left(\frac{dP}{dV}\right)_{\text{adiabatic}} = -r\left(\frac{P}{V}\right) \text{ more slope than isothermal}$$

For an isobaric process,

$$dP = 0$$

$$\Rightarrow P = \text{const} \Rightarrow \left(\frac{dP}{dV}\right)_P = 0$$

For an isochoric process

$$dV = 0$$

$$\Rightarrow \left(\frac{dP}{dV}\right)_V = \infty$$

Work done in ideal gas undergoing adiabatic process

$$PV^r = \text{const}$$

$$\Rightarrow P = k V^{-r}$$


For adiabatic (ideal) : $-dV = P dV$
 $\Rightarrow W = - \int dV = -C_V \Delta T$

$$= C_V (T_1 - T_2)$$

$$= \frac{R}{r-1} (P_1 V_1 - P_2 V_2)$$

$$W = P dV = k V^{-r} dV = \frac{k V^{(r+1)}}{(r-1)} \Big|_{V_1}^{V_2} = k \left(\frac{V_2^{1-r} - V_1^{1-r}}{1-r} \right)$$

$$= \frac{P_2 V_2^{1-r} - P_1 V_1^{1-r}}{1-r} = \left[\frac{P_2 V_2 - P_1 V_1}{1-r} \right]$$




For an ideal gas,

$$W = -R \frac{(T_2 - T_1)}{(r-1)} = \frac{R (T_1 - T_2)}{(r-1)}$$

$$W = \frac{R}{(r-1)} (T_1 - T_2)$$

In adiabatic Process 'a',
 Temperature at any point along AB is less than TA
 i.e. $T_A \downarrow \Rightarrow V \downarrow \Rightarrow dV < 0$. since $dQ = 0$
 $\Rightarrow dW > 0$ in this process
 Reverse is true for process -a'
~~∴ Area under the curve is negative!!~~

Reversible Process :



If I am able to retrace back from f to i backwards, then its reversible process.

It is ideal concept.

All processes are irreversible in nature

Reversible Process assumes 0 dissipative losses. In real processes, some dissipation always occurs. In indicator diagrams, the Quasi-static Processes are assumed to be reversible.

Some Maths.....
 if $Z = f(x, y)$

$$\text{we know, } dz = \left(\frac{\partial f}{\partial x}\right)_y dx + \left(\frac{\partial f}{\partial y}\right)_x dy$$

log dot graph

$\frac{\log n > 0}{\log n < 0}$ for $n >$
 for $n <$

Usually z is const. We always separate out the constant variable $\Rightarrow dz = 0$
 eg. for Isothermal Process

→ const. वाले को separate out कर लेते हैं और उसी के नाम पर Process का नाम रख देते हैं !!

$$T = f(P, V)$$

$$\Rightarrow 0 = \left(\frac{\partial f}{\partial x}\right)_y dx + \left(\frac{\partial f}{\partial y}\right)_x dy$$

$$\Rightarrow \left(\frac{dy}{dx}\right)_z = - \frac{\left(\frac{\partial f}{\partial x}\right)_y}{\left(\frac{\partial f}{\partial y}\right)_x} \quad \text{or} \quad \left(\frac{dy}{dx}\right)_z = - \frac{\left(\frac{\partial z}{\partial x}\right)_y}{\left(\frac{\partial z}{\partial y}\right)_x}$$

remember by Physical interpretation


$$\begin{aligned} \text{if } x \uparrow &\Rightarrow z \uparrow \Rightarrow \left(\frac{\partial z}{\partial x}\right) > 0 \\ y \uparrow &\Rightarrow z \downarrow \Rightarrow \left(\frac{\partial z}{\partial y}\right) < 0 \end{aligned}$$

Heat Engine

Heat Engine has 3 components :

- 1) Source ' T_1 '
- 2) Working substance
- 3) Sink ' T_2 ' ($T_2 < T_1$)

$$\begin{aligned} \text{For const. } z \\ x \uparrow \text{ and } y \uparrow \\ \Rightarrow \left(\frac{\partial z}{\partial x}\right) > 0 \end{aligned}$$



Assuming heat absorption is ideal process, i.e. no wastage in conduction, radiation etc.

$$\Rightarrow Q_1 = W + Q_2$$

$$\eta = \left(\frac{W}{Q_1} \right) = \frac{Q_1 - Q_2}{Q_1} = 1 - \left(\frac{Q_2}{Q_1} \right)$$

(ISOTHERMAL + ADIABATIC)

Carnot Engine




diagram of
2nd lecture

In reality it is not possible a thermodynamically reversible engine, so practical application of Carnot engine is restricted.


[Isothermal Process are VERY slow processes]
[Adiabatic Process are VERY FAST processes.]

While supplying heat, conducting walls of source and cylinder are kept next to each other. Isothermal expansion occurs.

When adiabatic process is started, Insulating stand is kept next to cylinders conducting wall. Heat supply is cut off.

While removing heat, sink is brought in contact with cylinder. Isothermal contraction occurs.

Finally insulation stand is again brought to achieve adiabatic compression.



We should start the cycle with this point for sake of analysis

In adiabatic expansion, gas works on its own and hence useful work is done. Temperature falls down as internal energy is used up for performing the expansion or +ve work. Hence $T_C < T_B$.

In BC, useful work has been done. Work of engine has been done. Now we have to bring back the system to previous state to again do the work.

CD : isothermal contraction

DA : adiabatic

$$\text{Work done} = \Delta(ABCD)$$

Note that all processes are assumed to be reversible.

Net Work Done

$$\underline{AB} \quad dV = 0$$

$$dW_1 = Q_1 = RT_1 \ln\left(\frac{V_2}{V_1}\right)$$

$$\underline{BC} \quad dQ = 0$$

$$dW_2 = \frac{R}{(\gamma-1)} (T_1 - T_2)$$

CD $dU = 0$

$$dW_3 = Q_2 = T_2 R \ln \left(\frac{V_4}{V_3} \right) \quad \leftarrow \text{all the terms are <}$$

DA $dQ = 0$

$$dW_4 = \frac{R}{(r-1)} (T_2 - T_1)$$

$$\text{Net Work Done} = RT_1 \ln \left(\frac{V_2}{V_1} \right) + RT_2 \ln \left(\frac{V_4}{V_3} \right)$$

We know $T_1 V_2^{r-1} = T_2 V_3^{r-1}, \quad T_1 V_1^r = T_2 V_4^r$

$$\Rightarrow \left[\frac{V_1}{V_2} \right] = \left[\frac{V_4}{V_3} \right]$$

Use TV^{r-1} form

$$\Rightarrow \boxed{\text{Net Work Done} = R(T_1 - T_2) \ln \left(\frac{V_2}{V_1} \right)}$$

Also from Energy supplied

$$\begin{aligned} Q &= RT_1 \ln \left(\frac{V_2}{V_1} \right) = RT_2 \ln \left(\frac{V_4}{V_3} \right) \\ &= R(T_1 - T_2) \ln \left(\frac{V_2}{V_1} \right) \end{aligned}$$

$dU = 0$ for complete cycle

$$\Rightarrow \boxed{dQ = dW}$$

$$\eta = \frac{W}{Q_1} = \frac{R(T_1 - T_2) \ln \left(\frac{V_2}{V_1} \right)}{RT_1 \ln \left(\frac{V_2}{V_1} \right)} = \frac{T_1 - T_2}{T_1}$$

$\Rightarrow \eta = 1 - \left(\frac{T_2}{T_1} \right)$

$$\textcircled{O} \quad \eta = 1 - \frac{Q_2}{Q_1} \quad \text{in general}$$

$$= 1 - \left(\frac{T_2}{T_1} \right) \quad \text{for Carnot engine}$$

→ Remember that Carnot theorem has 2 components

- ★ All the reversible engines operating between same source and same sink will have same efficiency equals to $\left[1 - \left(\frac{T_2}{T_1} \right) \right]$: Carnot Theorem (1)

- ★ Irreversible Engine will have lower efficiency than the reversible engine working between same sink & source

$$\text{i.e. } \eta_{\text{reversible}} > \eta_{\text{irreversible}}$$

$$\text{i.e. } \left(1 - \frac{Q_2}{Q_1} \right)_{\text{irreversible}} < \left(1 - \frac{T_2}{T_1} \right), \quad \begin{matrix} \text{Proof in} \\ \text{lecture} \\ 4 \end{matrix}$$

Note that both proofs follow the same

things of
a flying
Opposite Carnot
cycle

- ★ For reversible cyclic engine,

$$\frac{Q_2}{Q_1} = \frac{T_2}{T_1} \Rightarrow \frac{Q_2}{T_2} = \frac{Q_1}{T_1}$$

→ We can give the logic that any closed reversible process that be thought of consisting of many Carnot cycles $\Rightarrow \oint \frac{dQ}{T} = 0$

For irreversible


~~For irreversible~~
i.e.
operating
high η as
engine and
lower η as
refrigerator!!

$$\frac{Q_2}{Q_1} > \frac{T_2}{T_1} \Rightarrow \frac{Q_2}{T_2} > \frac{Q_1}{T_1}$$

$$d\left(\frac{Q}{T}\right) = \left[\frac{Q_1}{T_1}\right] + \left[-\frac{Q_2}{T_2}\right] < 0 \Rightarrow$$

$$\Rightarrow \oint \frac{dQ}{T} < 0$$

irreversible



$\frac{dQ}{T} \leq 0$ (Carnot theorem)
at $T_{\text{high}} > T_{\text{low}}$

$$\Rightarrow \int \frac{dQ}{T} \leq 0$$

(equality for reversible process)

Irreversible के लिए $\frac{dQ}{T}$ कम है
 $\Rightarrow -[dQ] \text{ का कोई component नहीं है}$ for some incoming heat
 $\Rightarrow \int \frac{dQ}{T} < 0$

Clausius Inequality

④ Note that Clausius Inequality talks nothing about entropy change. It talks only about $\int \frac{dQ}{T}$


Internal Combustion Engines

Within cylinder-piston arrangement, provision of burning of fuel.

2 examples of ICEs are Otto Engine and Diesel Engine.

Otto Engine

(ISOCHORIC + ADIABATIC)



Working substance: 1 gram-mole of air

\rightarrow mole = gram-mole
 $=$ gram-molecule
 $=$ avogadro no. of molecules

\rightarrow Otto engine designed by Nicholas Otto in 19th century

\rightarrow Modified, Advanced and fuel controlled versions of Otto Engines are used in cars all over the world.

4 strokes, 2 valves engine


↳ Suction / Intake Stroke
 Compression

Working (Ignition)

Exhaust (Heat Release)

\rightarrow Also called Petrol or Gasoline engine.

① At the atmospheric pressure, 'I' is opened and a mixture of air and fuel is sucked in. T_1 : room temp.



② 2nd stroke is adiabatic compression
Maximum compression can be 6 times.

$$\text{Compression Ratio} = \left(\frac{V_1}{V_2} \right)^{\frac{1}{r-1}} < 6$$

More than 6, Pressure will be huge s.t. walls will burst.

$$T_1 V_1^{r-1} = T_2 V_2^{r-1}$$

$$T_2 = T_1 \left(\frac{V_1}{V_2} \right)^{r-1} = T_1 (6)^{0.5} = 2.4 T_1$$

③ Isochoric heat addition is done in 3rd stroke
Compressed mixture is exploded by an electric spark to release heat which increases P and T. Piston is driven out adiabatically.

Temperature is increased further.

Before commencing with derivation, it's mandatory to mention:

- ① Use 4 strokes, with diagram

④ Adiabatic Expansion Occurs due to high temperature. Crank is thrown out. Useful work is done in this step.

⑤ Isochoric Heat Rejection. This is the 4th stroke

$$\eta = 1 - \frac{Q_2}{Q_1} = 1 - \frac{C_v(T_4 - T_1)}{C_v(T_3 - T_2)}$$

$$\boxed{\eta = 1 - \frac{(T_4 - T_1)}{(T_3 - T_2)}}$$

↑
Beware of not writing $1 - \frac{C_v(T_1 - T_4)}{C_v(T_3 - T_2)}$

$$T_1 V_1^{\gamma-1} = T_2 V_2^{\gamma-1}$$

$$T_3 V_3^{\gamma-1} = T_4 V_4^{\gamma-1}$$

$$V_3 = V_2$$

$$V_4 = V_1$$

as ' Q_2 ' is a ~~positive~~ number
 $\Rightarrow -Q_2 = dW = C_v(T_1 - T_4)$
 $\Rightarrow Q_2 = \underline{C_v(T_4 - T_1)}$

(use ~~TV~~ $PV^{\gamma-1}$ form)

$$(T_4 - T_1) V_1^{\gamma-1} = (T_3 - T_2) V_2^{\gamma-1}$$

$$\Rightarrow \boxed{\eta = 1 - \left(\frac{V_2}{V_1} \right)^{\frac{1}{\gamma-1}}} = 1 - \frac{1}{(P)^{\frac{1}{\gamma-1}}}$$

$$\eta_{\max} = 1 - \frac{1}{6^{\frac{1}{\gamma-1}}} \approx 52\% \quad (\text{for } \gamma = 1.4)$$

Note that compression ratio is, UNLIKE WHAT ITS NAME SUGGESTS, $\rho = \left(\frac{V_{\text{more}}}{V_{\text{less}}} \right)$


★ There is a single ρ for both expansion & compression, therefore the 2nd and 4th process need to be ISOCHORIC.

Compression Ratio, expansion ratio etc. are always greater than unity. $\frac{P_e}{P_c} > 1$

Diesel Engine

(ADIABATIC, ISOBARIC, ADIABATIC, ISOCORIC)

- ① In this engine, air is compressed sufficiently to heat it enough to ignite the fuel & then fuel is sprayed in.




- ② Compression upto 34 atm.


✓ Walls have capacity to bear 34 atm.

✓ $\left(\frac{V_3}{V_2}\right)$ can be as high as 17

- Designed by Rudolph Diesel at end of 19th century.
- More Efficient than Otto Engine
- Diesel engines are found in virtually all heavy duty applications e.g. truck, ships



* Difference only in starting phase of Power stroke



$$\begin{aligned} \rightarrow P_3 &= P_2 \\ \text{isobaric expansion} \\ \Rightarrow V \propto T \text{ i.e. } \frac{V_3}{V_2} &= \frac{T_3}{T_2} \end{aligned}$$

isochoric heat rejection

$$V_1 = V_4$$

- ③ Note that after AB, $T \approx 1000^\circ\text{C}$, $P \approx 34 \text{ atm}$, $\rho_c \approx 17$, Now at this stage fuel is injected & combustion takes place.

$$Q_1 = C_p (T_3 - T_2)$$

★ Compression & Expansion

$$Q_2 = C_v (T_4 - T_1)$$

Ratios are defined only for adiabatic compressions and expansions....

$$\eta = 1 - \frac{C_v (T_4 - T_1)}{C_p (T_3 - T_2)}$$

$$= 1 - \frac{\frac{1}{r} \left[\left(\frac{1}{P_e}\right)^r - \left(\frac{1}{P_c}\right)^r \right]}{\left(\frac{1}{P_e}\right) - \left(\frac{1}{P_c}\right)}$$

Compression > Expansion $\Rightarrow P_c > P_e \Rightarrow \left(\frac{1}{P_c}\right) < \left(\frac{1}{P_e}\right)$

$$P_e : \text{expansion ratio} = \left(\frac{V_4}{V_3}\right) = \left(\frac{V_1}{V_3}\right)$$

$$V_1 = V_4$$

$$P_c : \text{compression ratio} = \left(\frac{V_1}{V_2}\right)$$

$$T_1 V_1^{r-1} = T_2 V_2^{r-1}$$

$$T_3 V_3^{r-1} = T_4 V_4^{r-1}$$

$$(T_4 - T_1) V_1^{r-1} =$$

$$T_3 V_3^{r-1} - T_2 V_2^{r-1}$$

$$\Rightarrow \eta = 1 - \frac{1}{r} \cdot \frac{T_3 V_3^{r-1} - T_2 V_2^{r-1}}{V_1^{r-1} (T_3 - T_2)}$$

$$= 1 - \frac{1}{r} \cdot \frac{\left(\frac{T_3}{T_2}\right) V_3^{r-1} - V_2^{r-1}}{(V_1)^{r-1} \left[\left(\frac{T_3}{T_2}\right) - 1\right]}$$

While only p, V equations for easy derivation done on next page

$$= \frac{\frac{V_3^r}{V_2} - V_2^{r-1}}{V_1^{r-1} \left[\frac{V_3}{V_2} - 1\right]}$$

$$= \frac{V_3^r - V_2^r}{V_1^r(V_3 - V_2)}$$

$$= \frac{\left(\frac{V_3}{V_1}\right)^r - \left(\frac{V_2}{V_1}\right)^r}{\left(\frac{V_3}{V_1}\right) - \left(\frac{V_2}{V_1}\right)} = \frac{\left(\frac{1}{P_e}\right)^r - \left(\frac{1}{P_c}\right)^r}{\left(\frac{1}{P_e}\right) - \left(\frac{1}{P_c}\right)}$$

$$\Rightarrow \eta = 1 - \frac{1}{r} \left[\frac{\left(\frac{1}{P_e}\right)^r - \left(\frac{1}{P_c}\right)^r}{\left(\frac{1}{P_e}\right) - \left(\frac{1}{P_c}\right)} \right]$$

$$\frac{\left(\frac{T_4 - T_1}{T_3 - T_2}\right)}{P_4 V_4 - P_1 V_1} = \frac{(P_4 - P_1) V_1}{(V_3 - V_2) P_2}$$

now eliminate
"pressure" terms

$$P_1 V_1^r = P_2 V_2^r$$

$$P_2 = P_3$$

$$P_3 V_3^r = P_4 V_4^r$$

$$V_4 = V_1$$

use PV^r
form

$$= \frac{P_4 V_4 - P_1 V_1}{P_3 V_3 - P_2 V_2}$$

$$= \frac{\left[P_3 \left(\frac{V_3}{V_4}\right)^r - P_2 \left(\frac{V_2}{V_1}\right)^r\right] V_1}{[V_3 - V_2] P_2}$$

$$= \frac{\left(\frac{V_3}{V_4}\right)^r - \left(\frac{V_2}{V_1}\right)^r}{\left(\frac{V_3}{V_1}\right) - \left(\frac{V_2}{V_1}\right)}$$

$$= \frac{\left(\frac{1}{P_e}\right)^r - \left(\frac{1}{P_c}\right)^r}{\left(\frac{1}{P_e}\right) - \left(\frac{1}{P_c}\right)}$$

① Clausius introduced the concept of entropy. He gave a quantitative criteria following which we can predict direction of evolution of a process. He showed that only such processes can occur for which entropy does not decrease.

② Boltzmann showed that entropy is a measure of disorder in molecular arrangement of a system.

③ Since entropy characterizes the approach to equilibrium, it can be related to the probability of occurrence of state. Thus, it helps to relate the results of thermodynamics to those of statistical mechanics.


Refrigerator

Brefrigerator = Coefficient of performance



$$Q_1 = Q_2 + W = \left(\frac{Q_2}{W} \right)$$

Opposite heat and work flows.

$$\eta = \frac{1}{\beta + 1}$$



Entropy



Going from 1 adiabatic to other via any isothermal process $\left(\frac{Q}{T}\right)$ remains const. [like equipotential lines, adiabats are equi-entropic lines]

$$\left(\frac{Q_1}{T_1}\right) = \left(\frac{Q_2}{T_2}\right) = \left(\frac{Q_3}{T_3}\right) = \dots = \left(\frac{Q_n}{T_n}\right)$$

We defined this change as entropy

$$S_2 - S_1 = \left(\frac{Q_1}{T_1}\right)$$

We can take any 2 isotherms A and B.

With the fixed adiabats α and β , they can make a Carnot engine

$$\Rightarrow \frac{Q_1}{Q_2} = \frac{T_A}{T_B} \Rightarrow \left(\frac{Q_1}{T_A}\right) = \left(\frac{Q_2}{T_B}\right)$$

Entropy is a thermodynamic variable. It is mathematically defined as, $ds = \left(\frac{dQ}{T}\right)_S$. It is a state function. It remains constant during reversible adiabatic process.

$$\begin{cases} s = \text{const.} \\ ds = 0 \end{cases}$$

$$\int_1^2 ds = \int_1^2 \frac{dQ_{rev}}{T} \rightarrow \text{going from equilibrium state 1 to equilibrium state 2 via a reversible process.}$$

1st law

We know, internal energy is difference of heat supplied and work produced. This increase in internal energy can be used to do more work via adiabatic expansion.

0th law

It gives the concept of temperature.

2nd law (IP even some processes do not violate conservation of energy, they may violate $\Delta S_{univ} > 0$. Hence not allowed)

It gives us the concept of entropy. Consider following points about Entropy: 2nd law \Rightarrow Carnot theorem \Rightarrow Clausius inequality

- ① Entropy change can be calculated as $\int \frac{dQ_{rev}}{T}$ only for reversible processes.
- ② Entropy change for reversible adiabatic process is 0.
- ③ Note that, it is a state function, we are interested only in ds , not actual S . Hence, this enables us to calculate ds for irreversible processes.

$$ds_{1,2 \text{ irreversible}} = ds_{1,2 \text{ reversible}}$$

Replace actual process by reversible process from i to f states. Calculate $\int \frac{dQ}{T}$

- ④ Entropy is a measure of disorder in the system. All natural processes, that occur spontaneously, create more chaos, more disorder, i.e. increase the entropy: law of increase of Entropy

$$\Delta S > 0$$

⑤ Equilibrium is the most probable state.
 Maximum disorder also means more probability.

Equilibrium \Rightarrow Prob_{maximum} \Rightarrow Maximum Entropy

Concentration gradient
 \Rightarrow mixing will lead to more and equilibrium

Entropy is an additive quantity while Probability is multiplicative \Rightarrow log function is the relation.

$$\Rightarrow S \propto \ln \Omega$$

Ω : thermodynamic probability.

$$S = k \ln \Omega$$

S : entropy

k : boltzmann's const

⑥ From Clausius Inequality, we have seen

$$\oint \frac{dQ}{T} \leq 0$$

$$\int_1^2 \frac{dQ}{T} = \left. dS \right|_1^2 = S_2 - S_1$$


$$\& \oint_1^1 \frac{dQ}{T} \leq 0$$

To prove Entropy is path independent function

Let A, B, C are reversible processes.

$$\int_1^2 dS = \int_1^2 \frac{dQ}{T} = S_2 - S_1$$

$$\oint_{AB} \frac{dQ}{T} = 0 \quad [\text{using Clausius inequality}]$$



$$\Rightarrow \int_A \frac{dQ}{T} + \int_B \frac{dQ}{T} = 0 \quad - \quad (1)$$

$$\oint_{BC} \frac{dQ}{T} = 0 \quad [\text{Clausius inequality}]$$

$$\Rightarrow \int_B \frac{dQ}{T} + \int_C \frac{dQ}{T} = 0 \quad - \quad (2)$$

\rightarrow Note that Clausius inequality talks nothing about dS . It talks about $\oint \frac{dQ}{T}$

From ①

$$\int_B \frac{dQ}{T} = - \int_A \frac{dQ}{T}$$

Carnot theorem



Clausius Inequality
 $\oint \frac{dQ}{T} \leq 0$

From ②

$$\int_B \frac{dQ}{T} = - \int_C \frac{dQ}{T}$$

↓
Concept of Entropy
 $\Delta S \geq 0$

$$\Rightarrow \int_A \frac{dQ}{T} = \int_C \frac{dQ}{T} \Rightarrow \boxed{\text{Path Independent}}$$

Now let us say process C is irreversible.

$$\oint_{AB} \left(\frac{dQ}{T} \right) = 0$$

$$\& \oint_{BC} \left(\frac{dQ}{T} \right) < 0$$

$$\Rightarrow \int_{A'}^2 \frac{dQ}{T} + \int_2^1 \frac{dQ}{T} = 0$$

$$\Rightarrow \int_B^2 \frac{dQ}{T} + \int_1^2 \frac{dQ}{T} < 0$$

$$(S_2 - S_1) + \int_2^1 \frac{dQ}{T} = 0$$

⇒ From ①

$$(S_1 - S_2) + \int_1^2 \frac{dQ}{T} < 0$$

$$\Rightarrow \int_B^1 \frac{dQ}{T} = (S_1 - S_2)$$

This shows that entropy of isolated system cannot decrease. For reversible changes, entropy will remain constant. while for irreversible changes, entropy must increase.

For isolated systems,
 $dQ = 0$
 $\Rightarrow \Delta S \geq 0$

This means that entropy of an isolated system tends towards maximum i.e. equilibrium. while it approaches equilibrium.

$$\Rightarrow S_2 - S_1 \geq \int_1^2 \frac{dQ}{T}$$

$$(S_2 - S_1) > \int_1^2 \frac{dQ}{T} \quad \text{irreversible}$$

$$\text{or } \int_1^2 \frac{dQ}{T} < (S_2 - S_1) \quad \text{irreversible}$$

equality holds good for reversible process.

$$\int_1^2 dS = \int_1^2 \frac{dQ}{T}$$

reversible process

Now dQ can be $C_V dT$ or $C_p dT$ or $m s dT$

let us say $dQ = C_V dT$ { solids
or liquids }

$$\Rightarrow \int_1^2 dS = \int_1^2 C_V \frac{dT}{T} = C_V \ln\left(\frac{T_2}{T_1}\right)$$

~~Let us take ice (1kg) at -20°C . It is to be converted to steam (1kg) at 200°C . Calculate Entropy Change!!~~

1 kg Ice at -20°C (253K) to 0°C (273K)

$$\Delta S_1 = \int dS = \int \frac{dQ}{T} = \int C_V \frac{dT}{T} = C_{V_{\text{ice}}} \ln\left(\frac{273}{253}\right)$$

(assumed ^{all} along reversible process)

1 kg ice to 1 kg water at 0°C : state change

$$\Delta S_2 = \int dS = \int \frac{dQ}{T} = \left(\frac{mL}{273}\right)$$

1 kg of water at 0°C to water at 100°C

$$\Delta S_3 = \int dS = C_{V_{H_2O}} \ln\left(\frac{373}{273}\right)$$

1 kg of water at 100°C to steam at 100°C

$$\Delta S_4 = \left(\frac{mL}{373}\right)$$

1 kg of steam at 100°C to 200°C

$$\Delta S_5 = C_{\text{steam}} \ln\left(\frac{473}{373}\right)$$

$$\Delta S = [\Delta S_1 + \Delta S_2 + \Delta S_3 + \Delta S_4 + \Delta S_5]$$

Entropy is an additive process!!

Note that $\Delta S_1 = k \ln \Omega_1$,

$$\Delta S_2 = k \ln \Omega_2$$

:

$$\Rightarrow \Delta S = \sum_{i=1}^5 \Delta S_i = k \sum_{i=1}^5 \ln \Omega_i = k \ln (\Omega_1 \Omega_2 \Omega_3 \Omega_4 \Omega_5)$$

Entropy change calculations for 1 gram-mole of ideal gas.

$$We have, PV = RT$$

$$PdV + VdP = RdT$$

$$S = f(T, V)$$

We know

$$dS = \frac{dQ}{T} = \frac{dU + PdV}{T} = \frac{C_V dT + PdV}{T}$$

$$= C_V \left(\frac{dT}{T} \right) + R \left(\frac{dV}{V} \right)$$

$$\Rightarrow S = C_V \ln T + R \ln V + C_1$$

or

$$\Delta S = C_V \ln \left(\frac{T_2}{T_1} \right) + R \ln \left(\frac{V_2}{V_1} \right)$$

For isothermal :

$$\Delta S = R \ln \left(\frac{V_2}{V_1} \right)$$

For isochoric :

$$\Delta S = C_V \ln \left(\frac{T_2}{T_1} \right)$$

Also we can write, $S = f_2(T, P)$

$$\begin{aligned} dS &= \frac{C_V dT + P dV}{T} = \frac{C_V dT + R dT - V dP}{T} \\ &= (C_V + R) \left(\frac{dT}{T} \right) - \left(\frac{V}{T} \right) dP \\ &= C_P \left(\frac{dT}{T} \right) - R \left(\frac{dP}{P} \right) \end{aligned}$$

$$\Rightarrow S = C_P \ln T - R \ln P + C_2$$

$$\Delta S = C_P \ln \left(\frac{T_2}{T_1} \right) - R \ln \left(\frac{P_2}{P_1} \right)$$

For isobaric : $\Delta S = C_P \ln \left(\frac{T_2}{T_1} \right)$

Also we can write $S = f_3(P, V)$

$$\begin{aligned} dS &= \frac{C_V dT + P dV}{T} = \frac{\frac{C_V}{R} (P dV + V dP) + P dV}{\frac{PV}{R}} \\ &= \frac{(C_V + R) dV + \frac{C_V}{P} dP}{\frac{PV}{R}} \end{aligned}$$

$$= \frac{(C_V + R) dV}{PV} + \frac{C_V}{P} \left(\frac{dP}{P} \right)$$

$$= \frac{C_P}{V} \left(\frac{dV}{V} \right) + \frac{C_V}{P} \left(\frac{dP}{P} \right)$$

$$\Rightarrow S = C_P \ln V + C_V \ln P + C_3$$

$$\Delta S = C_P \ln \left(\frac{V_2}{V_1} \right) + C_V \ln \left(\frac{P_2}{P_1} \right)$$

For 1 gram-mole gas undergoing :-

1) Isothermal

$$S_2 - S_1 = \underline{R \ln\left(\frac{V_2}{V_1}\right)} = \underline{R \ln\left(\frac{P_1}{P_2}\right)}$$

2) Isobaric

$$S_2 - S_1 = \underline{C_p \ln\left(\frac{T_2}{T_1}\right)} = \underline{C_p \ln\left(\frac{V_2}{V_1}\right)}$$

3) Isochoric

$$S_2 - S_1 = \underline{C_V \ln\left(\frac{T_2}{T_1}\right)} = \underline{C_V \ln\left(\frac{P_2}{P_1}\right)}$$

(4) Adiabatic Process: $S_2 - S_1 = 0$

✓ Illustrate, using 3 different paths, that ΔS is path independent.

Path 1

$$S_2 - S_1 = C_p \ln 2 + C_V \ln\left(\frac{1}{2}\right)$$

$$= (C_p - C_V) \ln 2$$

$$= R \ln 2$$

(rather than any reversible curve, take Path ① as isothermal curve. $\Rightarrow S_2 - S_1 = R \ln\left(\frac{2V}{V}\right) = \underline{R \ln 2}$)


Path 2

$$\Delta S = \Delta S_1 + \Delta S_2$$

$$= C_V \ln\left(\frac{1}{2}\right) + C_p \ln\left(\frac{2}{1}\right)$$

$$= (C_p - C_V) \ln 2$$

$$= R \ln 2$$



Path 3

$$\Delta S = \Delta S_1 + \Delta S_2 = C_p \ln 2 + C_V \ln\left(\frac{1}{2}\right) = (C_p - C_V) \ln 2 = R \ln 2$$

Thermodynamics (4)

29/12/2011

Tut 21

(Q6)

$\Delta S_{\text{resistor}}$

$$\Delta S_{\text{universe}} = \Delta S_{\text{resistor}} + \Delta S_{\text{surrounding}}$$

Resistance remains same. } NO change in state of resistor.
 Temperature remains same. } $\Rightarrow \Delta S_{\text{res}} = 0$

$$\Delta S_{\text{surrounding}} = \left(\frac{I^2 R t}{T} \right) / t = \left[\frac{I^2 R}{T} \right]$$

(Q7)

$$\Delta S = \Delta S_A + \Delta S_B$$

$$Q = 10^4 \text{ J/s}$$

$$= -\frac{Q}{1500} + \frac{Q}{500}$$

$$= Q \cdot \frac{2}{1500} \text{ J/s}$$

$$= 10^4 \cdot \frac{2}{1500} = \frac{200}{15} = \left(\frac{40}{3} \right) \frac{\text{J}}{\text{s}}$$

(Q9)

outside the course but do cover it from some book !!

(Q14)

Carnot's Theorem :- $\eta_R > \eta_{Ia}$ working b/w same T_1 & T_2

(Q20)


Proof

Proof is by contradiction.

let us suppose $\eta_I > \eta_R$

$$\eta_I = \frac{W}{Q_1}, \quad \eta_R = \frac{W}{Q'_1}$$

$$\Rightarrow Q'_1 > Q_1$$



Now reversible engine can be performed in backward direction also.

Let us couple I and R engine with R reversed

Now R acts as refrigerator

2nd law states that its impossible for any device to take heat from lower temperature and transfer it to higher temperature.

2nd law is violated in this system as $[Q_1' - Q_1] > 0$
↑
heat transferred

Hence contradiction

⇒ Our assumption is wrong

$$\Rightarrow \boxed{\eta_R > \eta_I}$$

Q15) $\Delta U = 0$ $\Delta S = R \ln\left(\frac{V_2}{V_1}\right) \Leftarrow$ assumption of ideal gas

$$H = U + PV$$


U is a state function $\Rightarrow H$ has to be a state function

$$dH = d(PV) = P_2 V_2 - P_1 V_1$$

Q17) ice \rightarrow water at 0 $\xrightarrow{\Delta S_1} \xrightarrow{\Delta S_2}$ water at $T \xleftarrow{\Delta S_3}$ water at 30

Tut 20

① Adiabatic Compression




Adiabatic Expansion




Slope: Θ_{re}

④



irrev.



rev

$$\Delta(\text{rev}) > \Delta(\text{irrev})$$


do not attempt such questions

GO FOR STANDARD QUESTIONS !!

⑥ Cooling ; adiabatic demagnetization ; Useful work in engine, travelling of sound waves

⑦ For diatomic gas : $\gamma = 1.4$

⑧ "increases n times".



→ Summing over all cycles of small isothermal-adiabatic curves

$$\frac{dq_1}{T_1} = -\frac{dq_2}{T_2}$$

$$\Rightarrow \oint \frac{dq}{T} = 0$$

Thermodynamic Relationships

1st Thermodynamic Potential : Internal Energy

1st law of Thermodynamics

$$dQ = dU + PdV$$

2nd law

$$dQ = T dS$$

[reversible process]

Combining,

$$TdS = dU + PdV$$

Note that for any P,V,T system $dW = PdV$.

It is not just for ideal gas but for all P,V,T systems i.e. system described by state (P,V,T).

$$\Rightarrow dU = T dS - P dV$$

1 of the 4

Internal Energy is a thermodynamic potential i.e.

Work done per unit quota against the Force.

e.g. $dV = \frac{dW}{m} = \frac{F \cdot de}{m}$ (gravitational)

$$dV = \left(\frac{dW}{q} \right) \quad (\text{electromagnetic})$$

Always create such definitions

Here, thermodynamic Potential = Work done on system in [dU] reversible adiabatic process is stored in form of internal energy.

$$dU = T dS - P dV$$

In adiabatic, $dU = -P dV$ = work done on the system

$$U = U(s, v)$$

$$dU = \left(\frac{\partial U}{\partial s}\right)_v ds + \left(\frac{\partial U}{\partial v}\right)_s dv$$

$$\Rightarrow \left(\frac{\partial U}{\partial s}\right)_v = T \quad \text{Also}$$

$$\left(\frac{\partial s}{\partial U}\right)_v = \frac{1}{T}$$

$$\left(\frac{\partial U}{\partial v}\right)_s = -P$$

~~✓~~ We know if $z = f(x, y)$

$$\left[\frac{\partial}{\partial y} \left(\frac{\partial z}{\partial x} \right)_y \right]_x = \left[\frac{\partial}{\partial x} \left(\frac{\partial z}{\partial y} \right)_x \right]_y \therefore \text{Mathematical definition of Path independence}$$

It is valid if z is state function i.e. path independence

$\Rightarrow dz$ is exact differential

\Rightarrow above relation is valid

$$\left[\frac{\partial}{\partial V} \left(\frac{\partial U}{\partial S} \right)_V \right]_S = \left[\frac{\partial}{\partial S} \left(\frac{\partial U}{\partial V} \right)_S \right]_V$$

$$\Rightarrow \left[\frac{\partial T}{\partial V} \right]_S = - \left[\frac{\partial P}{\partial S} \right]_V$$

1st Maxwell Relationship

Group 1	Group 2
P V	S T

OR $\left(\frac{\partial \alpha}{\partial \beta}\right)_Y : (\bar{\alpha}, \bar{Y})$: same group

belongs to
same group

group of 1 variable vary करे और दूसरे constant रखो !!

$$\left\{ \begin{array}{l} \text{vary } \left(\frac{\partial P}{\partial S}\right)_V \\ \text{const.} \end{array} \right. = - \left(\frac{\partial T}{\partial V}\right)_S$$

2nd Potential

Helmholtz Free Energy

"Free energy" because it can be retrieved when the need arises.

Mathematically defined as $F = U - TS$

U is state function $\Rightarrow F$ is state function
 \Rightarrow only dF is important.

$$dF = dU - TdS - SdT$$

We know $dU = TdS - PdV$

IAPAC recommends symbol "A" for Helmholtz free energy

$$\Rightarrow dF = -PdV - SdT$$

If dT is zero \Rightarrow isothermal process.

ie, In Reversible Isothermal Process, work done on the system is stored in form of Helmholtz Free Energy

$$\left(\frac{\partial F}{\partial V}\right)_T = -P$$

$$\left(\frac{\partial F}{\partial T}\right)_V = -S$$

$$F = f(P, S)$$

F is a state function of V and T
 $\Rightarrow dF$ must be exact differential

$$\Rightarrow \left(\frac{\partial}{\partial T} \left[\frac{\partial F}{\partial V} \right]_T\right)_V = \left(\frac{\partial}{\partial V} \left(\frac{\partial F}{\partial T} \right)_V \right)_T$$

$$\Rightarrow \left(\frac{\partial P}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$$

2nd Maxwell Relation.

3rd Potential Enthalpy or Total Heat Function

Mathematically, $H = U + PV$

U : state function $\Rightarrow H$ is a state function

$$dH = dU + PdV + VdP$$

$$= Tds - PdV + PdV + VdP$$

$$dH = Tds + VdP$$

As C_V corresponds

to U ... on the similar lines

C_P corresponds to H

$$\Rightarrow H = f(S, P)$$

✓ In ^{reversible} Isobaric Process, Heat given is stored as enthalpy

$$dH = Tds + VdP$$

$$= Tds + 0 \quad (\text{Isobaric})$$

$$= dQ \quad (\text{reversible})$$

Components of def'n of H

$$\Rightarrow \left(\frac{\partial Q}{\partial T} \right)_P = \left(\frac{\partial H}{\partial T} \right)_P = C_P$$

$$\left(\frac{\partial H}{\partial S} \right)_P = T$$

$$\left(\frac{\partial H}{\partial P} \right)_S = V$$

- (1) Thermodynamic Potential
- (2) mathematically defined as $H = U + PV$
- (3) heat given to system in reversible isobaric process stored as enthalpy.
- (4) state function
 \Rightarrow total differential gives 3rd Maxwell eqn

H : state function \Rightarrow Path independent \Rightarrow exact differential

$$\Rightarrow \left[\frac{\partial}{\partial P} \left(\frac{\partial H}{\partial S} \right)_P \right]_S = \left[\frac{\partial}{\partial S} \left(\frac{\partial H}{\partial P} \right)_S \right]_P$$

$$\Rightarrow \left(\frac{\partial T}{\partial P} \right)_S = \left(\frac{\partial V}{\partial S} \right)_P$$

3rd Maxwell Relation

4th Potential

Gibbs Free Energy

Mathematically,

$$G = H - TS$$

H: state function \Rightarrow G: state function $\Rightarrow dG$: important

$$\begin{aligned} dG &= dH - TdS - SdT \\ &= TdS + VdP - TdS - SdT \end{aligned}$$

$$dG = \underbrace{VdP}_{\text{Mechanical Eqb.}} - \underbrace{SdT}_{\text{Thermal Eqb.}}$$

✓ At equilibrium $dG = 0$ and minimum.

*importance
in chemistry
if $dG \neq 0$ \Rightarrow chemical reaction is
going on \Rightarrow not yet equilibrium*

i.e. $\frac{\partial G}{\partial P} = 0$
& $\frac{\partial^2 G}{\partial P^2} > 0$

$$\left(\frac{\partial G}{\partial P} \right)_T = V$$

$$\left(\frac{\partial G}{\partial T} \right)_P = -S$$

Gibbs free energy is the maximum amount of non-expansive work that can be extracted from a closed system in a reversible process.

G: state function

$$\left[\frac{\partial}{\partial T} \left(\frac{\partial G}{\partial P} \right)_T \right]_P = \left[\frac{\partial}{\partial P} \left(\frac{\partial G}{\partial T} \right)_P \right]_T$$

$$\Rightarrow \left(\frac{\partial V}{\partial T} \right)_P = - \left(\frac{\partial S}{\partial P} \right)_T$$

4th Maxwell Equation

Refer to End of ~~lecture~~!!

★ $\underbrace{P, V, T, U, S}_{\text{state variables}}$, $\underbrace{F, G, H}_{\text{state functions}}$:- only 2 are independent variables
 state variables dx: state functions
 ↓ ↓
 define a state define process

- ⑤ We subtract the entropy part to get free energies
eg.

$$F = U - TS$$

$$G = H - TS$$

- ⑥ Natural tendency of a spontaneous reaction is to decrease its free energy (Gibbs Free Energy)
 $\Rightarrow \Delta G_{\text{spontaneous reaction}} < 0$

This energy released equals the maximum amount of work that can be performed as a result of the chemical reaction.

In contrast, if ΔG is > 0 \Rightarrow energy (in the form of work) has to be added to the reacting system to make the reaction go.

- ⑦ In Maxwell's relations, $\frac{\partial V}{\partial T}$ is negative i.e. either $\left(\frac{\partial T}{\partial V}\right)_S$ or $\left(\frac{\partial V}{\partial T}\right)_P$ relationships bear a negative sign !!
- ⑧ In Maxwell Relations, variable of LHS is constant of RHS and variable of RHS is const. of LHS.

Thermodynamics (5)

30 / 12 / 11 Yd2d

- All relations derived were of form

$$PV = ST$$

From this relation, we have to do cross 1 by 1.

$$\textcircled{I} \quad \left(\frac{\partial P}{\partial S}\right)_V = -\left(\frac{\partial T}{\partial V}\right)_S \quad : \quad dU = TdS - PdV$$

$$\textcircled{II} \quad \left(\frac{\partial P}{\partial T} \right)_V = \left(\frac{\partial S}{\partial V} \right)_T \quad : \quad dF = -PdV - SdT$$

$$\text{III} \left(\frac{\partial V}{\partial S} \right)_P = \left(\frac{\partial T}{\partial P} \right)_S \quad : \quad dH = TdS + VdP \quad 1.10$$

$$\textcircled{IV} \quad \left(\frac{\partial V}{\partial T} \right)_P = - \left(\frac{\partial S}{\partial P} \right)_T \quad : \quad dG = VdP - SdT \quad 11$$

अनुपात स्वता !! } refer Qds Cdt end of Thermo Notes

- These Equations are very important. Questions like, "Does C_p varies with Pressure ?" or "Does C_V varies with volume ?"

$$\text{i.e. } \textcircled{1} \boxed{\left[\frac{\partial C_V}{\partial V} \right]_T} = 0 \quad ?$$

Solution $C_V = \left(\frac{\partial Q}{\partial T}\right)_V = T \left(\frac{\partial S}{\partial T}\right)_V \leftarrow$ no relation relating S and T as relations are cross in nature

$$\text{LHS} = \frac{\partial}{\partial V} \left[T \left(\frac{\partial S}{\partial T} \right)_V \right]_T = T \frac{\partial}{\partial T} \left[\left(\frac{\partial S}{\partial V} \right)_T \right]_V \quad (\because S \text{ is path independent})$$

$$= T \frac{\partial}{\partial T} \left[\left(\frac{\partial P}{\partial T} \right)_V \right]_V = \boxed{T \left(\frac{\partial^2 P}{\partial T^2} \right)_V}$$

$$\text{For ideal gas} \Rightarrow P = \left(\frac{R}{V}\right) T$$

$$\left(\frac{\partial P}{\partial T}\right)_V = \left(\frac{R}{V}\right)$$

$$\left(\frac{\partial^2 P}{\partial T^2}\right)_V = 0$$

$$\Rightarrow \left(\frac{\partial C_V}{\partial V}\right)_T = 0 \quad \text{for ideal gas}$$

→ It comes out to be zero even for a real gas !!

$$\text{if For a special gas : } PV = A + BT + CT^2$$

$$\Rightarrow \left(\frac{\partial P}{\partial T}\right)_V = \frac{1}{V} [B + 2CT]$$

$$\left(\frac{\partial^2 P}{\partial T^2}\right)_V = \frac{1}{V} [2C]$$

$$\Rightarrow \left(\frac{\partial C_V}{\partial V}\right)_T = \frac{2CT}{V} \neq 0$$

$$\textcircled{2} \quad \boxed{\left(\frac{\partial C_V}{\partial P}\right)_T} = 0 ?$$

$$\text{Solution } C_p = \left(\frac{\partial Q}{\partial T}\right)_P = T \left(\frac{\partial S}{\partial T}\right)_P$$

$$\text{LHS.} = \frac{\partial}{\partial P} \left(T \left(\frac{\partial S}{\partial T}\right)_P \right)_T = T \frac{\partial}{\partial T} \left(\left(\frac{\partial S}{\partial P}\right)_T \right)_P$$

$$= T \frac{\partial}{\partial T} \left(\cancel{\left(\frac{\partial S}{\partial P}\right)} \left(-\frac{\partial V}{\partial T}\right)_P \right)_P = -T \left(\frac{\partial^2 V}{\partial T^2}\right)_P$$

$$= -T \left(\frac{\partial^2 V}{\partial T^2}\right)_P$$

(Q1)

Ideal Gas

$$\left(\frac{\partial U}{\partial V}\right)_T = 0$$

for

$$PV = RT$$

$$\left(\frac{\partial U}{\partial V}\right)_T = f(v) \quad \text{for} \quad \left(P + \frac{a}{v^2}\right)(v - b) = RT$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - P = T \left(\frac{\partial P}{\partial T}\right)_V - P$$

$$\text{for ideal gas} = T \left(\frac{R}{V}\right) - P$$

$$\text{for real gas} \quad \left(\frac{\partial P}{\partial T}\right)_V = \left(\frac{R}{V-b}\right)$$

$$\Rightarrow T \left(\frac{\partial P}{\partial T}\right)_V - P = \frac{TR}{V-b} - P = \left(\frac{a}{V^2}\right)$$

From above relations, we can gather a useful result :-



$$U = U(T, V)$$

[It can be written in form of]
any 2 variables

$$\underline{dU} = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

for a general process

\uparrow
 Energy due to movement of molecules
 C_V

\uparrow
 Energy due to interaction of molecules
 0 for ideal gas

$\left(\frac{a}{V^2}\right)$ for real gas

इन दो में क्या
लिखा, इस पर टिकाए !!

For an isochoric process :

- $dQ = C_V dT$
- $dW = 0$
- $dU = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T \cdot dV$
 $= C_V dT + \left(\frac{\partial U}{\partial V}\right)_T \cdot 0 = C_V dT$

Q7) If It's written, "start from Maxwell Equation"

$$\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V$$

Multiply by T ,

$$T\left(\frac{\partial S}{\partial V}\right)_T = T\left(\frac{\partial P}{\partial T}\right)_V$$

$$\Rightarrow \left(\frac{\partial Q}{\partial V}\right)_T = T\left(\frac{\partial P}{\partial T}\right)_V$$

$$\Rightarrow \left(\frac{\partial U + PV}{\partial V}\right)_T = T\left(\frac{\partial P}{\partial T}\right)_V$$

$$\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T + P = T\left(\frac{\partial P}{\partial T}\right)_V$$

$$\Rightarrow \left(\frac{\partial U}{\partial V}\right)_T = T\left(\frac{\partial P}{\partial T}\right)_V - P$$

How to write in

exam....

if says "starting from
Maxwell's equations, prove
that...."

Coefficient of Elasticity of Gases : Bulk Modulus

Stress : dP Strain : $\left(\frac{dV}{V}\right)$ $k = - \frac{dP}{\left(\frac{dV}{V}\right)}$

$[P \uparrow \Rightarrow V \downarrow]$
but
 k is +ve

$$\Rightarrow k = -V \left(\frac{dP}{dV}\right)$$

So Process can be isothermal or adiabatic

- Isothermal Coefficient of Elasticity

$$k_T = -V \left(\frac{\partial P}{\partial V} \right)_T$$

- Isothermal Coefficient of Compressibility

$$\beta_T = \frac{1}{k_T}$$

Note that these symbols are not standard

- Adiabatic Coefficient of Elasticity

$$k_s = -V \left(\frac{\partial P}{\partial V} \right)_s$$

- Adiabatic Coefficient of Compressibility

$$\beta_s = \frac{1}{k_s}$$

- Volume Coefficient of Expansion of Gases

- We also know volume of gases increase with Temp.

$$V(T) = V_0 (1 + \alpha T) \quad \text{at const. Pressure} \checkmark$$

$$\Rightarrow \alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_P$$

$$Q/ \text{Show that } C_p - C_V = T \left(\frac{\partial P}{\partial T} \right)_V \left(\frac{\partial V}{\partial T} \right)_P \quad (1)$$

$$= -T \left(\frac{\partial P}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P^2 \quad (2)$$

$$= [b + \left(\frac{\partial V}{\partial P} \right)_T] \left(\frac{\partial V}{\partial T} \right)_P \quad (3)$$

$$C_p = T \left(\frac{\partial S}{\partial T} \right)_P$$

$$C_V = T \left(\frac{\partial S}{\partial T} \right)_V$$

$$\text{LHS} = T \left[\left(\frac{\partial S}{\partial T} \right)_P - \left(\frac{\partial S}{\partial T} \right)_V \right]$$

= 

Standard way !!

Let S be expressed as $f(T, V)$

① $\Rightarrow dS = \left(\frac{\partial S}{\partial T} \right)_V dT + \left(\frac{\partial S}{\partial V} \right)_T dV$

प्राप्ति न हो सकती
यद्यकि $\left(\frac{\partial S}{\partial T} \right)_P = \left(\frac{\partial S}{\partial T} \right)_V + \left(\frac{\partial S}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P$

$$\Rightarrow \left(\frac{\partial S}{\partial T} \right)_P - \left(\frac{\partial S}{\partial T} \right)_V = \left(\frac{\partial S}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P$$


$$\Rightarrow \text{LHS} = T \left[\left(\frac{\partial S}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P \right]$$

$$= T \left[\underbrace{\left(\frac{\partial P}{\partial T} \right)_V}_{\text{1st part}} \underbrace{\left(\frac{\partial V}{\partial P} \right)_P}_{\text{2nd part}} \right]$$

③ 2nd part
 $\text{let } U = f(T, V)$
 $\Rightarrow dU = \left(\frac{\partial U}{\partial T} \right)_V dT + \left(\frac{\partial U}{\partial V} \right)_T dV$
 $\Rightarrow TdS - PdV = \left(\frac{\partial U}{\partial T} \right)_V dT + \left(\frac{\partial U}{\partial V} \right)_T dV$
 $\Rightarrow TdS = \left(\frac{\partial U}{\partial T} \right)_V dT + \left(P + \frac{\partial U}{\partial V} \right)_T dV$
 Differentiating w.r.t. T at const P
 $\Rightarrow \left(\frac{\partial Q}{\partial T} \right)_P - \left(\frac{\partial U}{\partial T} \right)_V = \left[P + \left(\frac{\partial U}{\partial V} \right)_T \right] \left(\frac{\partial V}{\partial T} \right)_P$
 $\Rightarrow C_P - C_V = \text{RHS}$

Now if we know equation of state

$(C_P - C_V)$ can be calculated



at A : $\boxed{\left(\frac{\partial V}{\partial T} \right)_P = 0}$

$\Rightarrow C_P = C_V$

$C_P - C_V = - T \left(\frac{\partial b}{\partial v} \right)_T \left(\frac{\partial v}{\partial T} \right)_P^2$
 since $\left(\frac{\partial v}{\partial T} \right)_P > 0$ and $\left(\frac{\partial b}{\partial v} \right)_T < 0$ for all substances $\Rightarrow C_P - C_V > 0$ for all substances.
 At temp = 4°C for water, $\left(\frac{\partial v}{\partial T} \right)_P = 0$
 $\Rightarrow C_P = C_V$

As $T \rightarrow 0$, $C_P \approx C_V$



Q13) Similar 1st Part

$$\left[b + \cancel{a} + \frac{a}{V^2} \right] \left[\cancel{\frac{RT}{(P+a/V^2)}} \right] = R$$

$$C_p = C_v + R$$

$$C_v = \left(\frac{\partial U}{\partial T} \right)_V = C$$

$$\Rightarrow \boxed{C_p = C + R}$$

Using integration by Parts,

$$\left(\frac{\partial V}{\partial T} \right)_P = \frac{R}{\left(\frac{RT}{V-b} \right) [1 - \frac{2a(V-b)}{V^3 RT}]}$$

Using Binomial,

$$\Rightarrow R \left[1 + \frac{2a(V-b)^2}{V^3 RT} \right] + C = C_p$$

$$C_p \boxed{\left(\frac{\partial V}{\partial T} \right)_P = \frac{V-b}{T} \left[1 + \frac{2a}{RTV} \right]}$$

What about formula for real Gas

For Vander Waal's gas

$$\boxed{C_p - C_v = R + \frac{2a}{VT}}$$

$$\Rightarrow R \left[1 + \frac{2a}{VRT} \right] + C = C_p$$

$$\Rightarrow C_p = \cancel{R} + \left(\frac{2a}{VT} \right) + C$$

This result can be more easily obtained via

$$\boxed{C_p - C_v = T \left(\frac{\partial P}{\partial T} \right)_V \left(\frac{\partial V}{\partial T} \right)_P}$$

★ For Vander Waal's Gas, for derivation of $\left(\frac{\partial V}{\partial T} \right)_P$,

perform $+ \frac{a}{V^2} - \frac{a}{V^2}$ in denominator to form $P + \frac{a}{V^2} - \frac{2a}{V^2}$ term and $(RT/V-b)$ as common.

$$C_p - C_v = -T \left(\frac{\partial P}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P^2$$

We need to do "NO DERIVATION" in this answer !!

- (a) Reasoning :
- ① $\left(\frac{\partial P}{\partial V} \right)_T$ cannot be 0
For Water
 - ② $\left(\frac{\partial V}{\partial T} \right)_P = 0$ at $T = 4^\circ C$

(b) For Real Gas

For Ideal Gas



$$-T \left(\frac{R}{V^2} \right) \left(\frac{R}{P} \right)^2 T \\ = \frac{R^3}{R^2 T^2}, T = R$$

Proof

$$C_p - C_v = +T k_T \alpha^2 V \quad (E_T = k_T)$$

$$= -T \left(\frac{\partial V}{\partial T} \right)_P^2 \left(\frac{\partial P}{\partial V} \right)_T \quad (2)$$

$$\left(\frac{\partial P}{\partial T} \right)_V = - \left(\frac{\partial P}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P$$

From

$$\text{If } Z = f(x, y, z) \text{ or } f(x, y, z) = 0$$

$$\left(\frac{\partial z}{\partial x} \right)_y, \left(\frac{\partial z}{\partial y} \right)_x, \left(\frac{\partial z}{\partial z} \right)_x = -1$$

or

From

$$P = f(T, V)$$

$$dP = \left(\frac{\partial P}{\partial V} \right)_T dV + \left(\frac{\partial P}{\partial T} \right)_V dT$$

$$0 = \left(\frac{\partial P}{\partial T} \right)_P = \left(\frac{\partial P}{\partial T} \right)_V + \left(\frac{\partial P}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P$$

$$\Rightarrow \left(\frac{\partial P}{\partial T} \right)_V = - \left(\frac{\partial P}{\partial V} \right)_T \left(\frac{\partial V}{\partial T} \right)_P$$

Q1 In Isoentraltic Process, show that

Change ~~wrt~~ Temp wrt. Pressure is given by

$$\frac{1}{C_p} \left[T \left(\frac{\partial V}{\partial T} \right)_P - V \right]$$

Joule kelvin
Coefficient is tough
i.e. $T P H$ i.e. $\left(\frac{\partial T}{\partial P} \right)_H$

Solution

This is the Joule kelvin Process

To prove $\left(\frac{\partial T}{\partial P} \right)_H = \frac{1}{C_p} \left[T \left(\frac{\partial V}{\partial T} \right)_P - V \right]$

$$\boxed{\frac{1}{C_p} < >}$$

Ans bracket \Rightarrow volume \Rightarrow terms

Joule-kelvin coefficient

$$H = U + PV$$

$$dH = dU + \underbrace{PDV}_{PdV} + VdP$$

$$\Rightarrow dH = dQ + V dP$$

$$\begin{aligned} \text{also } Q &= \left(\frac{\partial Q}{\partial T} \right)_P \\ &= \left(\frac{\partial H}{\partial T} \right)_P \end{aligned}$$

at const. Pressure

$$dH = dQ \quad \Rightarrow$$

We know Cyclic Process

$$\left(\frac{\partial T}{\partial P} \right)_H = - \left(\frac{\partial T}{\partial H} \right)_P \left(\frac{\partial H}{\partial P} \right)_T$$

$$= -\frac{1}{C_p} \left[\left(\frac{\partial H}{\partial P} \right)_T \right]$$

deriving from function
 $H = f(P, T)$

$$\Rightarrow dH = \left(\frac{\partial H}{\partial P} \right)_T dP + \left(\frac{\partial H}{\partial T} \right)_P dT$$

\Rightarrow at const. H & doing $\left(\frac{\partial}{\partial T} \right)_H$

$$0 = \left(\frac{\partial H}{\partial P} \right)_P \left(\frac{\partial P}{\partial T} \right)_H + \left(\frac{\partial H}{\partial T} \right)_P$$

$$\left(\frac{\partial H}{\partial P}\right)_T = \left(\frac{\partial U}{\partial P}\right)_T + \left[\frac{\partial}{\partial P}(PV)\right]_T$$

For ideal gas $\left(\frac{\partial U}{\partial P}\right)_T = 0$

For real gas $\left(\frac{\partial U}{\partial P}\right)_T \neq 0$: deviation from Joule's law

Real gas $\left[\frac{\partial}{\partial P}(PV)\right]_T \neq 0$ deviation from Boyle's Law

Ideal gas $\left[\frac{\partial}{\partial P}(PV)\right]_T = 0$

$$\Rightarrow \mu = \left(\frac{\partial T}{\partial P}\right)_H = 0 \text{ for ideal gases}$$

For non-ideal gas in general

$$\begin{aligned} \mu &= -\frac{1}{C_p} \left[T \left(\frac{\partial S}{\partial P} \right)_T - P \left(\frac{\partial V}{\partial P} \right)_T + V + P \left(\frac{\partial V}{\partial P} \right)_T \right] \\ &= -\frac{1}{C_p} \left[-T \left(\frac{\partial V}{\partial T} \right)_P + V \right] = \frac{1}{C_p} \left[T \left(\frac{\partial V}{\partial T} \right)_P - V \right] \end{aligned}$$

$$dU = T dS - P dV$$

$$\left(\frac{\partial U}{\partial P}\right)_T = T \left(\frac{\partial S}{\partial P}\right)_T - P \left(\frac{\partial V}{\partial P}\right)_T$$

For ideal gas : 0

$$\text{For real gas : } \frac{1}{C_p} \left[\frac{2a}{RT} - b \right]$$

★ [Heat Pump (or refrigerator) is the opposite of]
heat engine

H.W.



Kelvin Planck statement from Clausius

Say Clausius is wrong

⇒ Heat Pump (R) can be there :

We can couple it with Heat Engine (H)

s.t. Heat remove by R be same as
Heat Rejected from H.



Now the combined is a heat engine that converts heat ($Q' - Q$) to work (W) without producing any external effect.


⇒ Violation of Kelvin-Planck statement



Clausius Statement from Kelvin Planck Statement

Say Kelvin-Planck statement is wrong.


Let us have Heat Engine (H) such that




We can couple it with Heat Pump, st.
(R)

Work Produced by H is used by R

such that





$$W = Q \Rightarrow Q'' = W + Q' > W \Rightarrow Q'' > Q$$

Now the combined device transfers heat from lower temp. to higher temperature without using any external work.

Thermodynamic Temperature Scale

We can use Carnot Principle to establish thermodynamic temperature scale. Carnot Theorem states that all reversible engines working between same heat reservoirs are equally efficient. Thus, any reversible heat engine operating between temperature T_1 and T_2 must have same efficiency, i.e. efficiency is a function of only temperatures.

$$\eta = 1 - \frac{q_C}{q_H} \Rightarrow \left(\frac{q_C}{q_H} \right) = f(T_C, T_H)$$

$$\text{let } T_1 > T_2 > T_3$$

$$f(T_1, T_3) = \frac{q_3}{q_1} = \left(\frac{q_3}{q_2} \right) \left(\frac{q_2}{q_1} \right) = f(T_3, T_2) f(T_2, T_1)$$

LHS is not a function of T_2
 \Rightarrow in order to achieve this

$$f(T_1, T_2) = \frac{g(T_2)}{g(T_1)}$$

$$\text{and } f(T_2, T_3) = \frac{g(T_3)}{g(T_2)}$$

$$\text{s.t. } f(T_1, T_3) = \frac{g(T_3)}{g(T_2)} \cdot \frac{g(T_2)}{g(T_1)} = \frac{g(T_3)}{g(T_1)}$$

As a matter of convenience, we choose $g(T) = T$. Now

Choosing 1 fixed reference as reference Temperature (i.e. Triple Point of Water), we establish Thermodynamic Temperature Scale, as the other one [note that we need 2 points to make scale] is Absolute Zero. Can define T .

By international standards, the unit is kelvin and Points are defined as:

- 1) Absolute zero, the lowest possible temperature at 0K and -273.15°C
- 2) Triple Point of water at 273.16 K and 0.01°C

This definition defines 3 things:-

- a) It fixes the magnitude of kelvin Unit as being precisely 1 part in 273.16 parts the difference between Absolute Zero & Triple Point of water.
- b) It establishes that one kelvin has precisely the same magnitude as 1-degree increment on Celsius scale
- c) It establishes difference between 2 scales' null points as being precisely 273.16 K.

$$\Rightarrow \boxed{T(V-b)^{\frac{R}{C_V}} = \text{const.}}$$

$$dU + PdV = 0 \quad \dots \quad (1)$$

$$\begin{aligned} dU &= \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV \\ &= C_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV \end{aligned} \quad (2)$$

$$\begin{aligned} \text{Now } \left(\frac{\partial U}{\partial V}\right)_T dV &= \left(\frac{Tds - PdV}{dV}\right)_T dV = T \left(\frac{\partial s}{\partial V}\right)_T dV \quad \left(\frac{\partial s}{\partial V} \text{ for real gas}\right) \\ &= T \left(\frac{\partial P}{\partial T}\right)_V dV - PdV \end{aligned} \quad (3)$$

$$\begin{aligned} \Rightarrow dU + PdV &= C_V dT + T \left(\frac{\partial P}{\partial T}\right)_V dV - PdV + PdV = 0 \\ \Rightarrow C_V dT + T \left(\frac{\partial P}{\partial T}\right)_V dV &= 0 \quad \Rightarrow C_V dT + \frac{RT}{V-b} dV = 0 \end{aligned}$$

Thermodynamics (6)

02/01/2012

Joule Thomson or Joule - kelvin Coefficient.

$$\mu = \left(\frac{dT}{dP} \right)_H = \frac{1}{C_p} \left[T \left(\frac{\partial V}{\partial T} \right)_p - V \right]$$

For ideal gases : $\mu = 0$ i.e. $dT = 0$

$$\text{For real gases} : \mu = \frac{1}{C_p} \left[\frac{2a}{RT} - b \right]$$

Joule - kelvin Effect

जॉले प्लग में Joule Thomson Effect



High Pressure at low Pressure
जॉले प्लग में गैस रिलीज होती है तो असरी Temperature drop कर जाती है !!

- Outcome of famous Porous Plug Experiment.
- Joule was interested in interaction of real gas molecules.
- In ideal gases, molecules are point particles while in real gases they occupy space, hence interact, while the former do not interact.
- H₂ and He are so small st. it can be treated as an ideal gas.

Joule's law : $U_{\text{ideal gas}} = U(T)$

i.e. $\left(\frac{\partial U}{\partial V} \right)_T = 0$; $\left(\frac{\partial U}{\partial P} \right)_T = 0$



Joule wanted to prove, for real gases $\left(\frac{\partial U}{\partial V} \right)_T \neq 0$ & $\left(\frac{\partial U}{\partial P} \right)_T \neq 0$



- Thomson improved the experimental set up and performed famous porous plug experiment, with Joule. It's also called Joule Thomson Experiment.

Aim: To find out inter-molecular interactions in real gases.

RESULT: "Whenever a real gas is subjected to constant Pressure difference in an irreversible adiabatic process, resulting into change of its temperature, this change in temperature w.r.t. Pressure difference ~~is given by~~ is called Joule-Thomson Coefficient"

$$\mu = \left(\frac{dT}{dP} \right)_H = \frac{1}{C_p} \left[T \left(\frac{\partial V}{\partial T} \right)_P - V \right]$$

Set up :



P₁: Const. High Pressure

P₂: Const. low pressure

⇒ ISOBARIC

T_{gas} = T_{sensitive heat bath}
(initially)

[Most of the gases, except H₂ & He ; T ↓ in Joule-Thomson Experiment]

Walls are totally insulated i.e. no heat flow allowed \Rightarrow Adiabatic

Porous Plug : Cotton or silk

We have to do work to maintain Pressure P_1 & P_2 .



$$\left. \begin{aligned} W_{\text{done by external force}} &= P_1 V_1 - P_2 V_2 \\ \rightarrow W_{\text{gas}} = W_{\text{sys}} &= W = (P_2 V_2 - P_1 V_1) \end{aligned} \right] \star$$

$$dW = P_2 dV_2 - P_1 dV_1$$

$$W = P_2 V_2 - P_1 V_1$$

$$dQ = 0 = dU + dW$$

$$\Rightarrow dW = -dU = U_1 - U_2$$

Irreversible



$$\Rightarrow U_1 - U_2 = P_2 V_2 - P_1 V_1$$

\star Adiabatic + Const. Pressure difference

$$\Rightarrow U_1 + P_1 V_1 = U_2 + P_2 V_2$$

\Rightarrow Isoenthalpic Process

This is an irreversible process

$$dP = P_1 - P_2 = \text{const.}$$

$$\mu = \left(\frac{\partial T}{\partial P} \right)_H = \frac{1}{C_p} \left(T \left(\frac{\partial V}{\partial T} \right)_P - V \right)$$

$$\star \text{Now } \left(\frac{\partial T}{\partial P} \right)_H = -\frac{1}{C_p} \left(\frac{\partial H}{\partial P} \right)_T$$

$$= -\frac{1}{C_p} \left[\left(\frac{\partial U}{\partial P} \right)_T + \left(\frac{\partial (PV)}{\partial P} \right)_T \right]$$

derivation from
Joules law
 $U = f(T)$

derivation
from Boyle's
law
 $(PV)_T = \text{const.}$

$$\text{For real gases, } \mu = \frac{1}{C_p} \left[\frac{2a}{RT} - b \right]$$

Since ideal gas follow
both the laws \Rightarrow

$\mu_{\text{ideal gas}} = 0$
↑
Joule-Kelvin
Coefficient

$$\frac{T_1 - T_2}{P_1 - P_2} = \frac{1}{C_p} \left[\frac{2a}{RT} - b \right]$$

$$T_1 - T_2 = \left(\frac{P_1 - P_2}{C_p} \right) \left[\frac{2a}{RT} - b \right]$$

Cooling

① $T_1 > T_2 \Rightarrow \frac{2a}{RT} - b > 0 \Rightarrow T < \frac{2a}{Rb}$

② T : initial temperature or Temperature of heat bath.

Heating

① $T_2 > T_1 \Rightarrow T > \left(\frac{2a}{Rb} \right)$

at $T = \left(\frac{2a}{Rb} \right)$: no change in Temperature.

It is the Temperature of inversion (T_i)

$$T_i = \left(\frac{2a}{Rb} \right)$$

$T < T_i$: Cooling

$T > T_i$: heating

$T = T_i$: no change in temp.

✓ Note the dP is negative....

① For H_2 and He , a and b are very small.

T_i for H_2 and He is very small.

$\Rightarrow T > T_i$: always heating

Other gases : $T_i \approx 1000^\circ K$

$\Rightarrow T < T_i$: always cooling

Classius-Clapeyron Equation

- ① (3 Proof)
 - ① 1st order: 2 applications
 - ② 2nd order: 2 examples

- ② It describes 1st order Phase Transition

e.g. Melting

Freezing

Vapourization

Condensation


3rd CT
Joule-Thomson
($\frac{\partial P}{\partial T}$)

i.e. $\Delta S \neq 0, \Delta V \neq 0$

★ On a Pressure-Temp. (P-T) diagram, the line separating the 2 phases is known as the Coexistence curve. This relation gives the slope of

this curve.

$$\frac{dP}{dT} = \left(\frac{mL}{T \Delta V} \right)$$



- ③ If C_p changes \Rightarrow 2nd order Phase Transition

- ④ Gibbs Function

$$dG = VdP - SdT$$

$$\text{If } \left(\frac{\partial G}{\partial P} \right)_T = V, \quad \left(\frac{\partial G}{\partial T} \right)_P = -S \quad \text{i.e. } 1^{\text{st}} \text{ derivatives}$$

are discontinuous \Rightarrow 1st Order Phase Transition.

★ discontinuous means that during change, both volume & entropy undergo a change. e.g. melting, vapourization since $dT=0, [C_p, \alpha, k_T] \rightarrow \infty$
If n^{th} derivative is discontinuous $\Rightarrow n^{\text{th}}$ order Phase Transition

e.g. $\left(\frac{\partial^2 G}{\partial P^2} \right)_T = \left(\frac{\partial V}{\partial P} \right)_T$: if its discontinuous \Rightarrow 2nd order Phase Change

$\left(\frac{\partial^2 G}{\partial T^2} \right)_P = -\left(\frac{\partial S}{\partial T} \right)_P = -\frac{C_p}{T}$: if C_p is discontinuous \Rightarrow 2nd order Phase Change

Note that here, volume & entropy are usually continuous (remain same)
Note that 2nd order if any 1 is discontinuous or both.


- ⑤ Note that we always talk about

Solid \rightarrow liquid transition

and

liquid \rightarrow vapour transition



$$\frac{\partial \ln P}{\partial \frac{1}{T}} = \frac{C_p}{V} : \quad \text{--- ---}$$



e.g. Ferromagnetic Material $\xrightarrow{\text{heating above Curie Temp.}}$ Paramagnetic Material

its 2nd order ϕ -change

Normal Conductor $\xrightarrow[\text{certain Temp.}]{\text{Cooled below}}$ Super Conductor



Note that slope of
liq-gas conversion is
 $\frac{\partial v}{\partial p} \text{ i.e. } v_{\text{vap}} > v_{\text{liq}}$
but
slope of solid-liq may
be ore or ore .
It's ore for H_2O .

Classius Clapeyron from Maxwell

⑥ Classius-Clapeyron Equation deals with change of Phase Change Temperature upon application upon Pressure

$$\frac{dp}{dT} = \frac{L}{T(V_2 - V_1)}$$

$$V: \text{specific volume} = \frac{1}{f}$$

Phase 1 $\xrightarrow{\text{at a}}$ Phase 2
 V_1 • Temp T V_2
 • Pressure P
 • $\Delta Q = L$

→ It is one of the most important formula of thermodynamics. It gives rate at which vapour pressure must change with Temperature for 2 phases to coexist in equilibrium.

Inversely, we may investigate effect of Pressure on Boiling Point

We know $\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V$ [IInd Maxwell equation]

Multiply by T, $T \left(\frac{\partial S}{\partial V}\right)_T = T \left(\frac{\partial P}{\partial T}\right)_V$

$$\Rightarrow \left(\frac{\partial Q}{\partial V}\right)_T = T \left(\frac{\partial P}{\partial T}\right)_V$$

dQ · heat given to system... तथा जप्त की ओर जाएगा !!

$$\Rightarrow \frac{L}{T(V_2 - V_1)} = \left(\frac{dP}{dT} \right)$$

(4)

liquid \rightarrow vapour

(6 cm of Hg)

dT

$$= \frac{2 \cdot 3 \times 10^6}{373 \left(\frac{1}{P_2} - \frac{1}{P_1} \right)}$$

for ex., For H_2O $(V_2 - V_1) < 0$

melting of ice

$\Rightarrow LHS$ is \ominus ve

If I increase Pressure,
T melting can be reduced i.e.

even at $-3^\circ C$ ice can be melted
Upon application of enough

Pressure: REGELATION

④ Note that V and L are for 1 kg $\Rightarrow V = \frac{L}{P}$

or if m kg and corresponding volumes


$$\Rightarrow \frac{mL}{T(V_2 - V_1)}$$

$$\Rightarrow \left(\frac{\partial b}{\partial T} \right)_{\text{saturation}} = \frac{L}{T(V_{\text{vap}} - V_{\text{liq}})}$$

Since $V_{\text{vap}} > V_{\text{liq}}$ $\Rightarrow \left(\frac{\partial b}{\partial T} \right)_{\text{sat.}} > 0$

\Rightarrow increase in Pressure raises the boiling point, and vice versa. Hence the concept of Pressure Cooker.

Clasius Clapeyron from Carnot Cycle



PV Curve for Carnot Cycle

$\rightarrow P$ and T are const.


$\rightarrow V$ changes

AB, CD: isotherms
BC, DA: adiabats

Quick I change the Pressure,
corresponding T changes quickly
 \Rightarrow adiabatic process.

Let ABCD be a reversible Carnot Cycle.

Ph 1: liquid
Ph 2: vapour




I quickly reduce the P
i.e. adiabatic expansion

$$\frac{Q_1}{Q_2} = \frac{T_1}{T_2}$$

$$\frac{L+dL}{L} = \frac{T+dT}{T}$$

$$\Rightarrow \frac{dL}{L} = \frac{dT}{T}$$

$$\Rightarrow dL = \left(\frac{L}{T}\right) dT$$



$$\text{Work Done in Carnot Cycle} = \frac{Q_1 - Q_2}{\cancel{L}}$$

$$= L + dL - L$$

$$= dL$$

$$= \frac{L}{T} dT$$

$$\text{Work Done} = \text{Area of Curve} \approx dP (V_2 - V_1)$$

$$\Rightarrow \frac{L}{T} dT = dP (V_2 - V_1)$$

$$\Rightarrow \frac{\frac{L}{T} dT}{T(V_2 - V_1)} = \frac{dP}{dT}$$

Hence, Proved

★ In case of vapourization, $V_2 > V_1$ in all cases $\Rightarrow \frac{dP}{dT} > 0$

\Rightarrow if $dP \uparrow \Rightarrow dT \uparrow$ and if $P \downarrow \Rightarrow T \downarrow$

In case of melting, $V_2 > V_1$ or $V_1 > V_2$.

e.g. ^{ice to water}: $V_2 < V_1$

$$\Rightarrow (V_2 - V_1) < 0 \Rightarrow \left(\frac{dP}{dT}\right) < 0$$

Similarly

ice, gallium, bismuth
 $V_2 < V_1$ **BIG**

\Rightarrow if $P \uparrow \Rightarrow T \downarrow$

if $P \downarrow \Rightarrow T \uparrow$

e.g. Wax on melting : $V_2 > V_1 \Rightarrow V_2 - V_1 > 0 \Rightarrow \left(\frac{dP}{dT}\right) > 0$

Sulphur

\Rightarrow if $P \uparrow \Rightarrow T \uparrow$

$P \downarrow \Rightarrow T \downarrow$

~~Non Adiabatic~~

$A \rightarrow B : L + dL$

$B \rightarrow C : -C_2 dT$: saturated vapour \rightarrow normal vapour

$C \rightarrow A : -L$

$D \rightarrow A$: normal liquid \rightarrow saturated liquid

$C_1 dT$

$$\Rightarrow dQ = (L + dL) - (C_2 dT) - (L) + (C_1 dT)$$

$$dQ = dL + (C_1 - C_2) dT$$

$$\frac{L}{T} dT = \frac{dL}{T} + (C_1 - C_2) dT$$

$$\Rightarrow C_1 - C_2 = \left(\frac{dL}{dT}\right) - \left(\frac{L}{T}\right)$$

$$\Rightarrow \frac{dQ}{dT} = \frac{L}{T} + C_1 - C_2$$

PTO


* For phase change, we draw PT curves

Also the slope is $\left(\frac{dP}{dT}\right)$ for calausius clapeyron equation.....

Clasius Clapeyron
from Gibbs

HW

Energy



$$g_1 > g_2$$

PT Curve
always
for phase change

At both a and b, phases are in equilibrium, so the specific Gibbs free energy of 2 phases at each point are same i.e.

If dg is Δg between a and b

$$\begin{matrix} g_1^{(a)} \\ g_2^{(a)} \end{matrix} = \begin{matrix} g_1^{(b)} \\ g_2^{(b)} \end{matrix}$$

$$\Rightarrow dg_1^{ab} = dg_2^{ab}$$

$$\Rightarrow -S_1 dT + V_1 dP = -S_2 dT + V_2 dP$$

$$\Rightarrow (S_2 - S_1) dT = (V_2 - V_1) dP$$

$$\Rightarrow \frac{dP}{dT} = \frac{(S_2 - S_1)}{(V_2 - V_1)} = \frac{L}{T(V_2 - V_1)}$$

do this
question
from Tut Q1

[refer end of notes]

From IVth law, $\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial V}{\partial T}\right)_P \Rightarrow \left(\frac{\partial Q}{\partial P}\right)_T = -TV \alpha$

Hence for substances that expand on heating $\Rightarrow \alpha > 0$

$$\Rightarrow \left(\frac{\partial Q}{\partial P}\right)_T < 0$$

i.e. Heat is released upon compression

Relation between Specific Heats of Vapour and liquid

(not related to Clausius Clapeyron eqn) HW

For a change in state from liquid to vapour at Temperature T , we have

$$[S_s - S_p] = \left(\frac{L}{T} \right) - \textcircled{1}$$

s: saturated vapour
l: liquid

differentiating $\textcircled{1}$ wrt T ,

$$\frac{dS_s}{dT} - \frac{dS_p}{dT} = \frac{1}{T} \frac{dL}{dT} + L \left(-\frac{1}{T^2} \right)$$

$$\Rightarrow T \cdot \frac{dS_s}{dT} - T \frac{dS_p}{dT} = \frac{dL}{dT} - \frac{L}{T} \quad [\text{multiplying by } T]$$

$$\Rightarrow [C_s - C_p] = \left(\frac{dL}{dT} \right) - \frac{L}{T}$$

Specific heat of saturated ~~vapour~~ vapour is negative, meaning $C_s < 0$, meaning that in order to raise the temperature, a certain quantity of heat is to be withdrawn from the body.

To change saturated vapour at one Temperature to saturated vapour at a higher temp, Pressure must be increased. If compression were effected adiabatically, i.e., without adding or subtracting heat, the vapour would become superheated, consequently, during compression, some heat must be abstracted in order to keep the vapour in saturated condition. Hence $T \uparrow$ is accompanied by removal of heat $\Rightarrow C < 0$.

★ Vapour $\begin{cases} \text{saturated} \\ \downarrow \text{Adiabatic compression} \\ \text{Superheated} \end{cases}$

Thermodynamics (7)

03/01/2012

Adiabatic Demagnetization of Paramagnetic Salt

- Magnetic Properties of a material:
 - ✓ Permeability μ
 - ✓ Susceptibility χ

[Corresponding electrical properties]

→ Permittivity

→ Electrical Susceptibility

$$\mu = \mu_0 [1 + \chi] = \mu_0 \mu_s$$

$\left(\frac{\mu}{\mu_0}\right)$: relative permeability : μ_s

- 3 vectors of Magnetism:

Effects of \vec{H}

\downarrow

\vec{B} , \vec{H} , \vec{M}

Basic or Natural

\vec{H} : Magnetizing force that is applied on a [material that can be magnetized] or [magnetic material]. Magnetic Material has some property of acquiring magnetism e.g. magnetic dipole via unpaired charge like unpaired electrons.

If these dipoles are aligned in direction of field \Rightarrow Magnetism

\vec{M} : no. of magnetic dipoles per unit volume

Fraction of \vec{H} that has caused M is χ

i.e. $\chi \vec{H} = \vec{M}$

\vec{B} or intensity of magnetic induction is the magnetic induced due to applied \vec{H}

i.e. $\vec{B} = \mu_0 \vec{H}$

In free space: $\vec{B} = \mu_0 \vec{H}$

In a material, some dipoles will align too

$$\begin{aligned}\Rightarrow \vec{B} &= \mu_0 \vec{H} + \mu_0 \vec{M} \\ &= \mu_0 (1+\chi) \vec{H} \\ &= \mu \vec{H}\end{aligned}$$

$$\Rightarrow \boxed{\mu = \mu_0 (1+\chi)}.$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ SI units}$$


On these properties, we can classify materials:

(1) DIAMAGNETIC:

- χ is small negative quantity

$$\Rightarrow \mu < \mu_0$$

- We know $\vec{M} = \chi \vec{H}$
if χ is negative \Rightarrow dipoles align weakly opposite to applied \vec{H} .



χ diamagnetic is independent of temperature.

- It happens in materials with paired electrons.
- Superconductors are perfect diamagnetic materials

④ χ is independent of Temperature for Diamagnetic materials

2) Paramagnetic Materials

- small +ve value of χ
- $\mu > \mu_0$
- weakly aligned dipoles in direction of \vec{H}
- dependent on Temperature as inversely proportional

st. $\chi = \frac{M}{H} \propto \left(\frac{1}{T}\right)$

$$\chi = \frac{C}{T} \quad C: \text{Curie's const.}$$

i.e.
$$\vec{M} = \frac{CH}{T} \vec{H}$$

3) Ferromagnetic Materials

- Large +ve value of χ
- $\mu \gg \mu_0$
- $\chi = \left(\frac{C}{T-\theta} \right)$

Valid For $T > \theta$: Paramagnetic Material

Note that for $T < \theta$ it is Ferromagnetic Material

θ : Curie's Temperature

Experiment

$$S = S(T, H)$$

Temp magnetic field applied

just remember
the starting
point

Below θ , ferromagnets has
spontaneous magnetization.
No law governs its.

$$dS = \left(\frac{\partial S}{\partial T} \right)_H dT + \left(\frac{\partial S}{\partial H} \right)_T dH$$

Multiply by T,

$$TdS = T \left(\frac{\partial S}{\partial T} \right)_H dT + T \left(\frac{\partial S}{\partial H} \right)_T dH$$

$$\Rightarrow TdS = \left(\frac{\partial Q}{\partial T} \right)_H dT + T \left(\frac{\partial S}{\partial H} \right)_T dH$$


[like Gases $dW = PdV$] [like $\left(\frac{\partial S}{\partial P} \right)_T = - \left(\frac{\partial V}{\partial T} \right)_P$, we have
here, $dW = -H dM$]

Work done by external field
on system = $H dM$
 \Rightarrow Work done by system = $-H dM$

$$\left(\frac{\partial S}{\partial H} \right)_T = + \left(\frac{\partial M}{\partial T} \right)_H$$

$$\Rightarrow TdS = \left(\frac{\partial Q}{\partial T} \right)_H dT + T \left(\frac{\partial M}{\partial T} \right)_H dH$$

C_H : specific heat at constant \vec{H}



$$\Rightarrow TdS = C_H dT + T \left(\frac{\partial M}{\partial T} \right)_H dH$$

Note that H is here applied magnetic field


For paramagnetic salt, $M = \frac{C_H}{T}$ and not enthalpy

$$\left(\frac{\partial M}{\partial T} \right)_H = - \frac{C_H}{T^2}$$


for paramagnetic

$$\Rightarrow TdS = C_H dT - \left(\frac{C_H}{T} \right) dH$$

\rightarrow Adiabatic demagnetization is accompanied by fall in Temperature. It's called Magneto Caloric Effect.



Characteristic Curve of Paramagnetic Material



Specimen is at Temp. of liquid He initially

$$T_0 = T_{\text{Helium}}$$

There is no scale to determine Temp. below -200°C

NO WAY TO MEASURE ABOUT ABSOLUTE ZERO ??

We will evolve a magnetic Temperature Scale.

$$\chi_1 = \left(\frac{C}{T_1} \right) : \quad \chi_1, T_1 : \text{known}$$

[assuming Curie's law is valid]

$$\chi_2 = \left(\frac{C}{T_2} \right) \quad \chi_2 : \text{known}$$

$$T_2 : ?$$

$$\frac{\chi_1}{\chi_2} = \frac{T_2}{T_1}$$

$$\Rightarrow T_2 = T_1 \left(\frac{\chi_1}{\chi_2} \right)$$



To measure χ : We put a coil in inner path and measure inductance.

Step 1 Isothermal Magnetization

- Let Temperature of liquid Helium be T_1
- Magnetic Field H is applied.
- Some dipoles will be aligned.
- Work Done = $\left[\int H dM \right]$ is stored in form of Potential Energy

Step 2 Adiabatic Demagnetization

- Quickly switch off the field.
- Internal Energy is used to bring back dipoles



$$TdS = C_H dT - \frac{CH}{T} dH$$

Adiabatic : $dS = 0$

$$\Rightarrow \int_{T_1}^{T_2} T C_H dT = + \left. \frac{CH}{2} \right|_H^0$$

$$\Rightarrow T_2^2 - T_1^2 = - \frac{CH^2}{C_H}$$

$$\Rightarrow (T_2 - T_1)(T_1 + T_2) = - \frac{CH^2}{C_H}$$

AB

$$\begin{aligned} dQ &< 0 \\ dU &= 0 \\ dW &< 0 \\ dH_{ext} &> 0 \end{aligned}$$

BC


$$\begin{aligned} dQ &= 0 \\ dW &> 0 \\ dU &< 0 \end{aligned}$$

$$\Rightarrow T_2 - T_1 = -\left(\frac{C}{C_H}\right) \frac{H^2}{2T}$$

★ $\Delta T \propto \frac{1}{T}$: At high Temp, ΔT will be insignificant. ∴ first cool by normal methods.
Make T low... then apply Magneto-Caloric effect
 $H=0$

• Issues of dimensions in S.I. units, but in CGS units, $\mu_0 = 1$ and is dimensionless. Hence this question is always solved in CGS units

(H: oersted)



"When we apply Magnetic Field, ~~dipole moment is reduced to align~~ dipoles are aligned in accordance.

~~This is due to Potential Energy~~ Note that entropy is reduced in this process. \Rightarrow Heat released.

In adiabatic demagnetization, the internal energy is used to de-align the dipoles in absence of H .

Hence the temperature drops".

"Temperature is measured by assuming the validity of Curie's law".

Disadvantages

- ① Cooling capacity is small and one cannot cool any large sized material.
- ② One cannot perform experiments that require absence of magnetic fields:

③ T already should be low.

④ Can we achieve $T=0$? During isothermal magnetization, heat is removed which continuously needs to be removed from salt, for which at surrounding, an even lower T is reqd., which becomes increasingly difficult.

Walls Van der Waals Equation of State for Real Gas

Thermometry:

For 1 mole of an ideal gas,

① Boyle's Law : $P \propto \frac{1}{V}$ ($T = \text{const.}$)

② Charles law : $V \propto T$ ($P = \text{const.}$)

Combining the ① and ②,

$$\boxed{PV = RT}$$

Kinetic Theory:

$$\frac{1}{2}mv^2 = \frac{3}{2}kT$$

$$PV = \frac{1}{3}NmV^2$$

$$= \frac{1}{3}N\cancel{mV^2} = \frac{1}{3}N3kT$$

$$= NKT$$

$$\boxed{PV = RT}$$


$$\boxed{R = Nk}$$

Particles themselves

○ Ideal Gas do not occupy space \Rightarrow No interaction

But real gas interact !!

Correction for Real Gas



① Finite size of Molecules:

Since volume of real gas particles $\neq 0$

\Rightarrow Space for their movement or interaction is reduced

$$\Rightarrow V_{\text{real}} = \underbrace{(V - b)}_{\text{available}}$$

b: Correction for volume occupied by molecules

② Correction for Intermolecular attraction.

$$\left(\frac{\partial U}{\partial V}\right) = \frac{\partial U}{A \cdot \partial r} = \frac{F}{A} = P$$

Vander Wall observed :

$$\frac{\partial U}{\partial V} \propto \frac{1}{V^2}$$

[Actually he observed :]

$$\left(\frac{F(r)}{A}\right) = \frac{A}{r^6} - \frac{B}{r^{12}}$$

* drop in pressure caused due to intermolecular interactions in the outermost layer

$$Hence Considered : \boxed{\frac{\partial U}{\partial V} = \frac{a}{V^2}}$$

$$= \left(\frac{a}{V^2}\right)$$

\Rightarrow Pressure Correction : ~~$P = P + \frac{a}{V^2}$~~ \Rightarrow in the ideal gas, we must replace p by the sum of observed pressure for a real gas and drop caused by intermolecular forces. i.e. $(P + \frac{a}{V^2})$

$$\boxed{\left[P + \frac{a}{V^2}\right] [V - b] = RT}$$

here V is for 1 mole of gas

$$\text{For } n \text{ moles, Put } V = \left(\frac{V}{n}\right)$$

$$\text{Real Gas} \quad \left(P + \frac{an^2}{V^2}\right) \left(\left(\frac{V}{n}\right) - b\right) = RT$$

If V_0 is the volume for n moles
 \Rightarrow for 1 mole
 Put $V = \left(\frac{V_0}{n}\right)$

$$\text{Ideal Gas} \quad P \left(\frac{V}{n}\right) = RT$$

① Van der Waal is a cubic in V.

$$\text{i.e. } V^3 - bV^2 + \frac{aV}{P} - \frac{ab}{P} - \frac{RT}{P} V^2 = 0$$

$$= V^3 - \left[b + \frac{RT}{P} \right] V^2 + \frac{a}{P} V - \frac{ab}{P} = 0$$

Liquification of Gases




Important Curve

Andrew's Experiment on CO_2 liquification

Andrew carried very exhaustive experiments on compressibility P of gases while attempting to liquify them. He found that

- (a) Above a certain temperature (48°C) CO_2 behaves like a perfect gas.
- (b) At 31.4°C , a kink is observed which suggests that gas can be liquified under compression.



Application of P : $\xrightarrow{(a)} A \rightarrow B$: CO_2 behaves as a gas $B \rightarrow C$: liquid and vapour coexist

P_c : Critical Pressure

T_c : Critical Temperature

$C \rightarrow D$: liquid

$\xrightarrow{(b)}$ in liquid phase slope is very high.

For (VAPOUR) $T < T_c$: Gas can be liquified via application of Pressure

(GAS) $T > T_c$: Gas can't be liquified just by application of Pressure

If $P = P_c \quad \left\{ \begin{array}{l} \\ T = T_c \end{array} \right. \Rightarrow V = V_c$

For CO_2 : $T_c : 31.4^\circ\text{C}$
 $P_c : 73 \text{ atm}$

Liquid state में दबाव से ज़रूरी कम्प्रेस करने के लिए अकेले उपयोगी Pressure apply करना पड़ता है। \therefore Very High slope of PV curve in liquid state.

$$V_c^3 - \left(b + \frac{RT_c}{P_c} \right) V^2 + \frac{a}{P_c} V_c - \frac{ab}{P_c} = 0 \quad \text{--- (1)}$$

Since V_c satisfies $V \Rightarrow V - V_c = 0$
or $(V - V_c)^3 = 0$

$$\Rightarrow V^3 - V_c^3 - 3VV_c(V - V_c) = 0$$

$$\Rightarrow V^3 - 3V_c V^2 + 3V_c^2 V - V_c^3 = 0 \quad \text{--- (2)}$$

(1) and (2) are equivalent

→ Temperature below which it becomes possible to liquefy gas under compression is known as Critical temp.

$$\Rightarrow 3V_c = b + \frac{RT_c}{P_c} \quad \text{--- (3)}$$

$$3V_c^2 = \frac{a}{P_c} \quad \text{--- (4)}$$

$$V_c^3 = \frac{ab}{P_c} \quad \text{--- (5)}$$

→ Pressure just required to liquefy a gas at critical temperature is called Critical pressure. and corresponding volume (for 1 mole of a gas) is called Critical volume

Dividing (4) and (5)

$$\frac{3}{V_c} = \frac{1}{b} \Rightarrow V_c = 3b$$

$$b = \frac{V_c}{3}$$

Putting in (4)

$$27b^3 = \frac{ab}{P_c}$$

$$\Rightarrow P_c = \frac{a}{27b^2}$$

$$a = 3P_c V_c^2$$

Putting in (3)

$$8b = \frac{R T_c}{a} 27b^2 \Rightarrow$$

$$T_c = \frac{8ab}{27Rb}$$

Critical Coefficient

$$= \frac{RT_c}{P_c V_c}$$

$$= \left(\frac{8}{3}\right)$$

Reduced Equation of State

$$\alpha = \frac{P}{P_c}$$

$$\beta = \frac{V}{V_c}$$

$$\gamma = \frac{T}{T_c}$$

$$\left(\frac{\partial P}{\partial V}\right)_T = -\frac{RT}{(V-b)^2} + \frac{2a}{V^3}$$

$$\left(\frac{\partial^2 P}{\partial V^2}\right)_T = \frac{2RT}{(V-b)^3} - \frac{6a}{V^4}$$

At critical point.

Also note that for critical isotherm the point P is ~~maxima~~ ~~minima~~ inflection point. $\Rightarrow \left(\frac{\partial P}{\partial V}\right)_T = 0$ and $\left(\frac{\partial^2 P}{\partial V^2}\right)_T = 0$

This offers 2nd way to find out critical parameters. The two equations on Van der Waals gas give:

$$P = \frac{RT}{V-b} - \frac{a}{V^2}$$

$$\frac{\partial P}{\partial V} = \frac{RT}{(V-b)^2} - \frac{2a}{V^3} = 0$$

$$\Rightarrow V_c = 3b \quad \& \quad T_c = \frac{8a}{27bR}$$

Replacing in equation we get P_c

$$\left(P + \frac{a}{V^2}\right) (V - b) = RT \rightarrow \text{we can get same results by placing } P, V, T \text{ in}$$

$$\left(\alpha P_c + \frac{3P_c V_c^2}{\beta^2 V_c^2}\right) \left(\beta V_c - \frac{V_c}{3}\right) = \gamma R T_c \text{ terms of } \alpha, \beta, \gamma \text{ and } a, b.$$

$$\Rightarrow \left[\alpha P_c + \frac{3P_c}{\beta^2}\right] \left[\beta V_c - \frac{V_c}{3}\right] = \gamma R T_c$$

$$\Rightarrow (P_c V_c) \left(\alpha + \frac{3}{\beta^2}\right) \left(\beta - \frac{1}{3}\right) = \gamma R T_c$$


$$\Rightarrow \left(\alpha + \frac{3}{\beta^2}\right) \left(\beta - \frac{1}{3}\right) = \frac{8}{3} \gamma$$

$$P_c V_c = \frac{3}{8} R T_c$$

Validity of derivation of critical constants

Vander Waal's Equation is cubic in V . This means that for each pair of (P, T) , V will have three values either all of them real or 1 real and 2 complex. For low temperatures, 3 real roots exist. As $T \uparrow$, these roots approach one another. They become equal at $T = T_c$. Above $T > T_c$, only 1 real root exists for all P i.e. there can be no condensation in this region.

$$\frac{4}{27} \frac{b^2}{\beta^2} \cdot \frac{35}{R} \frac{27 R b}{8a} = \left(\frac{3}{8}\right)$$



law of Corresponding States

For gas, : $[\alpha_1, \beta_1, r_1]$

For gas_2 : $[a_2, \beta_2, r_2]$

Only 2 Parameters are independent. 3rd can be derived from those 2.

says, if $\alpha_1 = \alpha_2$
 $\&$ $\Rightarrow r_1 = r_2$

And the 2 gases are said to be in corresponding states.

$$\text{eg. Gas 1 : } \begin{bmatrix} P_c : 2 \text{ atm} \\ V_c : 5 \text{ litre} \\ T_c : 300 \text{ K} \end{bmatrix} @ \begin{array}{l} P = 1 \text{ atm} \\ V = 25 \text{ litre} \\ T = 4500 \text{ K} \end{array} \quad \alpha_1 = \frac{1}{2} \quad \beta_1 = 5 \quad \gamma_1 = 15$$

$$\text{Gas 2: } \left[\begin{array}{l} P_c = 5 \text{ atm} \\ V_c = 30 \text{ litre} \\ T_c = 800 \text{ K} \end{array} \right] @ \begin{array}{l} P = 2.5 \text{ atm} \\ V = 15 \text{ litre} \\ T = 1200 \text{ K} \end{array} \quad \alpha_1 = \frac{1}{2} \quad \beta_1 = 5 \quad \gamma = 15$$

\Rightarrow Gas 1 and Gas 2 are in corresponding states.

→ Generic definition of corresponding states is that their reduced parameters are same !!

Q1 Calculate critical temperature of He if $P_c = 2.26 \text{ atm}$.
and critical density is 0.069 g cm^{-3}

Ans Critical volume is defined for 1 mole of gas.

$$\Rightarrow V_c = \frac{\text{Mass of 1 mole}}{\text{density}} = \frac{4g}{0.069 \text{ g/cm}^3} = \underline{\underline{57.97 \text{ cm}^3}}$$

$$b = \frac{V_c}{3} = \underline{\underline{19.32 \times 10^{-6} \text{ m}^3}}$$

$$\Rightarrow \frac{RT_c}{P_c V_c} = \frac{8}{3}$$

$$\frac{a}{27b^2} = 2.26 \times 10^5 \text{ N/m}^2$$

$$\Rightarrow T_c = \underline{\underline{4.2 \text{ K}}}$$

$$\begin{aligned}\Rightarrow T_c &= \frac{8a}{27Rb} = \frac{8}{R} \left(\frac{a}{27b^2} \right) b = \frac{8b}{R} P_c \\ &= \frac{8 \times 19.32 \times 10^{-6} \times 2.26 \times 10^5}{8.31} = \underline{\underline{4.2 \text{ K}}}\end{aligned}$$

Ans b is mathematically equivalent to 4 times the volume occupied by all the molecules

$$\text{i.e. } b = 4 \times \left(\frac{4}{3} \pi r^3 \right) \times N_A$$

$$\text{eg. } b = 2.42 \times 10^{-5} \text{ m}^3/\text{mole}$$

$$\Rightarrow b = 4 \times \frac{4}{3} \times \pi \times \frac{d^3}{8} \times N_A$$

$$\Rightarrow d^3 = \frac{3}{2} \frac{b}{\pi N_A} = 1.92 \times 10^{-29} \text{ m}^3$$

$$\Rightarrow \boxed{d = 2.68 \text{ \AA}}$$

Thermodynamics (8)

04/1/2012

Gibbs Phase Rule & Chemical Potential

$$dU = T dS - P dV$$

$$dF = -P dV - S dT$$

$$dH = T dS + V dP$$

$$dG = V dP - S dT$$



All these relations are valid for only HOMOGENOUS systems.
i.e. single composition system.

↑
(Chemical composition)

e.g. Ice + Water + Steam : Homogenous Multi-Phase System

Ice + (water + sulphur) + Steam : Heterogenous or
Multicomponent Multiphase

For HETEROGENOUS systems,

$$F = U - TS + \mu N$$

↑ no. of molecules
characteristic
of
added component called
Chemical Potential

$$dF = -P dV - S dT + \mu dN$$

Similarly

$$dU = T dS - P dV + \mu_i dN_i$$

$$dF = -P dV - S dT + \mu_i dN_i$$

$$dH = T dS + V dP + \mu_i dN_i$$

$$dG = V dP - S dT + \mu_i dN_i$$

eg.

$$dU = T dS - P dV + \mu_1 dN_1 + \mu_2 dN_2$$

$$\mu_1 = \left(\frac{\partial U}{\partial N_1} \right)_{S, V, N_2}$$

$$\text{or } \mu_i = \left(\frac{\partial U}{\partial N_i} \right)_{S, V, N_j} \text{ read this}$$

Similarly $\mu_i = \left(\frac{\partial F}{\partial N_i} \right)_{V, T, N_j}$

$$\mu_i = \left(\frac{\partial H}{\partial N_i} \right)_{S, P, N_j}$$

$$\mu_i = \left(\frac{\partial G}{\partial N_i} \right)_{P, T, N_j}$$

- All these definitions are of **chemical Potential**. But in applications, we use

$$\boxed{\mu_i = \left(\frac{\partial G}{\partial N_i} \right)_{P, T, N_j}}$$

P, T are constant \rightarrow Thermodynamic Equilibrium.

Gibbs Phase Rule

Coexistence

It describes thermodynamic equilibrium conditions for multi-component multi-phase systems.

'c' components
(coexisting in equilibrium)

'p' phases (coexisting in equilibrium)

$p \in [$ solid
liquid
gas
Plasma
Bose Einstein Condensate $]$

- Degree of Freedom at equilibrium

$$F = C - P + 2$$

^{independent}

[i.e. F is the number of ^{independent} thermodynamic coordinates required to specify the state of thermodynamic equilibrium (variables)]

given that equilibrium is not influence by forces of gravitational, electrostatic, & surface tension forces.
magnetic

We know,

$$F = \text{(Total number of thermodynamic variables)} \\ - \text{(constraints)}$$

① For triple point of water,

$$F = 1 - 3 + 2 = 0$$

i.e. everything is fixed. It is a unique point.

It's called 'INVARIANT SYSTEM'.

→ For ice-water state,

$$F = 1 - 2 + 2 = 1$$

Volume is changed in the process.

It is required to specify the system.

It's called 'MONOVARIANT SYSTEM'

★ Refer [PTO] for Proof of Gibbs Phase Rule
[to last page]

G describes the equilibrium conditions

$$G = H - TS + \mu'_1 N_1 + \mu'_2 N_2 + \dots + \mu'_c N_c \quad 1: \text{specifies solid}$$
$$+ \mu^2_1 N_1 + \mu^2_2 N_2 + \dots + \mu^2_c N_c$$
$$\vdots$$
$$+ \mu^b_1 N_1 + \mu^b_2 N_2 + \dots + \mu^b_c N_c \quad b: b^{\text{th}} \text{ phase}$$

$$dG = V dP - S dT + \sum_{i=1}^c \sum_{j=1}^b \mu_i^{(j)} dN_i^{(j)}$$

At equilibrium,

$$dG = 0$$

$$dP = 0$$

$$dT = 0$$

$$\Rightarrow \sum_{i=1}^c \sum_{j=1}^b \mu_i^{(j)} dN_i^{(j)} = 0$$

let us consider, c component system existing in single phase :

$$G = H - TS + \mu'_1 N_1 + \mu'_2 N_2 + \dots + \mu'_c N_c$$

$$dG = V dP - S dT + \mu'_1 dN_1 + \mu'_2 dN_2 + \dots + \mu'_c dN_c$$

Equilibrium of c components in 1 phase

$$dP = 0 \quad dT = 0 \quad \underbrace{\sum \mu_i^{(1)} dN_1 + \mu_2^{(1)} dN_2 + \dots}_{= 0}$$
$$\Rightarrow \text{independent: } (c-1)$$

\Rightarrow For 'p' phase : independent = $p(c-1)$

Apart from that $\frac{dP}{dT} = 0$

$$\frac{dT}{dP} = 0$$

$\Rightarrow P, T$ are required

Similarly considering 1 component in p phases :

$$dG = 0 \quad dP = 0 \quad dT = 0$$


$$\sum \left[\begin{array}{l} \mu_1^{(1)} dN_1 \\ \mu_1^{(2)} dN_1 \\ \mu_1 dN_1 \\ \vdots \\ \mu_1^{(p)} dN_1 \end{array} \right] = 0 \quad \Rightarrow (p-1) \text{ constraints}$$

$$F = 2 + (c-1)p - (p-1)c$$

$$= 2 - p + c$$

$$F = 2 + c - p$$

★ For triple point, $F=0$ $(2+1-3=0)$
Hence $2+c-p$ only !!



Membrane opened and particles allowed to interact

$$N = N_1 + N_2$$

$$dN = dN_1 + dN_2$$

$$\Rightarrow dN_1 = -dN_2$$

$$dG = V dP - S dT + \mu_1 dN_1 + \mu_2 dN_2$$

at equilibrium

$$(\mu_1 - \mu_2) dN_1 = 0 \quad \leftarrow$$

dynamic equilibrium, dN_1 will not be zero at any instant. On the avg. dN_1 will be zero....

$$\Rightarrow \mu_1 = \mu_2 \quad (\text{i.e. chemical equilibrium})$$

[PPT to PDF] for Transport Phenomenon

Transport Phenomenon

It is used to describe

Transports are measured per unit area per unit time per gradient.

Viscosity of Gases : Transport of Momentum
(to achieve mechanical eqb.)

Thermal Conductivity of Gas : Transport of Heat Energy
(to achieve Thermal eqb.)

Diffusion of Gas : Transport of Mass
(to achieve chemical eqb.)

Heat Flow :

$$\frac{Q}{t} = k A \left(\frac{dT}{dz} \right)$$

$$k = \frac{(Q/t)}{A \cdot \left(\frac{dT}{dz} \right)}$$

Coefficient of
Conductivity

Temperature
Gradient

Viscous Flow :

$$\frac{(F/A)}{(\frac{dv}{dz})} = \boxed{\eta} \quad \text{Viscosity}$$

$$= \boxed{\frac{(\frac{dp}{dt})}{A \cdot \left(\frac{dv}{dz}\right)}} \quad \text{velocity gradient}$$

Diffusion


$$D = \frac{(\text{No. of Particles crossing per unit area per unit time})}{\text{Concentration gradient}}$$

$$= \frac{(\frac{dN}{dt})}{A (\frac{dn}{dz})}$$

Viscosity of Gases as transport of Momentum

velocity
←
velocity gradient
 $z=0$
 λ

[note that velocity and velocity gradient are perpendicular]



λ : mean free path

→ There is relative motion between layers,

Δp when single molecule crosses upwards or downwards = $2 \lambda m \left(\frac{dv}{dz}\right)$

$n = \left(\frac{N}{V}\right)$: molecules per unit volume is known.

⑥ No. of molecules per unit area per unit time

$$\frac{N}{A \Delta t} = \frac{N}{A} dx \left(\frac{dx}{dt} \right) = \frac{N}{V} \cdot \left(\frac{d\vec{v}}{dt} \right) = n \vec{v}$$

In 1 particular direction, no. of particles crossing per unit area per unit time,

$$\frac{n \vec{v}}{3}$$

This is a statistical result that can be proved by calculating the avg. distance d below the

No. of particles crossing only upwards = $\frac{n \vec{v}}{6}$ reference plane where molecule

= No. of particles crossing only downwards makes its last collision It comes out to be per unit area per unit time either $\frac{(2)}{3}$. MENTION THIS

Change in Momentum, when a layer crosses upwards or

$$\text{downwards} = \left(\frac{n \vec{v}}{6} \right) * 2 \lambda m \left(\frac{dv}{dz} \right)$$

$$= \frac{1}{3} mn \bar{v} \lambda \left(\frac{dv}{dz} \right)$$

→ $\boxed{\left[\frac{(F)}{A} \right]} = \frac{1}{3} mn \bar{v} \lambda = \eta$

Tangential stress per unit $\boxed{\left(\frac{dv}{dz} \right)} = \frac{1}{3} \rho \bar{v} \lambda$

velocity gradient $\boxed{\therefore \frac{mN}{V} = \rho}$

η is dependent only on temperature. However at very high Pressure, λ becomes equal to molecular diameter & at very low Pressure, λ becomes comparable to dimensions of apparatus. So, η is $f(P)$ only at very high & very low pressure

$$\lambda = \frac{1}{\sqrt{2} \pi d^2 n}$$

d : diameter of molecule

For ideal gas : $PV = NkT \Rightarrow P = nkT$

$$\Rightarrow \eta = \frac{1}{3} \frac{m \bar{v}^2}{\sqrt{2} \pi d^2}$$


(*) $\bar{v} = \sqrt{\frac{8 k T}{m \pi}}$

\bar{v} : avg velocity

For gases : $n \propto T$
 For liquids : $n \propto \frac{1}{T}$

theoretically, not mathematically.

Thermal Conductivity



Heat flow

→ Heat flow and Temp. Gradient are Parallel.

$$E + \lambda \left(\frac{dE}{dz} \right)$$

$$T + \lambda \frac{dT}{dz}$$

$$E = C_v T$$

$$T$$

$$E - \lambda \left(\frac{dE}{dz} \right)$$

$$T - \lambda \frac{dT}{dz}$$

For 1 molecule crossing : $\Delta Q = 2 \lambda \left(\frac{dE}{dz} \right)$

For whole layer : $\frac{1}{6} n \bar{v} * 2 \lambda \frac{dE}{dz}$

writing
in
this fashion
will be
better

Net flow of heat from higher Temperature region to lower Temperature region per unit area per unit time = $\frac{1}{3} n \bar{v} \lambda \left(\frac{dE}{dz} \right)$

$$\begin{aligned} dQ &= m C_v \Delta T \\ &= m C_v 2 \lambda \frac{dT}{dz} \end{aligned}$$

$$E = m C_v T$$

$$\left(\frac{dE}{dz} \right) = m C_v \left(\frac{dT}{dz} \right)$$

$$\Rightarrow k = \frac{\left(\frac{dQ}{dt} \right)}{A \left(\frac{dT}{dz} \right)}$$

$$\left(\frac{dQ}{dt} \right) = \frac{1}{3} n \bar{v} \lambda m C_v \left(\frac{dT}{dz} \right)$$

$$= 2 m C_v \lambda \frac{n \bar{v}}{6}$$

$$= \frac{1}{3} m n \bar{v} \lambda C_v$$

$$=$$

$$\frac{\left(\frac{dQ}{dt} \right)}{A \left(\frac{dT}{dz} \right)} = \frac{F m n \bar{v} \lambda C_v}{3} = k$$

If give molar heat capacity C

$$C = \frac{(C)}{M}$$

We get $k = C_v n$. However actually due to variation of molecular density with Temperature gradient.

$$k = \epsilon \eta C_v \quad \text{where } 1 \ll \epsilon \ll 2.5$$

\uparrow flow of molecules

n^* represent concentration i.e. $\frac{(N)}{V}$

Diffusion

Conc. gradient \uparrow




$$n^* + \lambda \left(\frac{dn^*}{dz} \right)$$

New Flow of molecules

where n^* is the no. of particle

in reference layer travelling in same direction per unit time.



$$n^*$$



$$n^* - \lambda \left(\frac{dn^*}{dz} \right)$$

No. of particles crossing from higher conc. region to lower conc. region per unit area per unit time = $\frac{n^-}{6}$

$$\frac{\text{no. of molecules}}{6}$$

This is molecules per unit volume

$$D = \frac{1}{3} \bar{v} \lambda = \frac{\left(\frac{dN^+}{dt} \right)}{A \left(\frac{dn^*}{dz} \right)}$$

$$= \left(\frac{m}{P} \right)$$

$$\textcircled{O} \text{ Frequency of Collision} = \left(\frac{\bar{v}}{\lambda} \right) = \left(\frac{1}{T_{\text{Collision}}} \right)$$

$$\textcircled{O} \text{ Density } \rho = mn \quad \underline{m}: \text{ mass of 1 molecule}$$

We know numbers of particles crossing per unit area per unit time = $\frac{n\bar{v}}{6}$

$$\Delta N \text{ from upper surface} = \left(n + \frac{dn}{dz} \cdot \lambda \right) \frac{\bar{v}}{6}$$


$$\Delta N \text{ from lower surface} = \left(n - \frac{dn}{dz} \cdot \lambda \right) \frac{\bar{v}}{6}$$

$$\Rightarrow \Delta N_{\text{net}} = 2 \frac{dn}{dz} \cdot \lambda \frac{\bar{v}}{6}$$

per unit area per unit time

$$\Rightarrow \boxed{\left[\frac{\left(\frac{\Delta N}{\Delta t} \right)}{\Delta A \cdot \left(\frac{dn}{dz} \right)} \right] = \frac{1}{3} \bar{v} \lambda = D}$$

Whenever in a gas, stored in a container, there is concentration gradient, energy gradient etc. some phenomenon, singly or jointly come into play to destroy the cause of non-uniformity this phenomenon is called Transport Phenomenon.



- If particles are distinguishable, a specification of the energy state of each particle is said to define a microstate

Theermodynamics (9)

05/01/2012

STATISTICAL THERMODYNAMICS

Thermodynamic state is described by thermodynamic variables. These variables are measured at equilibrium. To measure, one way is thermometry or calorimetry. 2nd way is kinetic theory : to track down motion of every particle. 3rd way is statistical Physics.

Statistics

"Statistic" implies Probability distribution. (NOT STATISTICS)

① Thermodynamic Probability = ' Ω ' = No. of accessible Microstates to a macrostate.

This is definition of Thermodynamic Probability. Do not confuse it with $0 < p < 1$ Probability of a Microstate may be $(\frac{1}{\Omega})$

② Entropy $S = k \ln \Omega$ (bridge between statistics & Thermo)

Equilibrium \Rightarrow Maximum $\Omega \Rightarrow$ Maximum S

S : additive Ω : ~~Multiplication~~

$$S = S_1 + S_2 \quad \Omega = \Omega_1 \cdot \Omega_2$$

\Rightarrow we chose $S \propto \ln \Omega$


$$\Rightarrow \boxed{S = k \ln \Omega} \quad k: \text{Boltzmann Const.}$$

③ Whether the particles are distinguishable or non distinguishable, a specification of the number of particles n_i in each energy level is said to define a macrostate of the assembly.



In thermodynamics, we encounter two types of distributions:

Statistical Distribution



Energy

Energy emitted or absorbed is continuous.

Nature of Particle

Valid for distinguishable particle.

(Concept of Spin came in 1925)

Hence, spin is not the criteria.

No. of Particles per cell

No Restriction

Quantum Statistics (1926 - 30)

e.g. ① Bose - Einstein (1926)

② Fermi - Dirac (1928)

Energy

Energy is quantized taking discrete values

Nature of Particles

Valid for Indistinguishable particle.

Spin

Here, spin is a criteria

Full Integral :- Bose Einstein
Spin

Half Integral :- Fermi Dirac

Spin

No. of Particles per Cell

- No restriction for Bose Einstein
- Restrictions imposed on Fermi Dirac via Pauli's Exclusion Principle.

★ Here cell means an energy level. n particles in a cell means n -fold energy level degeneracy. But there can be g cells for energy level $E \rightarrow g$ level of degeneracy....

Occupation Index or No. of Particles per Cell of Energy E_i at equilibrium Temp

or [Probability distribution function
at Equilibrium temperature 'T']

$n_{(E_i)}$: no. of particles ~~with~~ energy level E_i

$g_{(E_i)}$: no of cells with energy level E_i

$E_i \rightarrow f_i \quad f_i \quad f_i \quad f_i \quad f_i \dots g_{\text{cells}}$

$$\left(\frac{n_{E_i}}{g_{E_i}} \right) = \left(\frac{n_i}{g_i} \right) = f(E_i)$$

: OCCUPATION INDEX

eg. 90 :	2	2	2
	2,2,1	1,2,2	2,1,2
$g=3$	$f=2$	$n=6$	

In Maxwell Boltzmann,

$$f(E_i) = A e^{-\beta E_i}$$

$$\beta = \frac{1}{kT}$$

$$f(E_i) = e^{-(\alpha + \beta E_i)}$$

where

$$e^{-\alpha} = A$$

$$\Rightarrow n_i = A g_i e^{-\left(\frac{E_i}{kT}\right)}$$

In Bose-Einstein,

$$f(E_i) = \frac{1}{e^{(\alpha + \beta E_i)} - 1}$$

$$\beta = \frac{1}{kT} \quad \alpha = \left(\frac{\mu}{kT} \right)$$

(μ : chemical Potential)

of BOSONS (integral spin)

For Photons: $\mu_{\text{Photon}} = 0$

$$\beta = \frac{1}{kT} \quad \alpha = \left(\frac{E_f}{kT} \right)$$

(E_f : fermi energy of FERMIONS)

(half integral spin)

In Fermi Dirac,

$$f(E_i) = \frac{1}{e^{(\alpha + \beta E_i)} + 1}$$

⑥ Power of 'e' is always a pure number.

Exponent

MB

BE

FD

$$\binom{g_i}{n_i}$$

$$(e^{\alpha + \beta E_i})$$

$$(e^{\alpha + \beta E_i} - 1)$$

$$(e^{\alpha + \beta E_i} + 1)$$


$$e^{\alpha + \beta E_i}$$

$$\binom{g_i}{n_i}$$

$$\binom{g_i}{n_i} + 1$$

$$\binom{g_i}{n_i} - 1$$

→ If $\binom{g_i}{n_i} \approx e^{\alpha + \beta E_i}$ is large, both Bose Einstein and Fermi Dirac will approach to statistical Thermodynamics i.e. indistinguishable particles have become distinguishable, when no. of cells are much more than the particles that have to be placed in the cells. [Concept of unique slot number]



→ Hence, Quantum statistics are required when $\binom{g_i}{n_i}$ is small. It happens when Temperatures are small.


In Ordinary temperatures, $\binom{g_i}{n_i}$ is large, hence Classical statistics is sufficient.

⇒ Basic criteria for distinguishability : $\left(\frac{g_i}{n_i}\right) \leftarrow$ seats or cells
 $\left(\frac{g_i}{n_i}\right) \leftarrow$ students or particles

⇒ If $g_i > n_i \Rightarrow$ Assign unique cell to each particle
 ⇒ distinguishable

g_i : No. of cells with energy E_i

- 0 Temp \Rightarrow No motion
 - 0 is unattainable \Rightarrow Particles are always in motion
 - ① $\sum n_i = N = \text{const.}$ (Total no. of particles)
 - ② $\sum E_i n_i = E = \text{const.}$
- } at equilibrium Temperature 'T'



$$\Omega(N, V, E) = ??$$

We have, 4 Assumptions :-

$$① \sum n_i = N \Rightarrow \sum dn_i = 0$$

$$② \sum G_i n_i = E \Rightarrow \sum E_i dn_i = 0$$

$$③ \text{Entropy is maximum} \Rightarrow dS = 0 \\ \Rightarrow d(\ln \Omega) = 0$$

④ Assumption of a priori equal probability

} at Equilibrium

④ Assumption of Apriori Probability :

Before occupancy, all cells are equiprobable to be occupied.

- ① Macro state : 1 particular value of $\{n_1, n_2, \dots, n_i\}$

e.g. $\{4, 10, 5\}$ is 1 macrostate

$\{10, 4, 5\}$ is other macrostate.


(कौसली number की specify किया दूँ, परमाणु में 4 ... कौन से चार, वह नहीं कहता कि it won't matter in indistinguishable Particles but imb in distinguishables)

- o Micro States : within 1 macro state, there can be multiple micro states depending upon filling of n_i in g_i (all g_i 's do)

(all g_i 's are distinguishable)

$$\begin{bmatrix} E_1 \\ n_1 \\ g_1 \end{bmatrix}, \begin{bmatrix} E_2 \\ n_2 \\ g_2 \end{bmatrix}, \dots, \begin{bmatrix} E_i \\ n_i \\ g_i \end{bmatrix}$$

$$\begin{array}{ll}
 N_5 = 2 & g_5 = 3 \\
 N_4 = 1 & g_4 = 1 \\
 N_3 = 1 & g_3 = 2 \\
 N_2 = 2 & g_2 = 4 \\
 N_1 = 1 & g_1 = 1
 \end{array}$$



ways of filling n_i into g_i

$$\Omega = \prod_i \overbrace{(n_i, g_i)}^{\sim}$$

For 1 particle,

We will measure

x and p_x
 y and p_y
 z and p_z

$$E_x = \frac{px^2}{2m} = \frac{1}{2}mv_x^2$$

$$Ey = \frac{by^2}{2m} = \frac{1}{2} m v y^2$$

$$E_2 = \frac{p_z^2}{2m} = \frac{1}{2} m v_z^2$$

$$1 \times 3 \times 2 = 6$$

In general,

1 x f x 2

Phase Space

• deviation of $n(E)$ $\delta(E)$

- ① To specify dynamical configuration of any assembly of particles, we need 6 coordinates viz x, y, z, p_x, p_y, p_z

Canonical Conjugate Variable: Measurement of 1 affects other
 $\Delta x \Delta p_x$ depends on Heisenberg Principle.

say f : degrees of freedom = 3

\Rightarrow Dimension of Phase space = $3 \times 2 = 6$

Volume Element in Phase Space : Cell

[general 2f dimensions of phase space]

$$\Delta x \Delta y \Delta z \Delta p_x \Delta p_y \Delta p_z$$

$$\text{If continuous, } dx dy dz dp_x dp_y dp_z = d^3x d^3p : \text{Cell}$$

Cells represents mathematical space between

$$x \text{ and } (x + dx)$$

$$y \text{ and } (y + dy)$$

$$z \text{ and } (z + dz)$$

$$p_x \text{ and } (p_x + dp_x)$$

$$p_y \text{ and } (p_y + dp_y)$$

$$p_z \text{ and } (p_z + dp_z)$$

⊗ the ... is smallest volume element

$$= \Delta x \Delta y \Delta z \Delta p_x \Delta p_y \Delta p_z$$

$$= \Delta x \Delta p_x \Delta y \Delta p_y \Delta z \Delta p_z$$

since $\Delta x \Delta p_x \approx h$

$$= h^3$$

- ② In general, volume element in phase space,
 $(dx dy dz \dots dz_f) (dp_x dp_y dp_z \dots dp_f)$

$$\text{If } dx dp_x = h \Rightarrow \boxed{\text{Volume Element} = h^f}$$

- ✓ In classical statistics, \hbar can be any small const.
It can be as small as possible \Rightarrow no. restriction on no. of cells
 \Rightarrow distinguishable.
- ✗ In quantum statistics, \hbar is Planck's const.
 \Rightarrow Restricted number of cells. \Rightarrow indistinguishable particles if more particles.

For 3-dimensional space,

$$\text{No. of cells with energy } E_i = \left[\frac{b^2 x^2 + b^2 y^2 + b^2 z^2}{2m} \right]$$

$$= g(E_i) \quad \text{Volume in phase space with energy } E_i \\ \text{Volume of 1 cell}$$

$$= \frac{\int_{\text{Phase Space}} d^3x d^3p}{h^3} = \frac{V}{h^3} \iiint d\beta_x d\beta_y d\beta_z$$

$$\text{No. of cells with momenta b/w } p \text{ & } (p+dp) = g(p) dp$$

$$= g(p) dp : \text{Representation Only !!}$$

We know, a sphere:

$$b^2 x^2 + b^2 y^2 + b^2 z^2 = p^2$$

$$(p_x + dp_x)^2 + (p_y + dp_y)^2 + (p_z + dp_z)^2 = (p + dp)^2$$

$$dp_x dp_y dp_z = \underbrace{\frac{4\pi}{3} [(p + dp)^3 - p^3]}_{4\pi p^2 dp}$$

\nearrow Both represent space b/w
 $(p_x, p_x + dp_x), (p_y, p_y + dp_y) \rightarrow (p_z, p_z + dp_z)$

$$g(p) dp = \int d^3x \, d^3p$$

$$\boxed{g(p) dp = \frac{V}{h^3} \cdot 4\pi p^2 \, dp} \quad \textcircled{*}$$

~~Integration~~
~~g(p)~~ ~~V~~ ~~h^3~~

similarly,

$g(v) dv$: no. of cells with velocity between ~~v~~ and $(v+dv)$

$g(E) dE$: no. of cells with energy between E and $(E+dE)$

$$\boxed{E = \frac{p^2}{2m}}$$

$$\Rightarrow \boxed{p = \sqrt{2m} \sqrt{E}}$$

$$\& \boxed{dp = \sqrt{2m} \frac{1}{2} E^{-\frac{1}{2}} dE}$$

[for non-relativistic energy]

we have,

$$g(E) dE = \frac{V}{h^3} \cdot 4\pi \cdot 2mE \sqrt{2m} \frac{1}{2} E^{-\frac{1}{2}} dE$$

$$\Rightarrow \boxed{g(E) dE = 2 \frac{V}{h^3} \cdot (2m)^{\frac{3}{2}} E^{\frac{1}{2}} dE} \quad \textcircled{*}$$

$$\Rightarrow \boxed{g(E) dE = C \sqrt{E} dE}$$

~~g(E)~~ ~~dE~~

Density of state

$$\rightarrow n(E) = \int g(E) dE$$

No of cells/states
↓
 $d n(E)$ {
Note that density of state has no relation with number of particles.

$$\rightarrow g(E) = \boxed{\frac{dn(E)}{dE}}$$

We know

$$g(p) dp = \frac{V}{h^3} 4\pi p^2 dp$$

In relativistic case,

$$E = \sqrt{p^2 c^2 + m_0^2 c^4}$$

$$E^2 - m_0^2 c^4 = p^2 c^2$$

$$p^2 = \frac{E^2}{c^2} - m_0^2 c^2$$

Photon For $m_0 = 0$, $p^2 = \frac{E^2}{c^2} \Rightarrow p = \frac{E}{c} \Rightarrow dp = \frac{dE}{c}$


$$g(E) dE = \frac{V}{h^3} \cdot 4\pi \cdot \frac{E^2}{c^3} dE = \underline{\underline{\frac{4\pi V}{c^3 h^3} E^2 dE}}$$

$$g(\gamma) dy = \frac{V}{h^3} \cdot 4\pi \cdot \frac{h^2 \gamma^2}{c^2} \frac{1}{c} dy \quad p = \frac{h\gamma}{c}$$

$$f(\gamma) dy = \boxed{\frac{4\pi V \gamma^2}{c^3} dy}$$

Taking into account the polarization,

$$g(y) dy = \frac{8\pi V y^2 dy}{c^3}$$



Thermodynamics (10)

Conjunction !! [किंतु फलस्त है]

① $\Omega(N, V, E)$: Ensemble

(Plural of Assembly i.e. Assembly
of Assemblies)

$$\textcircled{1} \sum n_i = N \quad \textcircled{2} \sum E_i n_i = E \quad \textcircled{3} \sum d(\ln \Omega) = 0$$

$$\sum dn_i = 0 \quad \sum E_i dn_i = 0$$

④ A priori Probability equal.

$$\textcircled{5} g(E_i) dE_i = \frac{2\pi V}{h^3} (2m)^{3/2} E^{1/2} dE$$

No. of cells b/w energy E_i and $(E_i + dE)$

$$\int_0^E g(E) dE = \underline{n(E)} \leftarrow \text{energy cells upto energy } E$$


nu. of cells / density of state / Degree of degeneracy.

$$\textcircled{6} g(p) dp = \frac{d^3x d^3p}{h^3} = \frac{V}{h^3} d^3p = \frac{V}{h^3} 4\pi p^2 dp$$

⑦ Ω : Thermodynamic Probability

= NO. of accessible microstates to a macrostate

From $E = E_1$ upto $E = E_2$,
the cells are of energy E ,
and are called g_i




* Problem is to put

- (n_1 particles in g_1 cells) &
- (n_2 particles in g_2 cells) &
- (n_3 particles in g_3 cells) ...

$$= \prod_i (n_i, g_i)$$

To put n_i particles in g_i cells



Example

$$n_i = 2$$


$$g_i = 2$$

① decide distinguishable/indistinguishable $g_i = 2$


A B

② decide whether there is any restriction


no restriction



Case 1 : distinguishable & no restriction : MB



Case 2 : indistinguishable & no restriction : BE



Case 3 : indistinguishable & restriction : F.D.

(n_1, n_2, \dots, n_c)

Macrostate
 \updownarrow
Configuration

Maxwell Boltzmann.

Macrostate = (3, 5, 2)

Probability of this Macrostate

$$= N C_{n_1} \frac{N-n_1}{m_1} C_{n_2} \frac{N-n_1-n_2}{m_2} C_{n_3} \dots$$


$N = 10$ Balls of different colours.
 It's a 2 step problem.....

$i = 3$ Tim in box of different numbers.

$$= \frac{\underline{N}}{\underline{m_1} \underline{N-n_1}} \frac{\underline{N-n_1}}{\underline{m_2} \underline{N-n_1-n_2}} \frac{\underline{N-n_1-n_2}}{\underline{m_3} \underline{N-n_1-n_2-n_3}}$$

$$= \frac{\underline{N}}{\underline{m_1} \underline{m_2} \underline{m_3} \dots \underline{m_i}}$$

No. of ways of putting n_i balls in i^{th} box
 $= \frac{110}{13 \ 15 \ 12} = 2520$



Probability of Microstates to this Macrostate

$$= (g_1)^{n_1} (g_2)^{n_2} \dots$$

Now dividing the balls in each division = ~~2520~~ $(15)^3 (25)^5 (27)^2$

Hence,

$$\Omega = \underline{N} \prod_i \frac{\underline{m_i}}{(g_i)^{n_i}}$$

★ Hence total no. of ways = Ω
 $= 2520 (15)^3 (25)^5 (27)^2$
 = Total no. of MICROSTATES TO MACROSTATE (3, 5, 2)

$$\ln \Omega = \sum_i \ln \underline{N} - \ln \underline{m_i} + n_i \ln g_i$$

$$d(\ln \Omega) = \sum_i d(\ln \underline{N}) - d(\ln \underline{m_i}) + \ln(g_i) d n_i$$

$$\ln \underline{N} \approx n \ln n - n$$

$$d(\ln \underline{N}) \approx (n \cdot \frac{1}{n}) + \ln n - 1 d n$$

$$\approx \ln(n) d n$$

Characteristic
 Stirling's Approximation for large n

$$\Rightarrow d(\ln \Omega) = \sum_i -\ln(n_i) d n_i + \ln(g_i) d n_i$$

$$d(\ln \Omega) = \sum_i \ln \left(\frac{g_i}{n_i} \right) d n_i$$

We know

$$\sum d n_i = 0$$

$$\text{det } \alpha \sum d n_i = 0 \quad - \textcircled{1}$$

α, β : Langrange multipliers

$$\sum \epsilon_i d n_i = 0$$

$$\text{det } \beta \sum \epsilon_i d n_i = 0 \quad - \textcircled{2}$$

$$\sum \ln \left(\frac{g_i}{n_i} \right) d n_i = 0 \quad - \textcircled{3}$$

(at equilibrium)
 $S = \max$
 $dS = 0 \Leftrightarrow d(\ln \Omega) = 0$
 $\Rightarrow d(\ln \Omega) = 0$

$$\textcircled{1} + \textcircled{3} - \textcircled{2}$$

$$\sum_i \left(\ln \left(\frac{g_i}{n_i} \right) + \alpha - \beta \epsilon_i \right) d n_i = 0$$

If it holds for $\forall i$:


$$\Rightarrow \ln \left(\frac{g_i}{n_i} \right) + \alpha - \beta \epsilon_i = 0$$

$$\ln \left(\frac{g_i}{n_i} \right) = - (\alpha - \beta \epsilon_i)$$

$$\frac{g_i}{n_i} = e^{-(\alpha - \beta \epsilon_i)}$$

$$\frac{n_i}{g_i} = e^{\alpha - \beta \epsilon_i}$$

$$\rightarrow \frac{n_i}{g_i} = A e^{-(E_i/kT)}$$



$$n_i = g_i e^{\alpha - \beta E_i} = A g_i e^{-\beta E_i} \text{ where } A = e^\alpha$$

$$\sum n_i = N = A \sum g_i e^{-\beta E_i}$$

This equation is always used to find out random constants ...

$$\Rightarrow A = \frac{N}{\sum g_i e^{-\beta E_i}}$$

$$Z = \sum g_i e^{-\beta E_i}$$

[Partition Function]

$$\Rightarrow n_i = \frac{N}{Z} g_i e^{-\beta E_i}$$

$P(E_i)$: Probability of occupancy of energy level E_i at equilibrium temp T

$$\Rightarrow P(E_i) = \frac{n_i}{N} = \frac{g_i e^{-\beta E_i}}{\sum_i g_i e^{-\beta E_i}}$$

'i' represents on energy level. It is summation over all energy levels.

To determine β

$$S = k \ln \Omega$$

$$dS = k d(\ln \Omega)$$

$$TdS = kT d(\ln \Omega)$$

$$dE = TdS = \left[\sum E_i d n_i = kT d(\ln \Omega) \right]$$

$$\Rightarrow \sum_i E_i dn_i = kT \sum_i -(\alpha - \beta E_i) dn_i$$

$$= -kT \sum_i (\alpha - \beta E_i) dn_i$$

when total energy
is assumed as
constant. Here, a
little perturbation
considered to find
 $\frac{\partial E}{\partial T} \neq 0$
 $\sum E_i dn_i = 0$

$$\Rightarrow \sum E_i dn_i = -\cancel{\alpha kT} \sum dn_i + kT \beta \sum E_i dn_i$$

$[4 \text{ pages}]$

$$\Rightarrow \beta kT = 1$$

$$\boxed{\beta = \frac{1}{kT}}$$

→ Proof of $\beta = \frac{1}{kT}$ is huge
So simply write $\beta = \left(\frac{1}{kT}\right)$
Refer B.J. Mc Clelland
Statistical Thermodynamics.

Hence for Maxwell Boltzmann

$$\left(\frac{n_i}{g_i}\right) = A e^{-\beta E_i}$$

where

$$\boxed{A = \frac{N}{\sum g_i e^{-\beta E_i}}}$$

$$\boxed{\beta = \frac{1}{kT}}$$

$$P(E_i) = \frac{n_i}{N} = \left(\frac{g_i e^{-\beta E_i}}{\sum g_i e^{-\beta E_i}} \right)$$

Note that β is
const. at equilibrium

Z : Partition Function

Plays same role in Statistical Physics as Wave Function does in Quantum Physics.

We know,

$$\langle x \rangle = \frac{\sum x (p(x))}{\sum p(x)}$$

↑
[expectation value]

$$= \frac{\int_{-\infty}^{\infty} x p(x) dx}{\int_{-\infty}^{\infty} p(x) dx}$$



Q) E_i $2kT$ kT 0 N_1 N_2 $g_2 = 1$ $g_1 = 1$ $g_0 = 1$ 100 particles to be filled

Not specified
 \Rightarrow assuming 1 cell per energy level

$$n_0 = A \cdot 1 \cdot e^0 = A$$

$$n_1 = A \cdot e^{-1}$$

$$n_2 = A e^{-2}$$

$$\Rightarrow N = A(1 + e^{-1} + e^{-2}) \Rightarrow A = \left[\frac{N}{1 + e^{-1} + e^{-2}} \right]$$

$$A = \frac{100}{1+0.37+0.09} = \frac{100}{1.38} = 72$$

$$\Rightarrow n_0 = 72$$

$$n_1 = 26$$

$$n_2 = 2$$

$$P(E_i) = \frac{n_i}{N} = \begin{array}{ll} 0.72 & \text{for } E_0 \\ 0.26 & \text{for } E_1 \\ 0.02 & \text{for } E_2 \end{array}$$

$$E = \sum E_i N_i = 0 \times 72 + kT \times 26 + 2kT \times 2 \\ = 30kT$$

$$\langle E \rangle = \frac{\sum E_i P(E_i)}{\sum P(E_i)} = \sum E_i P(E_i) = \sum \frac{n_i}{N} E_i \\ = \frac{1}{N} \sum n_i E_i \\ = \left(\frac{E}{N} \right) \\ = \underline{0.3kT}$$

n_2 _____ E_2

n_1 _____ E_1

We know,

$$n_1 = A g_1 e^{-\beta E_1}$$

This is what we always do... remember Planck's derivation $u(g) \propto$

$$N_0 = N_0$$

$$N_1 = N_0 e^{-\frac{\Delta E}{kT}}$$

$$N_2 = N_0 e^{-\frac{2\Delta E}{kT}}$$

If not specified, take $g_i = 1$

Or take $g_i = \text{const.}$ & same for all states. In this way g_i will be absorbed in the const. A.

$$n_1 = A e^{-\beta E_1}$$

$$n_2 = A e^{-\beta E_2}$$

$$n_2 = \frac{n_1}{e^{-\beta E}} e^{-\beta E_2} = n_1 e^{-\beta(E_2 - E_1)} = n_1 e^{-\left(\frac{\Delta E}{kT}\right)}$$

$$n_2 = n_1 e^{-\left(\frac{\Delta E}{kT}\right)}$$

[at equilibrium]

Population Inversion : $n_2 > n_1$, $\Delta E > 0$

$$\Rightarrow T < 0$$

Concept of Negative Temperature

(Refer end of notes)

Population Inversion is a very short duration phenomenon.

* By applying magnetic properties, population inversion can be achieved for a longer period of time !!

No. of particle within energy level b/w E_i and $E_i + dE$

$$= n(E) dE \quad \leftarrow \text{Notation for } dn$$

$$= dn$$

$$= A e^{-\frac{E}{kT}} g(E) dE \quad \leftarrow \text{Notation for no. of cells with energy b/w } E_i \text{ and } (E_i + dE)$$

$$= A e^{-\frac{E}{kT}} \cdot \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} E^{\frac{1}{2}} dE$$

$$dn = B E^{\frac{1}{2}} e^{-\left(\frac{E}{kT}\right)} dE$$

This is ~~notation~~ formula of no. of particles between energy level E and $(E + dE)$

$$\int_0^\infty dn = B \int_0^\infty E^{\frac{1}{2}} e^{-\frac{E}{kT}} dE$$

$$\Rightarrow N = B \int_0^\infty E^{\frac{1}{2}} e^{-\frac{E}{kT}} dE$$

$$n_i = A g_i e^{-\beta E_i}$$

~~$$g_i = \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} E_i^{\frac{3}{2}}$$~~

$$n(p) = A g(p) e^{-\left(\frac{p^2}{2mkT}\right)}$$

$$n(p) dp = A g(p) dp e^{-\frac{p^2}{2mkT}}$$

$$g(p) dp = \frac{V}{h^3} 4\pi p^2 dp$$

Similarly,

$$\text{Similarly, } n(v) dv = A \frac{V}{h^3} 4\pi m^3 v^2 dv e^{-\left(\frac{mv^2}{2kT}\right)}$$

$$n(v) dv = B v^2 e^{-\left(\frac{mv^2}{2kT}\right)} dv$$

Maxwell Boltzmann Speed distribution

- ① Note that we cannot leave it in terms of h and V ; we have to find the const B in terms of N .

$$\textcircled{O} \int_0^{\infty} e^{-x} x^{n-1} dx = \sqrt{n} \quad (\text{gamma } n) \quad n > 1$$

$$\textcircled{1} \quad \sqrt{n+1} = \sqrt{n} \sqrt{1 + \frac{1}{n}} = \frac{\sqrt{n}}{\Gamma(1 + \frac{1}{n})}$$

$$\textcircled{1} \quad \sqrt{\frac{1}{2}} = \sqrt{\pi}$$

$$\sqrt{\frac{5}{2}} = \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi}$$

$$\textcircled{1} \quad \int_{0}^{\infty} e^{-dx^2} x^n dx = \frac{1}{2(d)^{\frac{n+1}{2}}} \sqrt{\frac{n+1}{2}}$$

$$\textcircled{1} \quad \int_0^{\infty} e^{-\alpha x^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{\alpha}}$$

$$\Rightarrow \int_0^\infty n(v) dv = B \int v^2 e^{-\left(\frac{mv^2}{2kT}\right)} dv$$

$$N = B \frac{1}{2 \left(\frac{m}{2kT}\right)^{3/2}} \cdot \sqrt{\frac{3}{2}} = \frac{B}{4} \left(\frac{\sqrt{\pi}}{\left(\frac{m}{2kT}\right)^{3/2}}\right)$$

$$N = \frac{B\sqrt{\pi}}{4} \left(\frac{2kT}{m}\right)^{3/2}$$

$$\Rightarrow B = \left(\frac{2kT}{m}\right)^{-3/2} \frac{4\sqrt{\pi} N}{\sqrt{\pi}}$$

GAMMA FUNC.

$$\textcircled{1} \quad \Gamma(\mu) = (\mu-1)! = \int_0^\infty x^{\mu-1} e^{-x} dx \quad : \mu > 0$$

$$\textcircled{2} \quad \Gamma(\mu+1) = \mu \Gamma(\mu) \quad \int_{-\infty}^\infty e^{-\alpha x^2 + \beta x} dx = \sqrt{\frac{\pi}{\alpha}} e^{\frac{\beta^2}{4\alpha}}$$

$\textcircled{3}$ The argument of Γ : μ can be a rational no.
and thereby $\Gamma(\mu+1) = \mu(\mu-1)(\mu-2)\dots(1+\beta) \not\in \Gamma(\beta)$
where $0 < \beta \leq 1$. For integer $\mu=n$, $\Gamma(n) = [n-1]$

$$\textcircled{4} \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

$$\textcircled{5} \quad I_\mu = \int_0^\infty x^\mu e^{-\alpha x^2} dx = \frac{1}{2} \frac{1}{\alpha^{\frac{\mu+1}{2}}} \Gamma\left(\frac{\mu+1}{2}\right)$$

$$I_0 = \int_0^\infty e^{-\alpha x^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{\alpha}} \quad \mu = -1$$

$$I_1 = \int_0^\infty x e^{-\alpha x^2} dx = \frac{1}{2} \alpha^{(1)} = \left(\frac{1}{2\alpha}\right) \quad \textcircled{6} \quad \textcircled{1} = 1$$

$$I_2 = \int_0^\infty x^2 e^{-\alpha x^2} dx = \frac{1}{2} \frac{1}{\alpha^{3/2}} \cdot \frac{1}{2} \cdot \sqrt{\pi} = \frac{1}{4} \left(\frac{\pi}{\alpha^3}\right)^{1/2}$$

THERMODYNAMICS (II)

$$f(E_i) = \left(\frac{n_i}{g_i} \right) = A e^{-\beta E_i} : \text{not valid at extremely low temperature !!}$$

Maxwell Boltzmann Speed Distⁿ Formula

$$\text{Energy distribution } g(E) f(E) = A g(E) e^{-E/kT}$$

$n(E) dE$ = no. of particles with energy between E and $(E+dE)$ at equilibrium temperature T .

$$= A e^{-E/kT} \frac{2\pi V}{h^3} (2m)^{3/2} E^{1/2} dE$$

$$= B E^{1/2} e^{-E/kT} dE$$

$$N = B \int_0^\infty E^{1/2} e^{-E/kT} dE$$


$$\text{Put } \frac{E}{kT} = x \rightarrow dE = kT dx$$

$$\Rightarrow N = B(kT)^{3/2} \int_0^\infty x^{1/2} e^{-x} dx = B(kT)^{3/2} \Gamma(\frac{3}{2})$$


$$\Rightarrow N = \frac{\sqrt{\pi}}{2} \cdot B(kT)^{3/2}. \Rightarrow B = \frac{2N}{\sqrt{\pi}(kT)^{3/2}} = \frac{2\pi N}{(\pi kT)^{3/2}}$$

$$\Rightarrow \boxed{n(E) dE = \frac{2\pi N}{(\pi kT)^{3/2}} \sqrt{E} e^{-\left(\frac{E}{kT}\right)} dE}$$

This is Maxwell-Boltzmann Energy distⁿ formula



With increase in Temperature, the curve flattens since no. of particles or Area = const. \Rightarrow the following shapes:



Most Probability

Energy

$$\text{when } \frac{d(n(E))}{dE} = 0$$

$$\frac{dn}{dE} = 0$$

$$\rightarrow \frac{1}{2} E^{-\frac{1}{2}} e^{-\left(\frac{E}{kT}\right)} + E^{\frac{1}{2}} e^{-\left(\frac{E}{kT}\right)} \frac{-1}{kT} = 0$$

$$\Rightarrow \frac{1}{2} \frac{1}{\sqrt{E}} = \frac{\sqrt{E}}{kT}$$

$$\Rightarrow E = \frac{kT}{2}$$

Most Probable Energy

$$\text{Mean Energy } \langle E \rangle$$

$$= \frac{\int_0^\infty E \frac{n(E) dE}{\int_0^\infty n(E) dE}}{\int_0^\infty n(E) dE} = \frac{2\pi}{(\pi kT)^{3/2}} \int_0^\infty E^{3/2} e^{-\frac{E}{kT}} dE$$

Note that $f(x)$ ~~sum~~ does not have ~~max~~ value. The area has a value i.e. the interval has a value

example If I measure width of an oak leaf, the result = 3.5 cm is possible but has Probability = 0, since there are uncountable possibilities even between 3.4 and 3.6. Each of the outcomes has Probability 0. But the outcome will fall b/w [3.4 and 3.6] has non-zero probability.

Put $\frac{E}{kT} = x \Rightarrow dE = kT dx$

$$\begin{aligned} \Rightarrow \langle E \rangle &= \frac{2\pi}{(\pi kT)^{3/2}} \int_0^\infty (kT)^{5/2} \cdot x^{3/2} e^{-x} dx \\ &= \frac{2\pi}{(\pi)^{3/2} (kT)^{3/2}} (kT)^{5/2} \int_0^\infty x^{3/2} e^{-x} dx \\ &= \boxed{\frac{2}{\sqrt{\pi}} (kT) \cdot \frac{3}{4} \sqrt{\pi}} \end{aligned}$$

$$\langle E \rangle = \frac{3}{2} kT$$

Mean Energy

$\star \star$ rather than getting confused with constant. use $\int_0^\infty x(E) dE$ and cut out constants

$$\langle E \rangle = \frac{\langle p^2 \rangle}{2m} = \frac{\langle px^2 \rangle}{2m} + \frac{\langle py^2 \rangle}{2m} + \frac{\langle pz^2 \rangle}{2m}$$

$$\Rightarrow \text{Per degree of freedom Energy} = \frac{1}{2} kT$$

Momentum distribution.

$$n(p) dp = A e^{-\frac{p^2}{2mkT}} g(p) dp$$

= No. of particles with energy between p & $(p+dp)$

$$n(p) dp = A \cdot \frac{4\pi V}{h^3} p^2 e^{-\frac{p^2}{2mkT}} dp$$

$$= B p^2 e^{-\frac{p^2}{2mkT}} dp \quad (\text{say})$$

$$\begin{aligned} N &= \int_0^\infty n(p) dp = B \int_0^\infty p^2 e^{-\frac{p^2}{2mkT}} dp \\ &= \frac{B}{4} \left(\frac{\pi}{\left(\frac{1}{2mkT} \right)^3} \right)^{1/2} \end{aligned}$$

$$\Rightarrow B = \frac{4N}{\sqrt{\pi} (2mkT)^{3/2}} = \frac{4\pi N}{(2\pi m kT)^{3/2}}$$


$$\Rightarrow n(p) dp = \frac{4\pi N}{(2\pi m kT)^{3/2}} p^2 e^{-\frac{p^2}{2mkT}} dp$$

Velocity Distribution

$$\text{Using } p = mv$$

$$n(v) dv = \frac{4\pi N}{(2\pi m kT)^{3/2}} v^2 e^{-\frac{mv^2}{2kT}} dv$$

$$n(v) dv = \frac{4\pi N m^3}{(2\pi m kT)^{3/2}} v^2 e^{-\frac{mv^2}{2kT}} dv$$



$\left[\frac{n(p)}{N} \right]$ = Area b/w p and $(p+dp)$
= fraction of particles
with momentum between
 p and $(p+dp)$

~~Component wise Distribution of Momentum~~

We can write

$$P(p) dp = P(p_x) dp_x \star$$

$$P(p) dp = \frac{n(p)}{N} dp$$

$$P(p_y) dp_y$$

$$P(p_z) dp_z$$

$$P(p) dp = [P(p_x) P(p_y) P(p_z)] [dp_x dp_y dp_z]$$

$$= P(p_x) P(p_y) P(p_z) \frac{4\pi p^2}{(2\pi m kT)^{3/2}} dp$$

$$P(p) dp = \frac{1}{(2\pi m kT)^{3/2}} 4\pi p^2 \left[e^{-\frac{p_x^2}{2mkT}} dp_x e^{-\frac{p_y^2}{2mkT}} dp_y e^{-\frac{p_z^2}{2mkT}} dp_z \right]$$

where $p^2 = p_x^2 + p_y^2 + p_z^2$

P: Probability distribution function

NOTE THAT IF IT IS NOT THE PROBABILITY



We can write component wise,

$$P(p_x) dp_x = \frac{1}{(2\pi mkT)^{1/2}} e^{-\left(\frac{p_x^2}{2mkT}\right)} dp_x$$

On R.H.S.
write

$$\frac{4\pi p_x^2 dp_x}{dp_x dp_y dp_z}$$

$$P(p_y) dp_y = \frac{1}{(2\pi mkT)^{1/2}} e^{-\left(\frac{p_y^2}{2mkT}\right)} dp_y$$

On L.H.S
write

$$n(p) dp$$

$$P(p_z) dp_z = \frac{1}{(2\pi mkT)^{1/2}} e^{-\left(\frac{p_z^2}{2mkT}\right)} dp_z$$

$$\frac{n(p_x) dp_x n(p_y) dp_y}{n(p_z) dp_z}$$

It's called Component Distribution formula

Component
wise formula

[Note that while Probabilities are multiplicative, the fractions or numbers are additive]

Different Velocities

$$n(p_q) dp_q = \frac{N}{(2\pi mkT)^{3/2}} e^{-\left(\frac{p_q^2}{2mkT}\right)} dp_q$$

Following the same steps as analysis of energy,

$$g(v) dv = \cancel{\frac{4\pi V}{h^3} v^2 dv} \quad \frac{d^3x d^3v}{h^3} = \frac{4\pi V}{h^3} v^2 dv$$

$$n(v) dv = f(v) g(v) dv$$

$$= A e^{-\frac{mv^2}{2kT}} \cdot \frac{4\pi V}{h^3} v^2 dv$$

$$N = \int n(v) dv = A \frac{4\pi V}{h^3} \int_0^\infty v^2 e^{-\frac{mv^2}{2kT}} dv$$

$$= B \int_0^\infty v^2 e^{-\left(\frac{m}{2kT}\right)v^2} dv = B \frac{1}{4} \left[\frac{\pi}{\left(\frac{m}{2kT}\right)^3} \right]^{\frac{1}{2}}$$

$$\Rightarrow B = 4 \left(\frac{m}{2kT}\right)^{3/2} \frac{N}{\sqrt{\pi}}$$

$$\Rightarrow n(v) dv = \frac{4\pi N}{(2\pi kT)^{3/2}} v^2 e^{-\frac{mv^2}{2kT}} dv$$

$$P(v) = \frac{n(v) dv}{N}$$

For most probable velocity

$$\frac{d P(v)}{dv} = 0$$

↑ we can this formula straight from Momentum dist'.... Momentum dist' can be derived from Energy dist'.

$$v_{M.P.} \Rightarrow \frac{dn(v)/dv}{v=v_{mp}} = 0$$

$$\Rightarrow \cancel{2\pi} e^{-\frac{(mv^2)}{2kT}} + v^2 \cancel{e^{-\frac{mv^2}{2kT}}} - \frac{m}{2kT} \cancel{2v} = 0$$

$$\Rightarrow 1 = \frac{mv^2}{2kT}$$

$$\Rightarrow v = \sqrt{\frac{2kT}{m}}$$

Most important expression is of that of $n(E) dE$

$$n(E) dE = \frac{2\pi N}{(\pi kT)^{3/2}} E^{1/2} e^{-\frac{E}{kT}}$$

Most Probable Velocity

Mean value or Expectation Value or Average velocity

$$\langle v \rangle = \frac{\int_0^\infty v \frac{n(v) dv}{\int_0^\infty n(v) dv}}{N} = \frac{4\pi N}{N} \left(\frac{m}{2\pi kT} \right)^{3/2} \int_0^\infty v^3 e^{-\frac{mv^2}{2kT}} dv$$

$$= 4\pi \left(\frac{m}{2\pi kT} \right)^{3/2} \frac{1}{2} \cdot \frac{1}{\left(\frac{m}{2kT} \right)^2} \cdot \Gamma(2)$$

$$= \frac{2\pi}{\pi^{3/2}} \cdot \frac{1}{\left(\frac{m\pi}{2kT} \right)^{1/2}} = \sqrt{\frac{4\pi}{m\pi} \cdot \frac{2kT}{m}} = \sqrt{\frac{8kT}{\pi m}}$$

$$\langle v \rangle = \sqrt{\frac{8kT}{\pi m}}$$

Average Velocity

Note that $\langle v \rangle_{avg} > v_{most\ probable}$

$$= \frac{\int_0^\infty v^3 e^{-\frac{mv^2}{2kT}} dv}{\int_0^\infty v^2 e^{-\frac{mv^2}{2kT}} dv}$$

Easier derivation will be done when we mutually cancel the constant and perform both integrations...

$$= \frac{\Gamma(3/2)}{\Gamma(2/2)} = \sqrt{\frac{8kT}{\pi m}}$$

Root Mean Square Velocity

$$v_{rms} = \sqrt{\frac{\int_0^{\infty} v^2 n(v) dv}{\int_0^{\infty} n(v) dv}}$$


$$= \sqrt{4\pi \left(\frac{m}{2\pi kT}\right)^{3/2} \int_0^{\infty} v^4 e^{-\frac{mv^2}{2kT}} dv}$$

$$= \sqrt{4\pi \left(\frac{m}{2\pi kT}\right)^{3/2} \frac{1}{2} \left(\frac{m}{2kT}\right)^{\frac{5}{2}} \Gamma\left(\frac{5}{2}\right)}$$

$$= \sqrt{\cancel{\frac{4\pi}{\pi^{3/2}}}} \frac{1}{\left(\frac{m}{2kT}\right)^{\frac{1}{2}}} \cdot \frac{1}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi}$$

$v_{rms} = \sqrt{\frac{3kT}{m}}$

Root Mean Square Velocity



$$v_{MP} = 1.41 \sqrt{\frac{kT}{m}}, \quad v_{avg} = 1.6 \sqrt{\frac{kT}{m}}, \quad v_{rms} = 1.73 \sqrt{\frac{kT}{m}}$$

 Component wise distribution of velocities

Following the same techniques as in momentum components

i.e. $dv_x dv_y dv_z = 4\pi v^2 dv$

$$\text{Hence } P(v) dv \left(= \frac{n(v) dv}{N} \right)$$

$$= \underline{P(v_x) P(v_y) P(v_z) dv_x dv_y dv_z}$$

$$= \underline{P(v_x) P(v_y) P(v_z) 4\pi v^2 dv}$$

$$= 4\pi \left(\frac{m}{2\pi kT} \right)^{3/2} v^2 e^{-\frac{mv^2}{2kT}} [v_x^2 + v_y^2 + v_z^2] dv$$

$$\Rightarrow \begin{cases} P(v_x) dv_x = \left(\frac{m}{2\pi kT} \right)^{1/2} e^{-\frac{(mv_x^2)}{2kT}} dv_x \\ P(v_y) dv_y = \left(\frac{m}{2\pi kT} \right)^{1/2} e^{-\frac{(mv_y^2)}{2kT}} dv_y \\ P(v_z) dv_z = \left(\frac{m}{2\pi kT} \right)^{1/2} e^{-\frac{(mv_z^2)}{2kT}} dv_z \end{cases}$$

Only thing to note is that it is of $A e^{-\frac{E(v_x)}{kT}}$

We used,

Ek dum normalized curves \neq !!

$$P(v) dv = \cancel{P(v_x) dv_x \cdot P(v_y) dv_y \cdot P(v_z) dv_z}$$

$$= P(v_x) P(v_y) P(v_z) dv_x dv_y dv_z$$


$$= P(v_x) P(v_y) P(v_z) 4\pi v^2 dv$$

Note that the curve for $P(v_x) dv_x$ is symmetric

about $v_x = 0$ i.e. $v_{x \text{ avg}} = 0$

$$\langle v_x^{2n-1} \rangle = 0$$

$$\langle v_x^2 \rangle = \frac{kT}{m}$$



$$\text{num} = \left(\frac{m}{2\pi kT} \right)^{3/2} \cdot \frac{1}{4} \left[\frac{\pi}{\left(\frac{m}{2kT} \right)^3} \right]^{\frac{1}{2}}$$

$$\text{Most Probable velocity of } v_x = 0 = \frac{\sqrt{\pi}}{4 \left(\frac{m}{2kT} \right)} \cdot \frac{1}{\sqrt{\pi}} = \frac{kT}{2m}$$

$$\text{Note that } \int_0^\infty n(v_x) dv_x = \left(\frac{N}{2} \right)$$

To get N , we need to integrate from $(-\infty, \infty)$

$$\text{den} = 2 \cdot \frac{\sqrt{\pi}}{2} \left(\frac{m}{2kT} \right)^{\frac{1}{2}} \cdot \left(\frac{m}{2\pi kT} \right)^{\frac{1}{2}}$$

$$= \frac{2 \sqrt{\pi}}{2 \sqrt{\pi}} \cdot \frac{1}{2} = \frac{1}{2}$$

$$\Rightarrow \langle v_x^2 \rangle = \left(\frac{kT}{m} \right)^2 = \frac{2}{2} = 1$$

Equipartition of Energy Principle

[PTO to last Page]
of notes for Proof

For Any thermodynamic system at equilibrium temperature T ,
the energy per degree of freedom is $(\frac{1}{2}kT)$,

provided it is expressible in form of $E_q = \alpha q^2$.

$$U_x = \frac{1}{2} kx^2 ; U_y = \frac{1}{2} ky^2 ; U_z = \frac{1}{2} kz^2$$

$$\left[\frac{n(q) dq}{N} \right] = \left(\frac{m}{2\pi kT} \right)^{\frac{1}{2}} e^{-\frac{\alpha q^2}{kT}} dq$$

$$P(q) = \frac{n(q)}{N}$$

$$\langle E_q \rangle = \frac{\int E_q P(q) dq}{\int P(q) dq} = \frac{\int_0^\infty \alpha q^2 P(q) dq}{\int_0^\infty P(q) dq}$$

$$= \frac{\int_0^\infty \alpha q^2 \left(\frac{m}{2\pi kT} \right)^{\frac{1}{2}} e^{-\frac{\alpha q^2}{kT}} dq}{\int_0^\infty \left(\frac{m}{2\pi kT} \right)^{\frac{1}{2}} e^{-\frac{\alpha q^2}{kT}} dq}$$

$$= \frac{\int_0^\infty \alpha q^2 e^{-\frac{\alpha q^2}{kT}} dq}{\int_0^\infty e^{-\frac{\alpha q^2}{kT}} dq}$$

$$= \alpha \left(\frac{I_2}{I_0} \right) = \frac{\alpha \frac{1}{4} \frac{\sqrt{\pi}}{\alpha^{3/2}}}{\frac{1}{2} \frac{\sqrt{\pi}}{\alpha^{1/2}}} = \frac{\alpha}{2} \frac{1}{\alpha}$$

$$= \frac{\alpha}{2} \cdot \frac{1}{\left(\frac{\alpha}{kT} \right)} = \underline{\underline{\left(\frac{kT}{2} \right)}}$$

For 1 Gm-Mole Ideal Gas,

$$U = N \cdot f \cdot \frac{1}{2} kT = f \cdot \frac{1}{2} RT$$

Read this page

$$\textcircled{1} \quad C_V = \frac{fR}{2}$$

$$\textcircled{2} \quad C_P = C_V + R = \left(\frac{f}{2} + 1\right) R$$

$$\textcircled{3} \quad \left[\frac{C_P}{C_V} \right] = \frac{1 + \frac{f}{2}}{\frac{f}{2}} = \underbrace{1 + \frac{2}{f}}_{= r}$$

$$\begin{aligned} dU &= C_V dT \\ &= \frac{fR}{2} dT \\ \Rightarrow C_V &= \frac{fR}{2} \end{aligned}$$

Monoatomic Gas

$$f = 3 \Rightarrow r = 1 + \frac{2}{3} = 1.66$$

Diatomic Gas

$$f = 5 \Rightarrow r = 1 + \frac{2}{5} = 1.4$$

Triatomic Gas

$$r = (1.29, 1.42) \quad \begin{bmatrix} 3N-k \\ N=3 \text{ (Triatomic)} \end{bmatrix}$$

or
Polyatomic Gas

Note that while deriving law of equipartition of energy, we assumed energy to have a continuous spectrum which is not the case.


Thus equipartition law breaks down at the level of low temperatures where kT is significantly smaller than spacing between energy levels. Here arises the need of quantum theory.

<https://ourstudycircle.in/upscpdf/>

FERMI DIRAC STATISTICS

(Thermodynamics)
(12)


9/1/2012



* Note that 2 spin states are actually 2 different cells

g_{cells} is multiplied by 2

→ always remember that cells (g_i) are always distinguishable because they are the unique slot numbers.



Difference b/w FD and BE is restriction on no. of particles per cell due to Pauli Exclusion Principle.

For F.D., $g_i \geq n_i$ For Pauli Exclusion Principle to hold true.

i.e. no. of cells must be more than no. of particles

⇒ way of choosing is to choose n_i cells out of g_i to fill particles in them. i.e. $\prod_i g_i^{n_i}$

$$\Omega = \prod_i \frac{g_i}{n_i! g_i - n_i!}$$

Here also conceiving it as 2 step problem $(3, 5, 2)$

$$\Omega = \Omega_1 \Omega_2$$

$\Rightarrow \Omega_1 = 1$ (as all particle are indistinguishable)

$$\Rightarrow \Omega_2 = \prod_i g_i^{n_i}$$

∴ which 3, which 5 etc. do not matter

$$\ln \Omega = \sum \ln L_{gi} - \ln L_{ni} - \ln \underline{g_{i-ni}}$$

$$\begin{aligned}
 d(\ln \Omega) &= \sum d(\ln L_{gi}) - d(\ln L_{ni}) - d(\ln \underline{g_{i-ni}}) \\
 &\quad \uparrow \\
 &\quad 0 \\
 &= \sum d \left\{ -n_i \ln n_i + n_i \right\} - d \left\{ (g_{i-ni}) \ln (g_{i-ni}) \right. \\
 &\quad \left. - (g_{i-ni}) \right\} \\
 &= \sum -n_i \cancel{\frac{1}{n_i}} - \ln n_i \cancel{d n_i} + \cancel{d n_i} \neq d n_i \\
 &\quad + (g_{i-ni}) \cancel{\left(\frac{1}{g_{i-ni}} \right)} + \ln (g_{i-ni}) d n_i \\
 &= \sum \ln \left(\frac{g_{i-ni}}{n_i} \right) d n_i \quad \text{--- } \textcircled{3}
 \end{aligned}$$

We know

$$\alpha \sum d n_i = 0 \quad \text{--- } \textcircled{1}$$

$$\beta \sum \epsilon_i d n_i = 0 \quad \text{--- } \textcircled{2}$$

Doing $\textcircled{1} + \textcircled{3} - \textcircled{2}$

$$\sum_i \left[\ln \left(\frac{g_{i-ni}}{n_i} \right) + \alpha - \beta \epsilon_i \right] d n_i = 0$$

$$\begin{aligned}
 \Rightarrow \text{ If hold good } \forall i, \Rightarrow d n_i \neq 0 \text{ for all } i \\
 \Rightarrow \ln \left(\frac{g_{i-ni}}{n_i} \right) + \alpha - \beta \epsilon_i = 0
 \end{aligned}$$

$$\Rightarrow \left[\frac{g_i - n_i}{n_i} \right] = e^{-(\alpha - \beta E_i)}$$

To find variables, $S = k \ln \Omega$

$$T dS = T k d(\ln \Omega)$$

$$\sum g_i d n_i = -kT \sum (\alpha - \beta E_i)$$

$$\Rightarrow \boxed{\beta = \frac{1}{kT}}$$

$d(\ln \Omega)$ may be different but in terms of α and β , it is same
Hence β की समीक्षा नहीं हो सकती !!

$$\Rightarrow \frac{g_i - n_i}{n_i} = \left(\frac{g_i}{n_i} \right) - 1 = e^{-(\alpha - \beta E_i)}$$

$$f(E_i) = \frac{1}{\left(\frac{g_i}{n_i} \right)} = \frac{1}{1 + e^{-(\alpha - \beta E_i)}} = \frac{1}{1 + e^{-(\alpha - \frac{E_i}{kT})}}$$

define $\alpha = \frac{E_f}{kT}$ — where E_f : characteristic energy of fermions called fermi energy at temperature T

$$\Rightarrow \boxed{f(E_i) = \left(\frac{n_i}{g_i} \right) = \frac{1}{1 + e^{\left(\frac{E_i - E_f}{kT} \right)}}}$$

Note that α can be $+\frac{E_f}{kT}$ or $-\frac{E_f}{kT}$ depending upon whether we do $(1) + (3) - (2)$ or $(1) + (3) - (2) \dots$ etc. Finally we should get this formula of distribution.

Fermi Dirac Distribution


$$f(E) = \frac{n(E)}{g(E)} = \frac{1}{1 + e^{(E-E_f)/kT}} \quad \text{--- (1)}$$

Note that $f(E)$ is always less than 1. It makes sense as no. of particles per energy ~~cell~~ cannot be more than 1.

@ $T=0K$ [just for definition; it's unattainable]

From (1),
if $E < E_f$, we get $f(E) = 1$ (all states are fully occupied)

if $E > E_f$, we get $f(E) = 0$ (no state is occupied)



Hence 2 parts of complete defⁿ of E_f :

✓ Characteristic of fermions at Temp T.

✗ @ 0K, energy below which all states are fully occupied and above which no state is occupied.

Now 2 analysis in syllabus


(1) @ $T=0$, (i) find E_f (iii) find P_0
(ii) find E

(2) @ $T=T$ use $\int_0^{\infty} f(x-x_0) x^n dx = \frac{x_0^{n+1}}{n+1} \left[1 + \frac{\pi^2}{6} \frac{n(n+1)}{x_0^2} + \dots \right]$



in order to

(i) find $E_f(T)$
(ii) Contribution of Energy by e⁻
(iii) find C_V, P_T

Use $\frac{4\pi N}{h^3} (2m)^{\frac{3}{2}} = \frac{3}{2} \frac{N}{E_f^{\frac{3}{2}}}$
in both



Note that $f = \frac{1}{2}$ at E_f and not E_{f_0} . E_f lies close to but not at E_{f_0} .



(10)

$$\begin{aligned} E &= \alpha q^2 \\ \langle E \rangle &= \frac{\int E P(E) dE}{\int P(E) dE} = \frac{\int_0^\infty E f(E) dE}{\int_0^\infty f(E) dE} \\ &= \frac{\int E \frac{e^{-(\alpha + \beta E)}}{\int e^{-(\alpha + \beta E)} dE} dE}{\int e^{-(\alpha + \beta E)} dE} = \frac{\int E e^{-\beta G_i} dE}{\int e^{-\beta G_i} dE} \end{aligned}$$

$$\text{Let } \beta E = x \Rightarrow dE = \frac{dx}{\beta}$$

$$= \frac{1}{\alpha\beta} \cdot \frac{\Gamma_2}{\Gamma_1} = \frac{1}{\alpha\beta}$$

$$\underline{\underline{\beta = \frac{1}{kT}}}$$

~~.....~~

∴

check for errors.

2nd way

$$\text{Put } E = \alpha q^2 \quad dE = 2\alpha q dq$$

$$\alpha\beta q^2 = x$$

$$\sqrt{\alpha\beta} q = \sqrt{x}$$

$$q = \frac{1}{\sqrt{\alpha\beta}} \sqrt{x}$$

$$dq = \frac{1}{2\sqrt{\alpha\beta}} x^{-1/2} dx$$

$$\boxed{\frac{1}{\beta}}$$

$$\underline{\underline{\beta = \frac{1}{kT}}}$$

$$f(\epsilon) = \frac{1}{1 + e^{(\epsilon - E_f)/kT}}$$

@ 0 K : $f(\epsilon) = 1 \quad \epsilon < E_f$
 $= 0 \quad \epsilon > E_f$

No. of fermions at 0 K

$$\int dn_i = \int f(\epsilon_i) g(\epsilon_i) d\epsilon_i$$

$$= \int_0^{E_f} 1 \cdot \frac{2\pi V}{h^3} (2m)^{3/2} \epsilon^{1/2} \cdot d\epsilon$$

Space degeneracy

$$N = \frac{2\pi V}{h^3} (2m)^{3/2} \left[\frac{\left(\frac{E_f}{3/2} \right)^{3/2}}{\left(\frac{3/2}{3/2} \right)} \right]$$

in general at temp T,

$$N = \frac{2\pi V}{h^3} (2m)^{3/2} \int_0^{\infty} \frac{f}{1 + e^{(\epsilon - E_f)/kT}} \epsilon^{1/2} d\epsilon$$

If spin degeneracy is also taken into account, $g = J_{\text{space}} * J_{\text{sp}}$

$$g(\epsilon) = \frac{2\pi V}{h^3} (2m)^{3/2} \epsilon^{1/2}$$

on space cells per state

degeneracy of state or no. of particular energy level on

no. of prob.
states per
energy level

$$g_{\text{spin}} = (2S+1)$$

$$\text{For } e^- : S = \frac{1}{2} \Rightarrow g_{\text{spin } e} = 2$$

Hence $2e^-$ can be put in same space or cell if spin is taken into account.

$\frac{6h^2}{8ml^2}$	(2, 1, 1)	(1, 2, 1)	(1, 1, 2)	
	(Corresponding 6 spin states)			
$\frac{3h^2}{8ml^2}$	(1, 1, 1)			
	(1, 1, 1, $\frac{1}{2}$)	&		
	(1, 1, 1, $-\frac{1}{2}$)			

* Fermions like e^- , protons, neutrons have $S = \frac{1}{2}$

\Rightarrow every energy level E can have 2 particles.

(here we mean every cell)

$$\Rightarrow N = 2 * N_{\text{space degeneracy}}$$

* वर्ता 1 constant की ही फूंक पड़ेगा !!

$$N = 2 \times \frac{2\pi V}{h^3} (2m)^{3/2} \left[\frac{(E_f)_0^{3/2}}{3/2} \right]$$

$$\Rightarrow \boxed{\frac{4\pi V}{h^3} (2m)^{3/2} = \frac{3}{2} \frac{N}{E_f_0^{3/2}}}$$

This is important !!

$$n = \frac{N}{V} \Rightarrow 3\pi^2 n \hbar^3 = (2m E_{f_0})^{3/2}$$

$$\hbar = \frac{h}{2\pi}$$

(h cross)

$$E_{f_0} = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

at 0 k

$$n = \frac{\rho N}{M} = \left(\frac{N}{V}\right)$$

If $T = T_f \neq 0 \Rightarrow$ we have to integrate

Note the limits

$$\int_0^\infty \frac{e^{Ex}}{1 + e^{(E - E_f)/kT}} dE$$

Use

$$\begin{aligned} & \int_0^\infty f(x-x_0) x^n dx \\ &= \frac{x_0^{n+1}}{n+1} \left[1 + \frac{\pi^2}{6} \frac{n(n+1)}{x_0^2} + \dots \right] \end{aligned}$$

Sommerfeld lemma

$$\text{by Putting } \frac{E}{kT} = x \quad \text{and} \quad \frac{E_f}{kT} = x_0$$

$$\downarrow \text{Total no. of particles at a finite temperature}$$

$$N = \left[\frac{4\pi V}{h^3} (2m)^{3/2} \right]$$

$$\int_0^\infty \frac{(kT)^{3/2} x^{1/2} dx}{1 + e^{x-x_0}}$$

$$N = \left[\frac{3}{2} \cdot \frac{N}{E_{f_0}^{3/2}} (kT)^{3/2} \right] (x_0)^{3/2} \frac{2}{3} \left[1 + \frac{\pi^2}{6} x_0^{1/2} \times \frac{8}{2x_0^2} + \dots \right]$$

$$\Rightarrow 1 = \left(\frac{E_f}{E_{f_0}} \right)^{3/2} \left[1 + \frac{\pi^2}{8} \cdot \left(\frac{kT}{E_f} \right)^2 + \dots \right]$$

$$E_f(T) = E_{f_0} \left[1 - \frac{\pi^2}{12} \left(\frac{kT}{E_f} \right)^2 + \dots \right]$$

defining $E_f = k T_f$
 ↑
fermi temperature

* Remember
 $\frac{E_f}{kT} = \alpha$ (const)
 $\Rightarrow E_f = f(T)$
 But remember approximations like $(g_i + n_i) \gg 1$ were used. Hence expression for E_f is complex

$$E_f(T) = E_{f_0} \left[1 - \frac{\pi^2}{12} \left(\frac{T}{T_f} \right)^2 + \dots \right]$$

$T_f > 10^4$ i.e. very high values hence $E_f(T)$ and E_0 have little difference
 For $E_f = 3.2 \text{ eV}$ $T_f \approx 10^4$

$$N = \left[\int f(p) g(p) dp \right] \times \left[(2^S + 1) \right] \quad (\text{if spin taken into account})$$

$$f(p) = \frac{1}{1 + e^{\frac{p^2 - p_f^2}{2mkT}}}$$

$$\text{define: } \cdot E_f = \frac{p_f^2}{2m}$$

$$\textcircled{a} \quad T=0 \text{ K} : \quad \begin{cases} \text{for } p < p_f & f(0) = 1 \\ p > p_f & f(0) = 0 \end{cases} \quad \}$$

$$\int 2 \times f(p) \frac{V}{h^3} 4\pi p^2 dp$$

$$N = \frac{8\pi V}{h^3} \frac{p_f^2}{3}$$

Total Energy

$$E = \cancel{\int \epsilon dn_e}$$

$$\int \epsilon dn_e = \int \epsilon n(\epsilon) d\epsilon$$

$$= \int \epsilon f(\epsilon) g(\epsilon) d\epsilon$$

$$= \int_0^\infty \epsilon \cdot \frac{4\pi V}{h^3} \cdot (2\pi)^{3/2} \frac{\epsilon^{1/2}}{1 + e^{(\epsilon - E_f)/kT}} d\epsilon$$

↙

$$E = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty \frac{\epsilon^{3/2} d\epsilon}{1 + e^{(\epsilon - E_f)/kT}}$$

~~(3/8)~~ @ T=0

$$\frac{4\pi V}{h^3} (2m)^{3/2} \int_0^{E_f} e^{-\epsilon/kT} d\epsilon$$

$$\frac{3N}{2} (E_{f_0})^{3/2} \frac{2}{5} \frac{e_f^{5/2}}{k_B T}$$

↙

$$E = \frac{3}{5} N E_{f_0}$$

This shows that unlike classical particle, a fermion has appreciable energy even at 0 K.
Also $C_V = \left(\frac{\partial E}{\partial T}\right)_V = 0$ at 0 K.

$\frac{E}{N} = \frac{3}{5} E_{f_0} = 60\% \text{ of fermi energy}$
at 0 K

$$PV = \frac{2}{3} E$$

(Considering fermions as an ideal gas)


$$P = \frac{2}{3} \left(\frac{E}{V} \right) = \frac{2}{3} \cdot \frac{3}{5} \frac{N E_f}{V} = \frac{2}{5} \frac{N E_f}{V}$$

$$= \frac{2}{5} n \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

$$\Rightarrow P \propto n^{5/3}$$

* This high pressure is counter balanced by Coulomb attraction of e- by ions.

⑤ In an ideal fermi system, there is no inter-fermionic interaction



$$T_1 < T_2 < T_3$$

Thermodynamics (13)

10/01/2012

$$\int_0^\infty f(x-x_0) x^n dx = \frac{x_0^{n+1}}{n+1} \left[1 + \frac{\pi^2}{6} \frac{n(n+1)}{x_0^2} + \dots \right]$$

$$\int_0^\infty \frac{e^{-x}}{x^{n-1}} dx = \Gamma_n G(n)$$

↑ ↑
 gamma zeta

We know : Contribution of e- to Internal Energy.

F.D.

$$f(E) = \frac{n(E)}{g(E)} = \frac{1}{1 + e^{(E-E_f)/kT}}$$

$$n(E) = f(E) g(E)$$

↑

In MB and BE, we can put all particles in these energy level as there is no restriction....

$$dn = n(E) dE = 2 * f(E) * g(E)$$

↑

$$n = \int_0^\infty 2 * f(E) * g(E) dE$$

spin degeneracy

$$E = \int_0^\infty E_i dn$$

$$\text{At } 0 \text{ K, } n = \int_0^{E_{f0}} 2 g(E) dE$$

$$E = \int_0^\infty \epsilon^2 f(\epsilon) g(\epsilon) d\epsilon$$

$$E = \frac{4\pi V}{h^3} (2m)^{3/2} \int_0^\infty \frac{\epsilon^{3/2} d\epsilon}{1 + e^{(\epsilon - E_f)/kT}}$$

$$E = \frac{3N}{2 E_{f_0}^{3/2}} \int_0^\infty \frac{\epsilon^{3/2} d\epsilon}{1 + e^{(\epsilon - E_f)/kT}}$$

Energy contribution to internal energy

by ~~lattice~~ / electrons.... Note that it is electronic contribution, not lattice contribution (ions)

$$\text{Put } \frac{\epsilon}{kT} = x$$

$$d\epsilon = kT dx$$

$$\Rightarrow E = \frac{3N}{2 E_{f_0}^{3/2}} (kT)^{5/2} \int_0^\infty \frac{x^{3/2} dx}{1 + e^{(x - x_0)}}$$

$$x_0 = \frac{E_f}{kT}$$

$$n = \frac{3}{2}$$

$$\Rightarrow E = \frac{3N}{2 E_{f_0}^{3/2}} (kT)^{5/2} \cancel{\leq} \frac{(E_f)^{5/2}}{(kT)^{5/2}} \left[1 + \frac{\pi^2}{6} \cdot \frac{3}{2} \frac{5}{2} \left(\frac{kT}{E_f} \right)^2 + \dots \right]$$

$$\Rightarrow \boxed{E = \frac{3}{5} N \frac{E_f^{5/2}}{E_{f_0}^{3/2}} \left[1 + \frac{5\pi^2}{8} \left(\frac{kT}{E_f} \right)^2 + \dots \right]}$$

$$@ T = 0 K, E = \frac{3}{5} N E_f$$

Simplifying further, expressing E_f in terms of E_0 .

$$E = \frac{3N}{5} E_{f_0}^{\frac{5}{2}} \left[1 + \frac{\pi^2}{12} \left(\frac{kT}{E_f} \right)^2 + \dots \right]^{\frac{5}{2}} \left[1 + \frac{5\pi^2}{8} \left(\frac{kT}{E_f} \right)^2 + \dots \right]$$

$$E = \frac{3N}{5} E_{f_0} \left[1 - \frac{5\pi^2}{288} \left(\frac{kT^2}{E_f} \right)^2 + \frac{5\pi^2}{8} \left(\frac{kT}{E_f} \right)^2 + \dots \right]$$

$$\boxed{E = \frac{3N}{5} E_{f_0} \left[1 + \frac{5\pi^2}{12} \left(\frac{kT}{E_f} \right)^2 \right]}$$

\uparrow
Contribution of Fermions or electronic contribution

Specific Heat of Gases

Here we talk about "Ideal Fermi Gas" or "Ideal Fermi System".

→ Fermi Gas : Particles obey Pauli Exclusion Principle.

→ Ideal Fermi Gas : Non interacting particles

This does not mean particles are in gaseous state. It just means no interaction among particles.

$$C_V = \left(\frac{dE}{dT} \right)_V = \underbrace{\frac{3N}{5} E_{f_0} \left[\frac{5\pi^2}{12} \left(\frac{k}{E_f} \right)^2 \right]}_A 2T + \dots$$

$\approx AT$

$$A = \frac{N \epsilon_F \pi^2 k^2}{2 E_f^2}$$

$$dQ = C_V dT$$

$$dS = \frac{dQ}{T} = \frac{C_V dT}{T} = A dT$$

$$P(T) = \frac{2}{3} \frac{E(T)}{V}$$

↑

$$\left[\begin{array}{l} \frac{1}{2} mnv^2 = E \\ \frac{1}{3} mnv^2 = PV \end{array} \right]$$

Pressure exerted by electrons = $\frac{2}{3} * \frac{3N\epsilon_{f_0}}{5V} \left[1 + \frac{5\pi^2}{12} \left(\frac{kT}{\epsilon_f} \right)^2 + \dots \right]$

$$P(T) = \frac{2}{5} n \epsilon_{f_0} \left[1 + \frac{5\pi^2}{12} \left(\frac{kT}{\epsilon_f} \right)^2 + \dots \right]$$

Even at 0K, $P = \frac{2}{5} n \epsilon_{f_0}$

This Pressure is balanced by attraction b/w ions and electrons. Whenever this balanced is maintained, star survives. The limit after which star collapses, is called Chandrasekhar limit.

Important Points in F-D:

- derivation along with graph is important
- fermi energy, total no. of particles: typically at $T=0 K$
- Total Energy $E = \frac{3}{5} N \epsilon_{f_0}$


$$\epsilon_f = \frac{\hbar^2}{2m} (3\pi^2 n)^{2/3}$$

Strong degeneracy: all levels below E_f : fully occupied
all levels above E_f : empty.

Bose - Einstein Statistics

Valid for indistinguishable particle having full integral spin, having no restriction on no. of particles per degeneracy level.
e.g. Photons (quanta of energy of EM waves)
Phonons (minimum energy or quanta of elastic waves)

even $\xrightarrow{^4_2 \text{He}}$ (all even A nuclei)
Mass Number A



$$\Omega = \prod_i \binom{g_i}{n_i}$$

← no of ways to put n_i in g_i degeneracy level

let

$$n_1 = 3$$

$$g_1 = 2$$



$$\text{No. of ways: } (g_1) (g_1 - 1) \dots (n_1 - g_1)$$

$$\text{Total no. of ways} = (n_i + g_i - 1)!$$

Consider it as 2 step
 Problem (3, 5, 2)
 $\Omega_1 = 1$ (as all are
 indistinguishable)
 Consider n=3 box:

$$= \frac{(n_i + g_i - 1)!}{n_i! g_i!}$$

Let particles be lettered
 (a, b, c, d, ...). Let levels
 be numbered (1, 2, 3, ...)
 The dist'n of particles among
 levels may be written as:
 1 (ab) 2 (c) 3 4 (def) ...

n_i 's are not distinguishable
but g_i 's are distinguishable
for sure

$$\Omega = \prod_i \frac{(n_i + g_i - 1)!}{n_i! g_i!}$$

approximating

do not miss this step

$$\Rightarrow \Omega = \prod_i$$

$$n_i + g_i \gg 1$$

$$\Rightarrow n_i + g_i - 1 \approx n_i + g_i$$

If the numbers are arranged in all possible sequences, provided the sequence begins with a no., the no. of ways are $g_i (N_i + g_i - 1)!$

But (I) is equivalent to

$$2(c) 3 1(a) 4(d, e, f) \dots \quad (\text{II})$$

Hence there will be $1g$ such sequences.

Also (I) is equivalent to

$$1(f, g) 2(a) 3 4(b, c, d) \dots$$

since the particles are indistinguishable,
distinguishable particles can be arranged
in l^m ways

$$\Rightarrow \text{Total No. of Ways} = \frac{(n_i + g_i - 1)!}{g_i! n_i!}$$

$$\Rightarrow \Omega_2 = \prod_i \frac{(n_i + g_i - 1)! / (g_i - 1)! n_i!}{g_i! n_i!}$$

$$\ln \Omega = \sum (n_i + g_i) \ln(n_i + g_i) - (n_i + g_i) - n_i \ln n_i + n_i - \ln(g_i - 1)$$

$$\begin{aligned} d(\ln \Omega) &= \sum \ln(n_i + g_i) d n_i - \ln(n_i) d n_i \\ &= \sum \ln \left[\frac{n_i + g_i}{n_i} \right] d n_i \end{aligned}$$

For $\Omega(N, V, E)$,

$$d \sum d n_i = 0$$

$$\beta \sum E_i d n_i = 0$$

$$\sum \ln \left(\frac{n_i + g_i}{n_i} \right) d n_i = 0$$

$$\left[d(\ln \Omega) = \frac{d\Omega}{k} \right]$$

= 0 at equilibrium

$$\Rightarrow \sum \left(\ln \left(\frac{n_i + g_i}{n_i} \right) + \alpha - \beta E_i \right) d n_i = 0$$

$$\Rightarrow \frac{n_i + g_i}{n_i} = e^{-(\alpha - \beta E_i)} \quad \left[\text{to be valid } \forall i \right]$$

$$\Rightarrow f(E_i) = \left(\frac{n_i}{g_i} \right) = \frac{1}{e^{-(\alpha - \beta E_i)} - 1} = \frac{1}{A e^{\beta E_i} - 1}$$

$A < 1$
FUGACITY

$$\beta = \left(\frac{1}{kT} \right) \quad \text{as we found for other 2 distributions}$$

Assume

$$e^{-\alpha} = A$$

or

$$\mu = -\frac{\partial F}{\partial T}$$

Chemical potential

of assembly of bosons

$$f(E_i) = \frac{1}{e^{\left(\frac{(E_i - \mu)}{kT} \right)} - 1}$$

In assembly of Photons, $\mu = 0$

$$f(\epsilon_i) = \frac{1}{e^{\epsilon_i/kT} - 1}$$

$$f(\gamma) = \frac{1}{e^{h\gamma/kT} - 1}$$

$\mu = 0$ for $T < T_B$
and
Photons

For Photon $g(\epsilon) d\epsilon = g(\gamma) d\gamma$



$$g(\gamma) d\gamma = \frac{d^3x d^3p}{h^3}$$

Blackbody Radiation

Planck's Radiation Formula :

$$u(\gamma) d\gamma = \frac{8\pi h\gamma^3}{c^3} \frac{d\gamma}{e^{h\gamma/kT} - 1}$$

Energy density radiated
in between frequency γ and
 $(\gamma + d\gamma)$ at equilibrium Temp 'T'



$T > T_{\text{sur}}$

If heat transfer w/o heating the intervening medium:
Radiation

BlackBody: Body at higher Temperature, able to radiate out heat.

E radiated per unit volume of blackbody = Energy density

As $T \uparrow$, colour of radiation shifts from Red to Blue.

$$\frac{E(\gamma) d\gamma}{V}$$

3 postulates of Planck's Quantum Theory

Atoms of Blackbody behave like oscillating particles.

① They radiate energy, not continuously, but discretely in quantum of energies.

EM quanta: Photons

Solid Body: Elastic waves: Phonons

② $E_{\text{radiation}} \propto \lambda^2$


$$E_{\text{radiated}} = h\nu$$

E_{absorbed}

h : Planck's const.

③ $E_{\text{emission or absorption}} = nh\nu$ $n = 1, 2, 3, \dots$

\Rightarrow energy levels in an atom are always discrete



→ Blackbody radiation chamber emits photons of energy $h\nu$.

$$E = \sum g_i n_i = \int \epsilon dn = \int h\nu dn$$

dn will come from BE distribution due to Photons being bosons.

$$= \int h\nu f(\epsilon) g(\epsilon) d\epsilon$$

$$= \int h\nu \frac{1}{e^{(h\nu)/kT} - 1} g(\nu) d\nu$$

To find $g(\nu) d\nu$:

$$\begin{aligned} g(b) d(b) &= \frac{d^3 r d^3 b}{h^3} \\ &= \frac{V}{h^3} \cdot 4\pi b^2 db \end{aligned}$$

$$\boxed{\begin{aligned} b &= \frac{h\nu}{c} \\ db &= \frac{h}{c} d\nu \end{aligned}}$$

$$= \frac{4\pi V}{h^3} \frac{h^2}{c^2} \nu^2 \frac{h}{c} d\nu$$

$$= \frac{4\pi V}{c^3} \nu^2 d\nu = g(\nu) d\nu$$

But EM wave can have 2 modes of polarization

$$\Rightarrow \boxed{g(\nu) d\nu = \frac{8\pi V}{c^3} \nu^2 d\nu}$$



$$\Rightarrow dN = f(\epsilon) g(\epsilon) d\epsilon$$

↑
No. of Photons Emitted

$$= \frac{1}{e^{\frac{h\nu}{kT}} - 1} \left(\frac{8\pi\nu}{c^3} \right) \nu^2 d\nu$$

No. of photons at temp 'T' in any blackbody radiating EM radiation, per unit volume

$$\left(\frac{N}{V} \right) = n = \int dN$$

$$= \int_0^{\infty} \left(\frac{8\pi\nu}{c^3} \right) \left(e^{\frac{h\nu}{kT}} - 1 \right)^{-1} \nu^2 d\nu$$

$$\text{Put } \frac{h\nu}{kT} = x$$

$$d\nu = \left(\frac{kT}{h} \right) dx$$

$$n = \frac{8\pi}{c^3} \int_0^{\infty} \left(\frac{kT}{h} \right)^3 \frac{x^2 dx}{(e^x - 1)}$$

$$= 8\pi \left(\frac{kT}{ch} \right)^3 \int \frac{x^2 dx}{e^x - 1}$$

$$\Rightarrow [n \propto T^3]$$

★ No. of photons in a cavity vary as T^3

while energy varies as T^4

$$dE = h\nu dN \Rightarrow u(\nu) d\nu = \frac{dE}{V} = \frac{8\pi h\nu^3 d\nu}{c^3 [e^{\frac{h\nu}{kT}} - 1]}$$

$$E = \int h\gamma dn$$

per unit volume

$$= \int h\gamma \cdot 8\pi \left(\frac{kT}{ch}\right)^3 \cdot \left(\frac{x^2}{e^{x-1}}\right) dx$$

$$= \int \frac{8\pi (kT)^4}{(ch)^3} \cdot \frac{x^3}{(e^{x-1})} dx$$

We can also write

$$dE dy = \frac{8\pi}{ck^3} \cdot \left(\frac{kT}{x}\right)^3 \cdot \frac{1}{(e^{\frac{h\gamma}{kT}} - 1)} \left(\frac{h}{kT}\right) dx$$

per unit volume

$$\Rightarrow u(y) dx = \frac{8\pi h}{c^3} y^3 \frac{dy}{e^{h\gamma/kT} - 1}$$



This formula that we got after solving G.P.s
and A.G.P.s of M.B. distⁿ is
found directly with B.E. distⁿ

Thermodynamics (14)

Criteria for Classical Statistics to hold good

$$\left(\frac{g_i}{n_i}\right) = e^{\alpha + \beta E_i} \pm 1$$

$$\underbrace{\left(\frac{g_i}{n_i}\right) \neq 1}_{\text{quantum mechanical}} = \underbrace{e^{\alpha + \beta E_i}}_{\text{classical}}$$

whenever $\boxed{\left(\frac{g_i}{n_i}\right) > 1}$, then quantum statistics (QS)
 converges to classical statistics (CS), i.e. indistinguishable
 particles become distinguishable.

(2) Volume occupied by 1 Particle = $\left(\frac{V}{N}\right)$

length available for 1-d motion (assume cube) = $\left(\frac{V}{N}\right)^{1/3}$

Characteristic of a Particle: DeBroglie's Wavelength

$$\lambda = \frac{h}{\sqrt{2mE}} = \frac{h}{\sqrt{2mkT}}$$

*we can also
use $\frac{3}{2}kT$...
no difference*

\uparrow
E for 2 degrees of freedom = kT

For distinguishable, there should be no overlap of
particles.

$$\text{i.e. } \lambda = \frac{h}{\sqrt{2mkT}} < \left(\frac{V}{N}\right)^{1/3}$$

\rightarrow This is the
usual case:
1 kinetic and
1 potential

Hence for this condition, Classical Statistics is valid.
 Otherwise, we have to consider Quantum Effects.

$$\frac{h}{\sqrt{2mkT}} < \left(\frac{V}{N}\right)^{1/3}$$

$$T > \left(\frac{N}{V}\right)^{\frac{2}{3}} \frac{h^2}{2mk}$$

$$\left(\frac{N}{V}\right)^{1/3} \frac{h}{\sqrt{2mkT}} < 1$$

$$n^{1/3} \frac{h}{\sqrt{2mkT}} < 1$$

$$\left[\frac{n}{T^{3/2}}\right] \left(\frac{h^3}{2mk}\right)^{3/2} < 1$$

i.e.

$$\boxed{\frac{n}{T^{3/2}} \left(\frac{h^2}{2mk}\right)^{3/2} < 1}$$

Region of validity of
Classical statistics

Specific Heat of Solids

$$C_p - C_v = T \left(\frac{\partial P}{\partial T} \right)_V \left(\frac{\partial V}{\partial T} \right)_P$$

↑
0 (for solids)

$$\Rightarrow C_p = C_v$$

$$C_v = \left(\frac{\partial Q}{\partial T} \right)_V = \left(\frac{\partial E}{\partial T} \right)_V$$

① Fulop - Petru Law

For ideal gas, 1 gm-Mole

$$E = N * f * \frac{1}{2} kT \\ = \frac{f}{Z} * R * T$$

$$\Rightarrow C_V = \frac{f R}{Z}$$

④ Dulong and Petit's law is experimentally valid at room temperature i.e. at room temperature all solids have

$$C_V \approx 25 \text{ J mol}^{-1} \text{ K}^{-1}$$

[Independent of Temperature]
for ideal gas

For Solids, $E(T) = ?$

Dulong Petit (1870) : Classical
Einstein Theory (1907) } : Quantum
Debye Theory (1910)

For solid to be solid, every molecule is bound to the solid even while moving / oscillating.

And at any Temp T, molecules are in motion

⇒ Molecules are under elastic force.


This elastic force is responsible for inter molecular forces.

For 1 particle, $E = \frac{1}{2} m \dot{x}^2 + \frac{1}{2} kx^2$ [for 1-d motion]

It is classical energy, as for any x we can have any E , thus E is continuous.

$$= \frac{1}{2} kT + \frac{1}{2} kT \quad (\text{equipartition of energy}) \\ = kT$$

★ Now we can consider each particle to be composed of 3 oscillators




$$\Rightarrow E = 3kT \quad \text{vibrating in all 3 directions.}$$

$$E_{\text{Total}} = N * 3kT = 3RT \Rightarrow C_V = 3R$$

Dulong & Petit formula

② Einstein Theory

- Dulong & Petit formula valid for high Temperatures.
 \therefore law of equipartition is valid only for higher temperatures
 But we have seen that C_v varies with Temperature



No one was able to interpret these graphs upto 30 years till 1907.

By applying Planck's Quantum Theory, Einstein was able to explain this variation.

Minimum energy which an atom can have is Phonon.
 Its vibrational energy.

$$\gamma = \frac{1}{2\pi} \sqrt{\frac{k_1}{m}}$$

$$E \propto \gamma$$

$$E = h\gamma$$

Note that elastic waves are not transverse waves.

~~1~~ Einstein said that all molecules are vibrating with same frequency. This frequency is characteristic of solid.

~~2~~ Each molecule has 3 degrees of freedom

~~3~~ Energy emission and absorption is in multiples of $h\gamma$

~~4~~ Planck gave avg. Energy of 1 molecule per degree of freedom

$$\langle E \rangle = \frac{h\gamma}{e^{\left(\frac{h\gamma}{kT}\right)} - 1}$$

(Proof of this @ end of copy)

For 1 gm mole,

$$E = N * 3 * \left(\frac{h\nu}{e^{\frac{h\nu}{kT}} - 1} \right)$$

it was (kT) in classical statistics

$\langle E \rangle$: average energy of a molecule using Planck's Quantum Theory. Also called average energy of Planck Oscillator.

$$C_V = \left(\frac{\partial E}{\partial T} \right)_V = 3NAk \left[\frac{+1}{\left(e^{\frac{h\nu}{kT}} - 1 \right)^2} e^{\frac{h\nu}{kT}} \left(\frac{h\nu}{kT} \right) \frac{1}{T^2} \right]$$
$$= 3NAk \left(\frac{h\nu}{kT} \right)^2 \frac{e^{\frac{h\nu}{kT}}}{\left(e^{\frac{h\nu}{kT}} - 1 \right)^2}$$

$$\boxed{C_V = 3R \left(\frac{h\nu}{kT} \right)^2 \frac{e^{\frac{h\nu}{kT}}}{\left[e^{\frac{h\nu}{kT}} - 1 \right]^2}}$$

$\left(\frac{h\nu}{k} \right) = \theta_E$: Einstein Temperature

ν = characteristic frequency.

$\left(\frac{\theta_E}{T} \right)$: reduced Temperature

do not think too much
frequency is same

these are
overtones of
same
frequency.


something like $\left(n + \frac{1}{2} \right) \hbar \nu$

\Rightarrow

$$C_V = 3R \left(\frac{\theta_E}{T} \right)^2 \frac{e^{(\theta_E/T)}}{\left[e^{(\theta_E/T)} - 1 \right]^2}$$

Specific Heat of solid, at given Temperature T , is given by Einstein Formula.

where R is (Nk) where k is $-$, N is $-$
 θ_E is (hv/k) where h is $-$, v is $-$, k is $-$
 T is temperature



Shortcoming : ① It does not explain observed C_V at very lower temperatures.
 ② How to know θ_E ??

At higher temperatures, $\frac{\theta_E}{T}$ is small

$$\text{s.t. } e^{\frac{\theta_E}{T}} = 1 + \left(\frac{\theta_E}{T} \right)$$

$$\Rightarrow C_V = 3R \left(\frac{\theta_E}{T} \right)^2 \cdot \frac{\left(1 + \frac{\theta_E}{T} \right)}{\left[1 + \frac{\theta_E}{T} - 1 \right]^2} \approx 3R$$

As $T \rightarrow 0$ $\frac{\theta_E}{T}$ is large

$$e^{\theta_E/T} \gg 1$$

$$\Rightarrow C_v = 3R \frac{\left(\frac{\theta_E}{T}\right)^2}{e^{\frac{\theta_E}{T}}} = \frac{3R \left(\frac{\theta_E}{T}\right)^2}{1 + \left(\frac{\theta_E}{T}\right) + \frac{1}{2!} \left(\frac{\theta_E}{T}\right)^2 + \frac{1}{3!} \left(\frac{\theta_E}{T}\right)^3}$$
$$= \frac{3R}{\left(\frac{\theta_E}{R}\right)^2 + \frac{1}{\left(\frac{\theta_E}{R}\right)} + \frac{1}{2!} + \frac{1}{3!} \left(\frac{\theta_E}{T}\right) + \dots}$$
$$\approx \frac{3R}{\frac{1}{2!} + \frac{1}{3!} \left(\frac{\theta_E}{T}\right) + \dots} \rightarrow \frac{3R}{\infty} \rightarrow 0$$

This is explained satisfactorily.

Method of determination of characteristic frequency of solids


Einstein gave an innovative method to determine ν_E .

Method of Curve Fitting

Here the shortcoming is Arbitrary fixation of ν_E without any empirical measure.


i.e. Plot empirically the variation of C_v

Now fix the Plot of C_v as given by Einstein's Formula upon the above Plot and find ν for the material.



③ Debye's Theory of Specific Heat of Solids

- ✓ All molecules are not vibrating with same frequency. They are vibrating forming a discrete frequency spectrum. ($\gamma_1, \gamma_2, \dots, \gamma_m$)
- ✓ Vibrations of molecules are said to form elastic waves. Elastic Waves are longitudinal & transverse. Minimum quanta = $h\nu$
- ✓ Total modes of vibrations of solids are limited in number = $3N$



Mode $g(p) dp = \frac{V}{h^3} \frac{4\pi p^2}{\cancel{\lambda}} dp \quad p = \frac{h\nu}{\lambda}$

★ Momentum of a wave = $\frac{h}{\lambda}$
 $= \left(\frac{h\nu}{\lambda}\right)$

$$\frac{V}{h^3} \cdot 4\pi \cdot \frac{h^2 \cdot h}{V^2} \cdot \frac{y^2 dy}{V}$$

$$g(y)dy = V \cdot \frac{4\pi^2 y^2 dy}{V^3}$$


In a solid of volume V ,

$$g(y)dy \text{ (longitudinal)} = \frac{4V\pi^2 y^2}{V_L^3} dy \Rightarrow 3N = \int_0^{y_m} g(y) dy$$

$$g(y)dy \text{ (transverse)} = \frac{8\pi V y^2}{V_T^3} dy$$

Total modes of vibration of solid in volume V in frequency range y to $y+dy$

$$\boxed{\text{Total Modes} = 4\pi V y^2 \left[\frac{1}{V_L^3} + \frac{2}{V_T^3} \right] dy}$$



$$y = \frac{1}{2\pi\sqrt{k/m}}$$

$$4\pi V \int y^2 \left(\frac{1}{V_L^3} + \frac{2}{V_T^3} \right) dy$$

④ No. of molecules vibrating in frequency range y and $y+dy$ is
 $dn = f(y) g(y) dy$
Assume $f(y) = 1$
{(like we assume $g(y) = 1$ in equipartition proof)
(b'coz we are finding no. of states)
 $\Rightarrow dn = g(y) dy$ }

Although it was conceptualized discrete mode, for simpler analysis we treat them as continuous from $(0 \text{ to } y_m)$

Since total number of Modes are finite, it puts a cap on maximum value of frequency in the spectrum.

$$4\pi V \left(\frac{1}{V_L^3} + \frac{2}{V_T^3} \right) \frac{\nu_m^3}{3} = 3N$$

$\nu_m : \nu_{max}$

$$\boxed{4\pi V \left(\frac{1}{V_L^3} + \frac{2}{V_T^3} \right) = \left[\frac{9N}{2\nu_m^3} \right]}$$

Hence,

$$\boxed{\frac{9N}{2\nu_m^3} \nu^2 d\nu : \text{Modes of vibration between frequency } \nu \text{ and } \nu + d\nu}$$

dE : Energy of solid at equilibrium Temperature
 T with frequency b/w ν and $(\nu + d\nu)$

= (Modes of vibration) * (Energy of 1-mode)

$$= \frac{9N}{2\nu_m^3} \nu^2 d\nu \underbrace{e^{\frac{h\nu}{kT}}}_{\text{Note that this already contains } f(E) \text{ as well}}$$

Note that this already contains $f(E)$ as well as E

Hence no need of separate $f(E)$

→

$$\boxed{dE = \left[\frac{9Nh}{2\nu_m^3} \right] \frac{\nu^3 d\nu}{(e^{h\nu/kT} - 1)}}$$

$$E = \int_0^{\nu_m} \frac{9Nh}{2\nu_m^3} \frac{\nu^3 d\nu}{(e^{h\nu/kT} - 1)}$$

$$\text{Let } \frac{h\nu}{kT} = x \quad \nu = \frac{kT}{h} x \quad d\nu = \frac{kT}{h} dx$$

$$E = \frac{qNh}{\gamma_m^3} \int_0^{\frac{h\gamma_m}{kT}} \left(\frac{kT}{h}\right)^4 \frac{x^3 dx}{e^{x-1}}$$

$$= \frac{qNh}{\gamma_m^3} \left(\frac{kT}{h}\right)^4 \int_0^{\frac{h\gamma_m}{kT}} \frac{x^3 dx}{e^{x-1}}$$

$$\star \frac{qNk}{\gamma_m} \left(\frac{k}{h\gamma_m}\right)^3 T^4$$

$$\frac{h\gamma_m}{k} = \Theta_D : \text{Debye Temperature}$$

$$E = qR \frac{T^4}{\Theta_D^3} \int_0^{(\Theta_D/T)} \frac{x^3 dx}{e^{x-1}}$$

where $\Theta_D = \frac{h\gamma_m}{k}$

$$x = \left(\frac{h\gamma_m}{kT}\right)$$



Note that

$E = \frac{1}{2} kT \cdot 3N$

step 1-1 \rightarrow $\frac{1}{2} kT$

change $\frac{1}{2} kT \rightarrow \frac{1}{2} E$!!

Dulong & Petit

Einstein

Debye

(degree of freedom) $< E_{av}$

$3N * kT$

$$3N * \frac{h\nu}{e^{\frac{h\nu}{kT}} - 1}$$

$$\frac{qN\gamma^2 dv}{\gamma_m^3} * \frac{h\nu}{e^{\frac{h\nu}{kT}} - 1}$$

although total modes
are still $3N$

Cv

$3R$

$$3R \left(\frac{\partial E}{\partial T}\right)^2 \frac{e^{\frac{\partial E}{T}}}{(e^{\frac{\partial E}{T}} - 1)}$$

$$qR \left[4 \left(\frac{1}{\Theta_D}\right)^3 \int_0^{\Theta_D} \frac{x^3 dx}{e^{x-1}} - \left(\frac{\Theta_D}{T}\right) \frac{1}{e^{\frac{\Theta_D}{T}} - 1} \right]$$

Thermodynamics (15)

12/01/2012

By Debye's Theory,

$$E = \frac{qR}{\theta_D^3} T^4 \int_0^{\hbar \gamma_m / kT} \frac{x^3 dx}{e^{x-1}}$$

$$\text{where } \theta_D = \frac{\hbar \gamma_m}{k}$$

= Debye's Temp.

$$x = \frac{\hbar \gamma}{kT}$$

$$C_V = \left(\frac{dE}{dT} \right)_V = \frac{qR}{\theta_D^3} \frac{d}{dT} \left[T^4 \int_0^{\hbar \gamma / kT} \left(\frac{x^3 dx}{e^{x-1}} \right) \right]$$

$$= \frac{qR}{\theta_D^3} \left[4T^3 \int_0^{\theta_D/T} \frac{x^3 dx}{e^{x-1}} + T^4 \frac{d}{dT} \int_0^{(\theta_D/T)} \frac{x^3 dx}{e^{x-1}} \right]$$

Leibnitz Rule

$$\frac{d}{dt} \int_{u(t)}^{v(t)} f(x) dx = f(v) \left(\frac{dv}{dt} \right) - f(u) \left(\frac{du}{dt} \right)$$

Using this rule

$$\begin{aligned} \frac{d}{dT} \int_0^{(\theta_D/T)} \frac{x^3 dx}{e^{x-1}} &= \frac{(\theta_D/T)^3}{e^{\theta_D/T} - 1} \cdot \left[-\frac{\theta_D}{T^2} \right] \\ &= \frac{-\theta_D^4}{T^5 (e^{\theta_D/T} - 1)} \end{aligned}$$

$$\Rightarrow C_V = \frac{qR}{\theta_D^3} \left[4T^3 \int_0^{\theta_D/T} \frac{x^3}{e^{x-1}} dx + T^4 \cdot \frac{-\theta_D^4}{T^5 (e^{\theta_D/T} - 1)} \right]$$

$$C_V = qR \left[4 \left(\frac{T}{\theta_D} \right)^3 \int_0^{\theta_D/T} \frac{x^3}{e^{x-1}} dx - \left(\frac{\theta_D}{T} \right) \frac{1}{e^{\theta_D/T} - 1} \right]$$

↑

This is "DEBYE SPECIFIC HEAT FORMULA"

Q4/2u

Einstein

Debye

① Const. γ

discrete frequency spectrum

② $\gamma_1 = \gamma_2 = \dots$

Definite upper bound

$\gamma_1, \gamma_2, \dots, \gamma_m$

③ 3 dof

No meaning of individual dof

At higher Temperature region : $T \gg \theta_D$

$$\Rightarrow \underbrace{\left(\frac{\theta_D}{T} \right)}_{x \text{ is small}} \ll 1 \quad \Rightarrow \underbrace{e^x \approx 1+x}_{\text{x is small}}$$

$$C_V = qR \left[4 \left(\frac{T}{\theta_D} \right)^3 \int_0^{\theta_D/T} \frac{x^3}{x} dx - \frac{\theta_D}{T} \frac{1}{\left(\frac{\theta_D}{T} \right)} \right]$$

$$= qR \left[4 \left(\frac{T}{\theta_D} \right)^3 - \frac{1}{3} \left(\frac{\theta_D}{T} \right)^3 - 1 \right]$$

$$= qR \cdot \left[\frac{4}{3} - 1 \right]$$

$$\boxed{C_V = 3R}$$

At lower Temperature region

$$T \ll \theta_D$$

x is large

$$\frac{\theta_D}{T} \gg 1 \quad i.e. \frac{\theta_D}{T} \rightarrow \infty$$

$$C_V = qR \left[4 \left(\frac{T}{\theta_D} \right)^3 \int_0^\infty \frac{x^3 dx}{e^x - 1} - 0 \right]$$

↑
by expanding e^x

$$= qR \left[4 \left(\frac{T}{\theta_D} \right)^3 \int_0^\infty \frac{x^3 dx}{e^x - 1} \right]$$

$$= \left[\frac{36R}{\theta_D^3} \int_0^\infty \frac{x^3}{e^x - 1} dx \right] T^3$$


→ also used in
derivation of
Stefan Boltzmann's
law

$$\boxed{\int_0^\infty \frac{x^3}{e^x - 1} dx = \frac{\pi^4}{90}}$$

≈ 6.5


$$\boxed{C_V = A T^3}$$

$$\text{where } A = \frac{36R}{\theta_D^3} \int_0^\infty \frac{x^3}{e^x - 1} dx$$



Q5 | Tut 2⁴

Debye Theory valid for isotropic solids



There are some variations in Anisotropic Solids.

Q8 Corresponding States

2 solids are said to be in correspondence states if their reduced parameters are same , then their specific heat are also same.


$$\frac{\theta_{D1}}{T_1} = \frac{\theta_{D2}}{T_2} \Rightarrow C_{V1} = C_{V2} : \text{Debye's Theory}$$

Similarly if,

$$\frac{\theta_{E1}}{T_1} = \frac{\theta_{E2}}{T_2} \Rightarrow C_{V1} = C_{V2} : \text{Einstein's Theory}$$

$$C_{V1} = C_{V2} \Rightarrow \frac{\theta_{E1}}{T_1} = \frac{\theta_{E2}}{T_1}$$

$$T_1' = T_1 \left(\frac{\theta_{E2}}{\theta_{E1}} \right)$$



Tut 26 | 10

$$n_i = g_i e^{-\alpha - \beta \epsilon_i}$$

$$\langle E \rangle = \frac{3}{2} kT = \frac{\int \epsilon_i d n_i}{\int d n_i}$$

$$= \frac{\int \epsilon g(\epsilon) e^{-\beta \epsilon} d\epsilon}{\int g(\epsilon) e^{-\beta \epsilon} d\epsilon}$$

$$\text{Use } \epsilon = a q^2$$

$$g(\epsilon) = c \epsilon^{1/2}$$

Bose-Einstein Condensation

In Bose-Einstein Distribution.

$$\frac{n(\epsilon)}{g(\epsilon)} = f(\epsilon) = \frac{1}{e^{(\epsilon-\mu)/kT} - 1} = \left(\frac{1}{A}\right) e^{\mu/kT} - 1$$

$$\frac{n_i}{g_i} > 0 \Rightarrow e^{\frac{\epsilon_i - \mu}{kT}} > 1$$

$$\Rightarrow \frac{\epsilon_i - \mu}{kT} > 0$$

$$\Rightarrow \epsilon_i \geq \mu \Rightarrow \mu \leq 0$$

as ϵ can be 0 also

$$\boxed{\text{Always : } \mu \leq 0}$$

$$\boxed{A = e^{\mu/kT}}$$

Fugacity of
Assembly of
Bosons

① is valid for all ϵ ,

Putting $\epsilon = 0$

$$\frac{n(\epsilon)}{g(\epsilon)} = \frac{1}{\left(\frac{1}{A}\right) - 1}$$

$$\Rightarrow \frac{1}{A} \geq 1 \Rightarrow \boxed{A < 1}$$

$$\ln A < \ln 1$$

$$\Rightarrow \frac{\mu}{kT} \leq 0$$

✓

$$\Rightarrow \boxed{\mu \leq 0}$$

✓ Bose Einstein characteristics are required when T is close to 0
 we are cooling down Boson.
 A stage is reached when μ becomes 0.

Number of Bosons in ground state (minimum energy state)

$$n(\epsilon) d\epsilon = g(\epsilon) f(\epsilon) d\epsilon$$

$$N = N_0 + \frac{2\pi V}{h^3} \cdot (2m)^{3/2} \cancel{\int_0^\infty} \frac{\epsilon^{1/2}}{e^{(\epsilon-\mu)/kT} - 1} d\epsilon$$

A stage is reached at $T = T_B$, $\mu = 0$!! [strongly degenerate gas]

$$N = N_0 + \frac{2\pi V}{h^3} (2m)^{3/2} \int_0^\infty \frac{\epsilon^{1/2}}{e^{\epsilon/kT} - 1} d\epsilon$$

$$= \left[\frac{2\pi V}{h^3} \cdot (2m)^{3/2} \right] (kT_B)^{3/2} \int_0^\infty \frac{x^{1/2}}{e^x - 1} dx$$

(brace under the integral)

$$N = N_0 + \left[\frac{2.612}{h^3} (2m\pi)^{3/2} V \right] (kT)^{3/2}$$

(brace under the first term) $\Gamma(\frac{3}{2})$ (brace under the second term) $(\frac{3}{2})$

Zeta

① We can write let this be $\left(\frac{N}{kT_B} \right)^{3/2} \frac{1}{2} \sqrt{\pi} \cdot 2.612$.

$$\Rightarrow (kT_B)^{3/2} = \frac{\hbar^3 4\pi^2}{(2m)^{3/2}} \left(\frac{N}{V} \right) \cdot \frac{1}{\Gamma(\frac{5}{2}) \zeta(\frac{3}{2})}$$

(brace under the first term) $\left(\frac{2.612 (2\pi m)^{3/2} V}{h^3} \right) (kT_B)^{3/2}$
 (brace under the second term) $= N$

$$\alpha = \frac{N}{(kT_B)^{3/2}}$$

$$(kT_B)^{3/2} = \left(\frac{\hbar^2}{2m}\right)^{3/2} 4\pi^2 n \frac{1}{\Gamma(\frac{3}{2}) \zeta(\frac{3}{2})}$$

$$\Rightarrow T_B = \frac{2\pi\hbar^3}{mk_B} \left[\frac{n}{\zeta(\frac{3}{2})} \right]^{2/3}$$

$$\Rightarrow kT_B = \left(\frac{\hbar^2}{2m} \right) (4\pi^2 n)^{2/3} \left(\frac{1}{\Gamma(\frac{3}{2}) \zeta(\frac{3}{2})} \right)^{2/3}$$

$$\Rightarrow T_B \propto n^{2/3} \quad \dots \dots \text{Bose Einstein Temperature}$$

Characteristic Temperature of Bosons when $\mu=0$.

② We can also write

$$\frac{N}{(kT_B)^{3/2}} = \frac{2\pi V}{\hbar^3} (2m)^{3/2} \Gamma(\frac{3}{2}) \zeta(\frac{3}{2})$$

nothing but written the same thing....

$$N_{ex} = \left[\frac{2\pi V}{\hbar^3} (2m)^{3/2} \Gamma(\frac{3}{2}) \zeta(\frac{3}{2}) \int_0^\infty \frac{e^{E/2} dE}{e^{E/kT} - 1} \right]^{3/2}$$

Note that this does not take into account energy at ground state.

 It does not matter in FD b'coz in F-D only some particles can come in ground state.

$$\Rightarrow N = N_g + \frac{N}{(kT_B)^{3/2} \Gamma(\frac{3}{2}) \zeta(\frac{3}{2})} \int_0^\infty \frac{e^{-E/kT}}{e^{E/kT} - 1} dE$$

$\mu=0$



$$= N_g + \frac{N (kT)^{3/2}}{(kT_B)^{3/2}}$$

✓

$$\Rightarrow N_g = N - N \left[\frac{T}{T_B} \right]^{3/2}$$

$$\Rightarrow N_g = N \left(1 - \left(\frac{T}{T_B} \right)^{3/2} \right)$$

✓




Refer to more theory @ end of lecture
& more maths from end of copy

As Temp falls below T_B , population increases rapidly in ground state.
At $T=0$, all particles will be in ground state.

5th state of matter

It is responsible for abnormal properties exhibited by ^4He when it is cooled below T_B . Below T_B , it is called Helium 2. It is anomalous fluid. No opposition to flow. Fountain Effect in Capillary Tubes immersed in bath of Helium 2.



VIRIAL THEOREM OF CLASIUS

$$\bar{T} = -\frac{1}{2} \sum \vec{F}_i \cdot \vec{x}_i$$

or

$$2\bar{T} + \bar{V} = 0$$

where $\bar{V} = \sum \vec{F}_i \cdot \vec{x}_i$
 = "Virial"

$$\text{where } \vec{F}_i = m_i \frac{d^2 \vec{x}_i}{dt^2} = m_i \frac{d \vec{v}_i}{dt} = \left(\frac{d \vec{p}_i}{dt} \right)$$

Derivation

$$\text{let } \phi = \sum \vec{p}_i \cdot \vec{x}_i \quad \boxed{\text{---}} \quad \left. \begin{array}{l} \phi: \text{mathematical} \\ \text{construct} \end{array} \right\}$$

ϕ is either periodic or aperiodic

If Periodic, $\phi(\tau) = \phi(0)$

If aperiodic, τ is very large.

$$\frac{d\phi}{dt} = \sum \vec{p}_i \cdot \frac{d\vec{x}_i}{dt} + \sum \frac{d\vec{p}_i}{dt} \cdot \vec{x}_i$$

$$\left(\frac{d\phi}{dt} \right) = 2T + \sum \vec{F}_i \cdot \vec{x}_i$$

Taking Time average,

$$\frac{\int_0^T \left(\frac{d\phi}{dt} \right) dt}{T} = 2\bar{T} + \overline{\sum \vec{F}_i \cdot \vec{v}_i}$$

$$\frac{\phi(T) - \phi(0)}{T} = 2\bar{T} + \overline{\sum \vec{F}_i \cdot \vec{v}_i}$$

$$= 2\bar{T} + \overline{V}$$

V: Virial
~~~~~


L.H.S. = 0 whether  $\phi$  is periodic or aperiodic

$$\checkmark \frac{\text{num} = 0}{\text{den} = \infty}$$

$$\Rightarrow 2\bar{T} + V = 0 \quad V = \overline{\sum \vec{F}_i \cdot \vec{v}_i}$$

### Application

Pressure exerted by molecules of an ideal gas



$$F_i = \frac{2m_i \vec{v}_i}{T} = P_i \vec{\Delta A}$$

$$T = +\frac{1}{2} \sum P_i \vec{\Delta A} \cdot \vec{r} = +\frac{P}{2} \sum \vec{\Delta A} \cdot \vec{r}$$

$$= +\frac{P}{2} \int_S d\vec{A} \cdot \vec{r}$$

$$= +\frac{P}{2} \int_V \nabla \cdot \vec{r} dv = +\frac{P}{2} \int_S 3 dv = +\frac{3PV}{2}$$

$\nabla \cdot \vec{r} = 3$

$$T = -\frac{3}{2} PV$$

$$\text{Time Avg.} = \bar{T} = \sum \frac{1}{2} m_i (\overline{v_i^2}) \\ = \frac{1}{2} m N \overline{v_i^2}$$

Also, we can use

$$\bar{T} = \frac{3}{2} N k T$$

$$\Rightarrow PV = RT$$

$$\Rightarrow \frac{3}{2} PV = \frac{1}{2} m N \overline{v_i^2}$$

$$\Rightarrow PV = \frac{1}{3} m N \overline{v_i^2}$$

### Bose Einstein Condensate

BEC is a state of matter of dilute gas of weakly interacting bosons cooled to temperatures very near absolute zero.

Under such conditions, very large fraction of bosons occupy the lowest quantum state at which point quantum effects become apparent on macroscopic scale.

This transition occurs below a critical temperature, which for a uniform non interacting boson gas is given by

$$T_B = \frac{2\pi \hbar^2}{m k_B} \left[ \frac{n}{\zeta(\frac{3}{2})} \right]^{\frac{2}{3}} \approx 3.31 \left( \frac{\hbar^2 n^{2/3}}{m k_B} \right)$$

$n$ : particle density

$m$ : mass per boson

$\hbar$ : reduced Planck Const.

$k_B$ : Boltzmann Const.

$\zeta(\frac{3}{2})$ : Riemann Zeta Function  $\approx 2.61$

Two examples of materials containing BECs are superconductors and superfluids. No resistance in superconductor.

No viscosity in fluid. e.g. He

Also refer to last page !!

## Basic Postulates:

## KINETIC THEORY OF GASES

- ✓ Gas consists of very small particles i.e. total volume of particles  $\ll V_{\text{container}}$
- i.e. avg. distance between particles  $\gg$  size of particles
- ✓ These particles have same mass.
- ✓ They are in const. random motion, colliding elastically with walls of container, without interacting with each other
- ✓  $N$  (no. of molecules) is so large that statistical treatment can be applied
- ✓  $k \cdot E_{\text{avg}}$  depends only on Temperature.

### Derivations:



Pressure is explained by kinetic theory by assuming it to arise from forces exerted by molecules impacting on walls of container.

$$\Delta p_x = 2m v_x^2 \quad \Delta t = \left( \frac{2L}{v_x} \right) \Rightarrow F = \frac{\Delta p}{\Delta t} = \left( \frac{mv_x^2}{L} \right)$$

$$F_{\text{Total}} = \frac{Nm v_x^2}{L} = \frac{Nm \overline{v^2}}{L}$$

Since particles are assumed to move randomly,

$$\overline{v_x^2} = \overline{v_y^2} = \overline{v_z^2} = \left( \frac{\overline{v^2}}{3} \right)$$

$$\Rightarrow F = \frac{Nm}{L} \frac{\overline{v^2}}{3}$$

$$P = \frac{Nm}{L^3} \left( \frac{\overline{v^2}}{3} \right) = \frac{nm \overline{v^2}}{3}$$

$P = \frac{1}{3} mn \overline{v^2}$

where  $n$  is the number density of gas

From ideal gas law

$$PV = N k_B T$$

$$\Rightarrow PV = \frac{Nm \bar{v}^2}{3} = N k_B T$$

$$\Rightarrow T = \frac{m \bar{v}^2}{3 k_B} = \frac{m v_{rms}^2}{3 k_B}$$

From this,

$$\boxed{(k \cdot E)_{\text{molecule}} = \frac{1}{2} m v_{rms}^2 = \frac{1}{2} \cdot 3 k_B T = \frac{3}{2} kT}$$

$$\boxed{(k \cdot E)_{\text{system}} = \frac{1}{2} N m v_{rms}^2 = \frac{1}{2} \cdot 3 N k_B T = \frac{3}{2} R T}$$

Also  $PV = N k_B \cdot \frac{2}{3} \frac{k \cdot E}{N k_B} = \frac{2}{3} k \cdot E$ .

$$\Rightarrow \boxed{PV = \frac{2}{3} kE}$$

### 3 Tds equations

$$S = f(T, V) : T dS = C_V dT + T \left( \frac{\partial P}{\partial T} \right)_V dV$$

$$S = f(T, P) : T dS = C_P dT - T \left( \frac{\partial V}{\partial T} \right)_P dP$$

$$S = f(P, V) : T dS = C_V \cdot \left( \frac{\partial T}{\partial P} \right)_V dP + C_P \cdot \left( \frac{\partial T}{\partial V} \right)_P dV$$

# Mathematics

- ① All thermodynamic quantities can be divided into 2 groups : functions of state & functions of a process.  
 $(S, U \text{ etc.})$        $(Q, W)$

Only functions of state are path or process independent  
and hence ONLY FOR THEM

$$\text{if } dz = M dx + N dy$$

$$\Rightarrow \left( \frac{\partial M}{\partial y} \right)_x = \left( \frac{\partial N}{\partial x} \right)_y$$

---

IT IS NOT VALID FOR PATH DEPENDENT VARIABLES.

} It's called Total Differential

- ② If  $dz = 0$  i.e.  $z = \text{Const.}$

$$\Rightarrow \left( \frac{\partial z}{\partial x} \right)_y \cdot dx + \left( \frac{\partial z}{\partial y} \right)_x \cdot dy = dz = 0 \quad \dots \quad (1)$$

$$\Rightarrow \left( \frac{\partial z}{\partial x} \right)_y dx + \left( \frac{\partial z}{\partial y} \right)_x dy = 0$$

$$\Rightarrow \boxed{\left( \frac{\partial z}{\partial x} \right)_y + \left( \frac{\partial z}{\partial y} \right)_x \left( \frac{\partial y}{\partial x} \right)_z = 0}$$

$$\Rightarrow 1 + \left( \frac{\partial z}{\partial y} \right)_x \left( \frac{\partial y}{\partial x} \right)_z \left( \frac{\partial x}{\partial z} \right)_y = 0$$

$$\Rightarrow \boxed{\left( \frac{\partial z}{\partial y} \right)_x \left( \frac{\partial y}{\partial x} \right)_z \left( \frac{\partial x}{\partial z} \right)_y = -1}$$

$$\boxed{\left( \frac{\partial x}{\partial y} \right)_z \left( \frac{\partial y}{\partial z} \right)_x \left( \frac{\partial z}{\partial x} \right)_y = -1}$$

$\leftarrow$  cyclic

Differentiating ① with respect to  $x$  keeping

a Property OF STATE  $\lambda$  Const.

i.e. differentiating

$$dz = \left(\frac{\partial z}{\partial x}\right)_y dx + \left(\frac{\partial z}{\partial y}\right)_x dy$$

We get,

$$\left(\frac{\partial z}{\partial x}\right)_\lambda = \left(\frac{\partial z}{\partial x}\right)_y + \left(\frac{\partial z}{\partial y}\right)_x \left(\frac{\partial y}{\partial x}\right)_\lambda$$

It allows us to get relation between partial derivatives of one & the same quantity calculated, however, on

the basis of different const. parameters, e.g.  $C_p$  and  $C_v$ .

So basically if we require  $\left(\frac{\partial x}{\partial a}\right)_{p_1} - \left(\frac{\partial x}{\partial a}\right)_{p_2}$ , write  $X = f(a, p_1)$

$$X = f(a, p_2)$$

- ★ Global Warming does not violate the 2<sup>nd</sup> law of thermodynamics. Some people suggest that as Earth is source of heat and  $CO_2$  is thus colder, it cannot make the Earth hotter.

Note that Global Warming is not that Earth surface is getting hotter. It's just that heat is being trapped in  $CO_2$  and its loss to outer space is slowed down.

It's similar to wearing a blanket. Body radiates heat. Blanket prevents heat getting lost. It's not that blanket transfers heat from outside to body.

# TRANSPORT PHENOMENON

o Bad Pointee.

The Proofs done in class are Perfect.

Revise the Proof  $\leftarrow \text{GTM} \rightarrow$

## Law of Equipartition of Energy

Energy of a particle is in general a function of a number of different parameters. These might be velocity components, amplitude of oscillation etc. Each of these parameters is called a degree of freedom.

Let  $z$  represent any such parameter and  $E(z)$  represent the energy associated with that parameter.

If energy can be represented as a continuous function of parameter  $E(z) = az^2$ , then MB dist<sup>n</sup> lead to result that avg. no. of particles within a range  $\Delta z$  of the parameter is given by

$$n(z) dz = A e^{-\left(\frac{E(z)}{kT}\right)} dz$$

[assuming  $g(z) = 1$ ]

e.g. velocity components.

$$N = \int n(z) dz$$

$$= \int A e^{-\frac{E(z)}{kT}} dz = \int A e^{-\frac{az^2}{kT}} dz$$

$$\text{Total Energy } E = \int E(z) dN = \int A az^2 e^{-\frac{az^2}{kT}} dz$$

$$\Rightarrow \bar{E}(z) = \frac{E(z)}{N} = \frac{\int A az^2 e^{-\frac{az^2}{kT}} dz}{\int A e^{-\frac{az^2}{kT}} dz} = \frac{1}{2} kT$$

That is, for every degree of freedom for which above conditions are fulfilled, the mean energy per particle, in an assembly in equilibrium at temp T, is  $(\frac{kT}{2})$

$$\text{eg. } U(x) = \frac{1}{2} kx^2$$

$$E(2x) = \frac{1}{2} m v_x^2$$

① Not fulfilled for  $E(y) = my$

② Not fulfilled for energies at quantum level that take discrete values and considering energy as a continuous function is erroneous. For eg. rotational motion.

## ✓ Expansivity and Elasticity

$$\textcircled{O} \quad b = b(V, T)$$

$$\Rightarrow db = \left(\frac{\partial b}{\partial V}\right)_T dV + \left(\frac{\partial b}{\partial T}\right)_V dT$$

We know  $\alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T}\right)_P$ , : Coefficient of volume expansion or expansivity

$$\text{Also } k = -V \left(\frac{\partial P}{\partial V}\right)_T$$

Isothermal Elasticity


$$\Rightarrow db = -\left(\frac{k}{V}\right) dV + \left(\frac{\partial k}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_P dT$$

$$= -\left(\frac{k}{V}\right) dV + \frac{k\alpha}{V} dT$$

$$\boxed{db = -\frac{k}{V} dV + k\alpha dT}$$

~~Q~~ A reversible process is an ideal process & can never be achieved in practice. Explain

~~Q~~ Consider a gas in cylinder. We wish to change the volume of gas from  $V$  to  $2V$ .



1st Way

Rapid Movement of piston from A to B.

This process is irreversible as all intermediate states are non equilibrium states.

2nd Way

Piston is spring loaded s.t. it remains at A when weight  $W$  is put on top and at once moves to B as soon as  $W$  is removed. But even now process is irreversible

3rd Way

Now we can place 2 weights ( $\frac{W}{2}$  each). On removing 1 weight, intermediate equilibrium state C is reached.

So if use  $n$  weights ( $\frac{W}{n}$  each), we can realize  $(n-1)$  equilibrium states between A and B. Thus if  $n$  is made very large ( $n \rightarrow \infty$ ), we can have equilibrium states that do not differ much from each other.

This obviously is not practicable.

The problem of non-realizability of reversible process in thermodynamics is overcome by quasi-static process i.e. process is worked out very slowly.

Q1 A block of copper whose expansivity is  $4.8 \times 10^{-6} \text{ K}^{-1}$  and isothermal elasticity is  $1.30 \times 10^{11} \text{ Nm}^{-2}$  is at atm. pressure and at a temp. of  $0^\circ\text{C}$ . Its temp. is raised to  $10^\circ\text{C}$ . Calculate final pressure when

(a) Volume is kept const.

(b) volume changes from 1000 cc. to 1000.1cc.

At let  $p = p(V, T)$

$$\Rightarrow dp = \left(\frac{\partial p}{\partial V}\right)_T dV + \left(\frac{\partial p}{\partial T}\right)_V dT$$

$$\text{Now given } \alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T}\right)_P = 4.8 \times 10^{-6} \text{ K}^{-1}$$

$$k_T = -V \left(\frac{\partial P}{\partial V}\right)_T = 1.3 \times 10^{11} \text{ N/m}^2$$

$$\Rightarrow dp = -\frac{k_T}{V} dV + \frac{\left(\frac{\partial V}{\partial T}\right)_P}{\left(\frac{\partial p}{\partial V}\right)_T} dT$$

$$= -\frac{k_T}{V} dV + \alpha V \cdot \left(\frac{\partial p}{\partial V}\right)_T dT$$

$$= -\frac{k_T}{V} dV + \alpha V \frac{k_T}{V} dT$$

$$\Rightarrow \boxed{dp = -\frac{k_T}{V} dV + \alpha k_T dT}$$

(a)  $dV = 0 \Rightarrow dp = \alpha k_T \cdot (10)$

$$\Rightarrow p_2 = p_1 + \alpha k_T \cdot 10 = 10^5 + 624 \times 10^5 = 625 \text{ atm}$$


(b)  $dV \neq 0 \Rightarrow dp = -\frac{1.3 \times 10^{11}}{1000 \times 10^5} \times 0.1 + 625 \text{ atm}$

$$\Rightarrow p_2 = 495 \text{ atm}$$

When specimen is allowed to expand, Pressure is very less


Q1 A Carnot engine is operated b/w ice point & steam point.

(a) If engine receives 746 calories from hot reservoir in each cycle, how many calories does it reject to cold reservoir? (b) If engine is operated as a refrigerator and receives the same heat from cold reservoir, how many calories does it deliver to hot reservoir. (c) How much work is done by engine in each case?



Heat Engine

Fig (a)



Refrigerator

Fig (b)

(a) Refer to Fig (a)

For Carnot cycle,  $1 - \frac{Q_2}{Q_1} = 1 - \frac{T_2}{T_1}$

$$\Rightarrow 1 - \frac{Q_2}{746} = 1 - \frac{273}{373} \Rightarrow Q_2 = \underline{\underline{546}} \text{ calories}$$

$W = \underline{\underline{200}} \text{ cal}$

(b) Refer to Fig (b)

For Carnot refrigerator, again  $\frac{Q_2}{Q_1} = \frac{T_2}{T_1}$

$$\Rightarrow Q_1 = \frac{Q_2 \cdot T_1}{T_2} = \frac{746 \times 373}{273} = \underline{\underline{1019}} \text{ calories}$$

(same formula holds as cycle is just reversed)

$W = \underline{\underline{-273}} \text{ cal}$




Note that in both cases,  $Q_1$  is more


Q1 Calculate during the cycle

a) Work Done

b)  $\eta$



A1



a) Work Done = Area of PV curve =  $(p_2 - p_1)(v_2 - v_1)$

b)  $\eta = \frac{W}{Q_{AB} + Q_{BC}}$

$$Q_{AB} = C_V dT = \frac{C_V}{R} (p_2 - p_1) v_1$$

$$Q_{BC} = C_P dT = \frac{C_P}{R} (v_2 - v_1) p_2$$

$$\Rightarrow \eta = \frac{(p_2 - p_1)(v_2 - v_1)}{\left(\frac{1}{r_1}\right)(p_2 - p_1)v_1 + \left(\frac{r}{r_1}\right)(v_2 - v_1)p_2}$$


$$= \frac{r-1}{\left(\frac{v_1}{v_2 - v_1}\right) + \left(\frac{r p_2}{p_2 - p_1}\right)}$$

$$\left(\frac{v_1}{v_2 - v_1}\right) + \left(\frac{r p_2}{p_2 - p_1}\right)$$

Q1 Calculate  $\eta$  in terms of  $(p_a, p_b, r)$

$$\text{A1 } \eta = 1 - \frac{Q_{da}}{Q_{bc}} = 1 - \left(\frac{T_d - T_A}{T_c - T_B}\right)$$

$$\left(\frac{T_d}{T_c}\right) = \left(\frac{p_a}{p_b}\right)^{\frac{r-1}{r}} ; \left(\frac{T_A}{T_B}\right) = \left(\frac{p_a}{p_b}\right)^{\frac{r-1}{r}} \Rightarrow \eta = 1 - \left(\frac{p_a}{p_b}\right)^{\frac{r-1}{r}}$$



Q) An ideal gas through a cycle consisting of

a) isochoric, adiabatic and isothermal lines.

When isothermal occurs at minimum Temperature, calculate the  $\eta$  if absolute Temperature varies  $n$  fold within the cycle.

Ans By the given conditions, fig.1 represents the appropriate cycle.

$$\begin{aligned}\eta &= 1 - \frac{\Omega_2}{\Omega_1} \\ &= 1 - \frac{RT \ln \left( \frac{V_C}{V_A} \right)}{C_V (nT - T)}\end{aligned}$$




Fig 1

$$= 1 - \frac{(r-1)}{(n-1)} \ln \left( \frac{V_C}{V_A} \right) = 1 - \frac{(r-1)}{(n-1)} \ln(n)^{\frac{1}{r-1}} = 1 - \frac{\ln n}{(n-1)}$$

$$\left[ T_B V_B^{r-1} = T_C V_C^{r-1} \Rightarrow nT \cdot V_A^{r-1} = T \cdot V_C^{r-1} \Rightarrow \left( \frac{V_C}{V_A} \right) = n^{\frac{1}{r-1}} \right]$$

Q) Calculate pressure of air in vessel being evacuated, as a function of evacuation time  $t$ . Volume of vessel is  $V$  and initial pressure is  $P_0$ . Process is assumed to be isothermal and evacuation rate (i.e. Volume being evacuated per second) is  $C$ .  $C$  is independent of Pressure.

Ans Now the number of molecules are changing.


$$PV = nRT \Rightarrow PdV + Vdp = RT dn \Rightarrow Vdp = RT dn$$

$$\text{Also } dn = -n \frac{Cd t}{V}$$

$$\Rightarrow Vdp = -\frac{RT}{V} n C dt \Rightarrow \int \frac{dp}{p} = -\int \frac{C}{V} dt$$

$$\Rightarrow \ln \left( \frac{p}{P_0} \right) = -\frac{Ct}{V} \Rightarrow \boxed{P = P_0 e^{-\frac{Ct}{V}}}$$

Q1 Calculate Efficiency of the cycle



A1 AB  $\rightarrow dQ > 0$  as  $T dS > 0$


BC  $\rightarrow dQ_2 < 0$  as  $T dS < 0$

CA  $\rightarrow dQ = 0$  : Adiabatic Process

$$\eta = \frac{Q_1 - Q_2}{Q_1} = \frac{\left(\frac{1}{2}\right) \times (T_2 - T_1) (S_2 - S_1)}{\frac{1}{2} \times (T_2 + T_1) (S_2 - S_1) + T_1 (S_2 - S_1)} = \left(\frac{T_2 - T_1}{T_2 + T_1}\right)$$

A1 Prove Clausius Inequality for refrigeration cycles.

$$\eta_{\text{rev}} \geq \eta_{\text{irr}} \Rightarrow \left(\frac{Q_2}{Q_1 - Q_2}\right)_{\text{rev}} \geq \left(\frac{Q_2}{Q_1 - Q_2}\right)_{\text{irr.}}$$



$$\Rightarrow \left(\frac{1}{\frac{Q_1}{Q_2} - 1}\right)_{\text{rev}} \geq \left(\frac{1}{\frac{Q_1}{Q_2} - 1}\right)_{\text{irr.}} \Rightarrow \left(\frac{1}{\frac{T_1}{T_2} - 1}\right)_{\text{rev}} \geq \left(\frac{1}{\frac{T_1}{T_2} - 1}\right)_{\text{irr.}}$$

A1 (For same temperature limits)

$$\Rightarrow \frac{Q_1}{Q_2} \geq \frac{T_1}{T_2} \Rightarrow \frac{Q_1}{T_1} \geq \frac{Q_2}{T_2} \Rightarrow \boxed{\frac{Q_2}{T_2} + \left(-\frac{Q_1}{T_1}\right) \leq 0}$$

Q1 m gram of water at temperature  $T_1$  is isobarically and adiabatically mixed with equal amount of water at temperature  $T_2$ . Show  $\Delta S = 2mC_p \ln\left(\frac{T_{\text{avg}}}{\sqrt{T_1 T_2}}\right)$

A1 Now  $dQ_1 = dQ_2 \Rightarrow mC_p(t - T_1) = mC_p(T_2 - t)$   
 $\Rightarrow t = T_{\text{avg}} = \left(\frac{T_1 + T_2}{2}\right)$

$$\begin{aligned} \Rightarrow \Delta S &= \Delta S_1 + \Delta S_2 \\ &= mC_p \ln\left(\frac{t}{T_1}\right) + mC_p \ln\left(\frac{t}{T_2}\right) \\ &= mC_p \ln\left(\frac{t^2}{T_1 T_2}\right) = 2mC_p \ln\left(\frac{T_{\text{avg}}}{\sqrt{T_1 T_2}}\right) \end{aligned}$$

(Q) A heat engine receives 100 kcal of heat from a source at 1000 K. It ejects 50 kcal, 75 kcal and 25 kcal of heat at 500 K to surroundings. Investigate nature of change.

A1 To know nature of changes we have Clausius Inequality with us.

$$(a) \int \frac{d\theta}{T} = \frac{100}{1000} - \frac{50}{500} = 0 \quad : \text{reversible process}$$

$$(b) \int \frac{d\theta}{T} = \frac{100}{1000} - \frac{75}{500} = -0.05 \text{ kcal K}^{-1} : \text{irreversible process}$$

$$(c) \int \frac{d\theta}{T} = \frac{100}{1000} - \frac{25}{500} = 0.05 \text{ kcal K}^{-1} : \text{CANNOT OCCUR.}$$

Note that in 3rd case efficiency is more than 50% which is violation of Carnot's Theorem.

[Carnot Theorem puts a theoretical limit on efficiency which can be obtained by Clausius Inequality.]

Q2 Consider a system at Temperature  $T_s$  in thermal contact with a heat bath at  $T_h$  ( $T_s < T_h$ ). If heat capacity of system remains const. over interval  $(T_s, T_h)$ . show that temperature equalization is accompanied by increase in entropy.

A1 Final temperature =  $T_h$  When we say entropy, we say that of universe.....

$$dS_{\text{system}} = \int \frac{d\theta}{T} = \int_{T_s}^{T_h} C_s \cdot \frac{dT}{T} = \underline{\underline{C_s \ln \left( \frac{T_h}{T_s} \right)}}$$

$$dS_{\text{surrounding}} = \int \frac{d\theta}{T} = -C_s \frac{(T_h - T_s)}{T_h} = -C_s \left( 1 - \frac{T_s}{T_h} \right)$$

$$dS_{\text{Total}} = C_s \left[ \ln \left( \frac{T_h}{T_s} \right) + 1 + \left( \frac{T_s}{T_h} \right) \right]$$

Now,  $\left[ \ln x + \left( \frac{1}{x} \right) > 1 \right] \forall x$  [irrespective of hot or cold bath]

Hence  $dS_{\text{Total}} > 0$

$$\ln x + \left( \frac{1}{x} \right)$$



Q Prove that entropy always increases

A1 Here we have 2 cases

① Isolated system : from Clausius inequality  $\Delta S \geq 0$

② Heat from body at  $T_1$  to body at  $T_2$

$$\Rightarrow dS = \frac{dQ}{T_2} - \frac{dQ}{T_1} = dQ \left[ \frac{1}{T_2} - \frac{1}{T_1} \right]$$

since heat flows from higher temp to lower temp.

$$\Rightarrow \frac{1}{T_2} \geq \frac{1}{T_1} \Rightarrow dS \geq 0$$

(placed at 300k)

Q 10A current is passed through insulated resistor of  $25\Omega$  for 1s. Find  $\Delta S_{\text{res}}$  and  $\Delta S_{\text{univ}}$ . if  $m = 10g$  and  $C_p = 0.84 \text{ J g}^{-1} \text{ K}^{-1}$

A1 Since resistor is insulated  $\Rightarrow \Delta S_{\text{sur}} = 0 \Rightarrow \Delta S_{\text{univ}} = \Delta S_{\text{res}}$

$$\text{Also } \Delta Q = m C_p \Delta T \Rightarrow 100 * 25 * 1 = 10 * 0.84 * \Delta T$$

$$\Rightarrow \Delta T = 297.61 \Rightarrow T_2 = 597.6 \text{ K}$$

$$\Rightarrow \Delta S_{\text{res}} = \int \frac{dQ}{T} = m C_p \int \frac{dT}{T} = 8.4 \times \ln \left( \frac{T_2}{T_1} \right) \\ = 5.78 \text{ J/K}$$

Q Calculate  $\Delta S_{\text{univ}}$  if  $1 \mu\text{F}$  capacitor is connected to  $100\text{V}$  reversible cell @  $0^\circ\text{C}$ .

A1 Since Heat is not involved in charging the capacitor  $dQ = 0$  and  $dS = 0$

Q 1 mole of He and 1 mole of  $\text{H}_2$  are mixed at a const. Temperature. Calculate entropy change.

A1 Assuming isothermal process and  $V_1 = V_2 = V_0$

$$\Delta S = \Delta S_{\text{H}_2} + \Delta S_{\text{He}} \\ = 2R \ln \left( \frac{V_f}{V_i} \right) = 2 \times 2 \text{ cal/K/mol} \times \ln 2 \\ = 2.77 \text{ cal/K}$$



$$\text{Q1) Show that } C_p = T \left( \frac{\partial V}{\partial T} \right)_P + \left( \frac{\partial k}{\partial T} \right)_S$$

$$\text{A1) By definition } C_p = \left( \frac{\partial Q}{\partial T} \right)_P = T \left( \frac{\partial S}{\partial T} \right)_P$$

To convert '1' differential into '2', we know cyclic rule

$$\left( \frac{\partial S}{\partial T} \right)_P \left( \frac{\partial T}{\partial P} \right)_S \left( \frac{\partial P}{\partial S} \right)_T = -1 \Rightarrow \left( \frac{\partial S}{\partial T} \right)_P = - \left( \frac{\partial P}{\partial T} \right)_S \left( \frac{\partial S}{\partial P} \right)_T$$

$$\Rightarrow C_p = -T \left( \frac{\partial P}{\partial T} \right)_S \left( \frac{\partial S}{\partial P} \right)_T = -T \left( \frac{\partial P}{\partial T} \right)_S * \left[ - \left( \frac{\partial V}{\partial T} \right)_P \right]$$

$$\Rightarrow C_p = T \left( \frac{\partial P}{\partial T} \right)_S \left( \frac{\partial V}{\partial T} \right)_P$$

P V S T

~~Q1)~~ A Gas obey equation  $[P(V-b) = RT]$  where  $b$  is a const.

Gas has const. value of  $C_V$ . Show that

(a)  $U$  is a function of ' $T$ ' only

(b) During adiabatic process, gas follows  $b(V-b)^{\gamma} = \text{const}$

A1) We will prove 1<sup>st</sup> part by contradiction.

$$\text{Let } U = f(V)$$

$$\Rightarrow \left( \frac{\partial U}{\partial V} \right)_T \neq 0 \quad \underline{\underline{\text{L.H.S.}}} = \left( T \frac{ds}{dv} - P dv \right)_T = T \left( \frac{\partial s}{\partial v} \right)_T - P$$

$$= T \cdot \left( \frac{\partial P}{\partial T} \right)_V - P = T \cdot \frac{R}{(V-b)} - P = P - P = 0 \neq \underline{\underline{\text{R.H.S}}}$$

Hence contradicted.

$$\text{Similarly let } U = f(P)$$

$$\Rightarrow \left( \frac{\partial U}{\partial P} \right)_T \neq 0 \quad \underline{\underline{\text{L.H.S.}}} = T \left( \frac{\partial S}{\partial P} \right)_T - P \left( \frac{\partial V}{\partial P} \right)_T$$

$$= -T \cdot \left( \frac{\partial V}{\partial T} \right)_P - P \left( \frac{\partial V}{\partial P} \right)_T = -T \cdot \frac{R}{P} - P \cdot \left( \frac{V-b}{P} \right)$$

$$= -\frac{RT}{P} + (V-b) = -(V-b) + (V-b) = 0 \neq \underline{\underline{\text{R.H.S}}}$$

Hence Contradicted  $\Rightarrow \boxed{U = f(T) \text{ only}}$

$$(ii) p(V-b) = RT$$

$$\text{Now } C_p - C_v = T \left( \frac{\partial P}{\partial T} \right)_V \left( \frac{\partial V}{\partial T} \right)_P = T \cdot \frac{R}{(V-b)} \cdot \frac{R}{b}$$

$$= \frac{T \cdot R^2}{RT} = R$$

$$\text{Also, } \frac{C_p}{C_v} = r$$

$$\Rightarrow \boxed{C_v = \left( \frac{R}{r-1} \right)}$$

Also during adiabatic process

$$dQ = 0 \Rightarrow dU + PdV = 0$$

$$\Rightarrow C_v dT + PdV = 0 \quad (\text{now we know } U = f(T) \text{ only})$$

$$\Rightarrow dV = C_v dT$$

$$\Rightarrow \frac{R}{r-1} \cdot dT + \frac{RT}{(V-b)} dV = 0$$

$$\Rightarrow \frac{dT}{T} + (r-1) \frac{dV}{(V-b)} = 0$$

$$\Rightarrow \ln T + (r-1) \ln (V-b) = A$$

$$\Rightarrow T(V-b)^{r-1} = A,$$

$$\Rightarrow \frac{p(V-b)}{R} (V-b)^{r-1} = A$$

$$\Rightarrow \boxed{p(V-b)^r = \text{Const}}$$

 A system undergoes phase transition at a const. temp T and pressure P. If it suffers a change  $\Delta V$  in its volume, show that  $\Delta U = T \left[ \frac{\partial}{\partial T} \left( \frac{p}{T} \right)_V \right] \Delta V$

$$\text{A/} \text{ let } U = f(T, V)$$

$$\Rightarrow dU = \left( \frac{\partial U}{\partial T} \right)_V dT + \left( \frac{\partial U}{\partial V} \right)_T dV = \left( \frac{\partial U}{\partial V} \right)_T dV = \left[ T \left( \frac{\partial S}{\partial V} \right)_T + P \right] \Delta V$$

$$= \left[ T \left( \frac{\partial P}{\partial T} \right)_V + P \right] \Delta V = \underline{\underline{T^2 \frac{\partial}{\partial T} \left( \frac{p}{T} \right)_V \Delta V}}$$

Q/ The coefficient of viscosity of He is  $2 \times 10^{-5}$  poise at 27°C. Calculate the diameter of He molecule.

$$N_A = 6 \times 10^{23} \text{ mol}^{-1}; k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}, A[\text{He}] = 4$$

All From Transport Phenomenon.

$$\eta = \frac{1}{3} mn \bar{v} \lambda = \frac{1}{3} m n \bar{v} \frac{1}{\sqrt{2} \pi d^2 n} = \frac{m \bar{v}}{3 \sqrt{2} \pi d^2}$$

$$\Rightarrow d^2 = \left( \frac{mn\bar{v}}{3\sqrt{2} \pi \eta} \right)$$

Now  $m = \frac{4g}{6 \times 10^{23}} = 6.6 \times 10^{-24} \text{ g} = \underline{\underline{6.6 \times 10^{-27} \text{ kg}}}$

$$\bar{v} = \sqrt{\frac{8kT}{\pi m}} = \sqrt{\frac{8 \times 1.38 \times 10^{-23} \times 300}{3.14 \times 6.6 \times 10^{-27}}} = \underline{\underline{\sqrt{159.81} \times 10^2}} \\ = \underline{\underline{1264 \text{ m s}^{-1}}}$$

$$\Rightarrow d^2 = \frac{6.6 \times 1264 \times 10^{-27}}{3\sqrt{2} \times 3.14 \times 2 \times 10^{-5}} = 313.1 \times 10^{-22} = \underline{\underline{}}$$

$$\Rightarrow \boxed{d = 1.769 \text{ \AA}}$$

Also note that  $\underline{\underline{mn}} = \rho$

Q/ Calculate the thermal conductivity of air using the following data:  $\rho = 1.29 \text{ kg m}^{-3}$ ,  $\bar{v} = 460 \text{ m s}^{-1}$ ,  $\lambda = 6.4 \times 10^{-8} \text{ m}$ ,  $M = 29$ ,  $r = 1.4$  and  $R = 8.31 \text{ kJ kmol}^{-1} \text{ K}^{-1}$

All We know  $k = \frac{1}{3} mn \bar{v} \lambda \frac{C_V}{C_V + R}$

$$mn = \rho = 1.29$$

$$\bar{v} = 460$$

$$\lambda = 6.4 \times 10^{-8}$$

$$C_V = \frac{R}{M} = \frac{8.31}{29} = 0.283 \text{ J K}^{-1} \text{ mol}^{-1} = 9.06 \times 10^{-3} \text{ W m}^{-1} \text{ K}^{-1}$$

$$\Rightarrow \boxed{k = 9.06 \times 10^{-3} \text{ W m}^{-1} \text{ K}^{-1}}$$

Q1 The coefficient of viscosity and diffusion for O<sub>2</sub> are  $1.95 \times 10^{-5}$  N s m<sup>-2</sup> and  $1.22 \times 10^{-5}$  m<sup>2</sup> s<sup>-1</sup>. If avg. molecular speed is 440 ms<sup>-1</sup>, calculate density and mean free path of molecules.

A)  $\eta = \frac{1}{3} mn \bar{v} \lambda$

$$D = \frac{1}{3} \bar{v} \lambda$$

$$\Rightarrow \rho = mn = \frac{\eta}{D} = 1.6 \text{ kg/m}^3$$

$$\lambda = \frac{3D}{\bar{v}} = \frac{3 \times 1.22 \times 10^{-5}}{440} \\ = 8.32 \times 10^{-8} \text{ m}$$

Q2 Calculate mass of N<sub>2</sub> diffusing through an area of  $10^{-2} \text{ m}^2$  in 10 s if the concentration gradient is  $1.26 \text{ kg m}^{-4}$ , the mean free path =  $10^{-7} \text{ m}$  and  $\bar{v} = 476 \text{ ms}^{-1}$ .

A)  $D = \frac{1}{3} \bar{v} \lambda = \frac{dN}{A \cdot t \cdot \left( \frac{dn}{dz} \right)}$

$$\Rightarrow dN = \frac{1}{3} \cdot \bar{v} \cdot \lambda \cdot A \cdot t \left( \frac{dn}{dz} \right)$$

$$\Rightarrow dm = \frac{1}{3} \cdot \bar{v} \cdot \lambda \cdot A \cdot t \left[ m \left( \frac{dn}{dz} \right) \right] = \frac{1}{3} \bar{v} \lambda A t \left( \frac{dp}{dz} \right)$$

$$= \frac{1}{3} \times 476 \times 10^{-7} \times 10^{-2} \times 10 \times 1.26 = 1.99 \times 10^{-6} \text{ kg}$$

Q3 What fraction of volume must liquid ether occupy at room temperature in order to pass into critical state when critical temperature is reached? For ether T<sub>c</sub> = 467 K; P<sub>c</sub> = 35.5 atm; density = 720 kg/m<sup>3</sup> & A = 74

A)  $T_c = \frac{8a}{27Rb} = \frac{8b}{R} \left( \frac{a}{27b^2} \right) \Rightarrow \frac{8b}{R} = 13.64 \times 10^{-5} \text{ m}^3$   
 $\Rightarrow V_c = 3b = \frac{40.94 \times 10^{-5} \text{ m}^3}{\frac{RT_c}{P_c V_c}} = \frac{8}{3}$

Now since it's in critical stage  
volume of vessel =  $V_c = 40.94 \times 10^{-5} \text{ m}^3$

Also since V<sub>c</sub> is defined for 1 mole, for liquid ether  
(before evaporation), Volume =  $\frac{74 \times 10^{-3}}{720 \text{ kg/m}^3} = 10.27 \times 10^{-5} \text{ m}^3$

$\Rightarrow$  Fraction occupied = 25%

अद्वैत मर्या !! Constants का देखने रखना

Q) Prove  $\left(\frac{\partial T}{\partial V}\right)_P = - \left(\frac{\partial P}{\partial S}\right)_T$

All From  $P \quad V \quad S \quad T$

if we see  $\left(\frac{\partial T}{\partial V}\right)_S = - \left(\frac{\partial P}{\partial S}\right)_V$

1<sup>st</sup> Maxwell

नोट कि Constants तो ज्ञात हैं !!

Therefore instead of Order of numerator and denominator,  
we can look for const.

For ex. if 4<sup>th</sup> const if  $P \Rightarrow$  denominator of RHS is  $P$

$\Rightarrow$  numerator of LHS is  $V$

$\Rightarrow$  equation to use is

$$\left(\frac{\partial V}{\partial S}\right)_P = \left(\frac{\partial T}{\partial P}\right)_S \quad \text{OR} \quad \left(\frac{\partial V}{\partial T}\right)_P = - \left(\frac{\partial S}{\partial P}\right)_T$$

We can see it is 2<sup>nd</sup> one


$$\left(\frac{\partial V}{\partial T}\right)_P = - \left(\frac{\partial S}{\partial P}\right)_T \quad : 4^{\text{th}} \text{ Maxwell Equation}$$

Remember


$$\left(\frac{\partial A}{\partial B}\right)_C$$

A and C are neighbours like PV  
or ST

# OPAMP



## Ideal Opamp




$$V_C = A(V_A - V_B)$$

$$A \rightarrow \infty \Rightarrow V_A = V_B$$

②  $R_{in} = \infty \Rightarrow I_A = I_B = 0$

$$\left[ \frac{V_A - V_B}{R_{in}} \right]$$

③  $R_{out} = 0$




## Opamp Circuit

### AMPLIFIER

#### a) Inverting


$$V_o = -\frac{V_{in}}{R_1} \cdot R_f$$




$$R_{in} = R_1$$

$$R_{out} = 0$$

If  $R' \ll R$  and  $V_o = 0$ ,  $R_{out} = R_1$




b) Non Inverting



$$V_o = V_{in} \left[ \frac{R_f}{R_1} + 1 \right]$$

$R_{in} = \infty$   
 $R_{out} = 0$


**INTEGRATOR**



$$V_o = - \frac{V_i}{R C s}$$

$R_{in} = R$   
 $R_{out} = 0$

**DIFFERENTIATOR**




$$V_o = - \frac{V_{in} R}{(1/Cs)}$$

$$= - V_{in} R C s$$

$R_{in} = \infty$   
 $R_{out} = 0$

5

Since molecules are not point particles, they cover some volume and the space available to molecules is less than the volume of container.



Sphere of influence of each molecule is of radius  $2r$  and since this is shared by 2 interacting molecules, value of  $b$  is half of total no. of molecules times the volume of sphere of influence

$$b = \frac{1}{2} * N * \frac{4\pi}{3} (2r)^3 = \underline{4Nv}$$

### Claude's Virial Theorem

Assuming that particles obey Newtonian mechanics.

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = \vec{p}_i$$

$$\text{define } \phi = \sum_i \vec{p}_i \cdot \vec{r}_i$$

$\vec{p}_i$ : momentum

$$\begin{aligned} \left( \frac{d\phi}{dt} \right) &= \sum \vec{p}_i \cdot \vec{\dot{r}}_i + \sum \vec{p}_i \cdot \vec{r}_i = \sum \vec{F}_i \cdot \vec{r}_i + \sum m_i \vec{v}_i \cdot \vec{r}_i \\ &= \sum \vec{F}_i \cdot \vec{r}_i + 2T \end{aligned}$$

Taking time average of  $\left( \frac{d\phi}{dt} \right)$

$$\langle \frac{d\phi}{dt} \rangle = \frac{1}{2} \int_0^T \left( \frac{d\phi}{dt} \right) dt = \frac{1}{2} \int_0^T \sum \vec{F}_i \cdot \vec{r}_i dt + \frac{2}{2} \int_0^T T dt$$

$$\Rightarrow \frac{1}{T} (\phi_T - \phi_0) = \overline{\sum \vec{F}_i \cdot \vec{x}_i} + \overline{2T}$$

Choose  $T$  st.  $\phi_T = \phi_0$ , we get

$$\boxed{\sum \vec{F}_i \cdot \vec{x}_i = -2T}$$

L.H.S. is called Virial

- $F_i$  includes
- Force of impact on molecules due to collision with the walls of container
  - Force of collision between molecules
  - Force of interaction between molecules.


### Application

#### 1) Pressure due to ideal gas

Only 1<sup>st</sup> component of force is there.

$$\vec{F} \cdot \vec{x} = -p da \hat{n} \cdot \vec{x}$$

$$\text{Virial} = -\frac{1}{2} \overline{\sum \vec{F}_i \cdot \vec{x}_i}$$



$p$ : pressure

$$\begin{aligned} & \frac{1}{2} p \overline{\sum d\vec{s} \cdot \vec{x}_i} \\ &= \frac{1}{2} p \cdot \overline{\int \vec{x}_i \cdot d\vec{s}} \\ &= \frac{1}{2} p \cdot \int (\vec{\nabla} \cdot \vec{x}_i) dV \\ &= \frac{1}{2} p \cdot \overline{3V} \\ &= \frac{3}{2} p V \end{aligned}$$

$$[\vec{\nabla} \cdot \vec{x}_i = 3] \\ (\overline{3V} = 3V)$$

$$\Rightarrow \frac{3}{2} p V = \overline{T} = \frac{3}{2} RT \quad (\text{for 1 mole of gas})$$

$$\Rightarrow \boxed{pV = RT}$$

$$\& \boxed{p = \frac{2}{3} \frac{\overline{T}}{V}}$$

## Quasi static Process

If during state of non-equilibrium, if external forces be varied only slightly, so that resulting unbalanced forces are infinitesimally small, the process is assumed to be in equilibrium & is called Quasi static Process.

For quasistatic process,  $dW = P dV$

## Efficiency increase of Carnot engine

$$\eta = 1 - \left( \frac{T_2}{T_1} \right)$$

$$\left( \frac{d\eta}{dT_1} \right) = \left( \frac{T_2}{T_1^2} \right)$$

$$\left( \frac{d\eta}{d(-T_2)} \right) = \left( \frac{1}{T_1} \right)$$

$$\Rightarrow \left( \frac{d\eta}{d(-T_2)} \right) > \left( \frac{d\eta}{dT_1} \right) \Rightarrow \text{It's preferable to reduce } T_2 \text{ than to increase } T_1$$

## Internal vs External Combustion

- In steam engine, heat is produced in a separate boiler outside the cylinder : External Combustion Engine
- When heat is produced inside the cylinder by explosion of fuel, its called Internal Combustion Engine. It has much higher efficiency.

## Application of Clausius Inequality : Maximum Work

Q/2 bodies of identical mass and heat capacity C  
are at temp.  $T_1$  and  $T_2$ . What is the max. work  
that can be done?

Ans let final temp be  $T_f$


$$Q_1 = C(T_f - T_1)$$

$$Q_2 = C(T_f - T_2)$$

Now we cannot equate the two, as  $W_{done} \neq 0$

$$W = Q_1 - Q_2$$

$$= C[T_1 + T_2 - 2T_f]$$



For maximum work, we need to find out minimum value  
of  $T_f$ .

We know

$$\int \frac{dQ}{T} \leq 0$$

$$\Rightarrow \int_{T_1}^{T_f} -\frac{CdT}{T} + \int_{T_2}^{T_f} \frac{CdT}{T} \leq 0$$

~~$T_1$~~        ~~$T_2$~~


increase in entropy      decrease in entropy

$$\Rightarrow -C \ln \left( \frac{T_f}{T_1} \right) + C \ln \left( \frac{T_f}{T_2} \right) \leq 0$$

$$\Rightarrow C \ln \left[ \frac{T_f^2}{T_1 T_2} \right] \geq 0$$

$$T_f^2 \geq T_1 T_2$$

$$\Rightarrow W_{max} = C [T_1 + T_2 - 2\sqrt{T_1 T_2}]$$



Heat released by hot reservoir  
=  $-CdT$


Heat gained by cold reservoir  
=  $+CdT$

①



Find Entropy change after a long time.

Ans



For element  $dx$

$$T_i = T_1 - \left(\frac{T_1 - T_2}{L}\right)x$$

$$T_f = \frac{T_1 + T_2}{2} \quad (\text{for the whole rod})$$

$$dm = \frac{M}{L} \cdot dx$$

$$\begin{aligned} dS &= \frac{dQ}{T} = dm \cdot c \cdot \frac{dT}{T} = \frac{Mc}{L} dx \int_{T_i}^{T_f} \frac{dT}{T} \\ &= \frac{Mc}{L} dx \ln \left[ \frac{\frac{(T_1 + T_2)}{2}}{T_1 - \left(\frac{T_1 - T_2}{L}\right)x} \right] \end{aligned}$$



For whole rod

$$\begin{aligned} \Delta S &= \int dS \\ &= \int_0^L \frac{Mc}{L} \ln \left[ \frac{T_1 + T_2}{2T_1 - \left(\frac{T_1 - T_2}{L}\right)x} \right] dx \\ &= \int_0^L \frac{Mc}{L} \ln \left( \frac{T_1 + T_2}{2} \right) dx - \int_0^L \frac{Mc}{L} \ln \left( T_1 - \left(\frac{T_1 - T_2}{L}\right)x \right) dx \end{aligned}$$



IBP

$$\begin{aligned} &\int \ln \left( T_1 - \left(\frac{T_1 - T_2}{L}\right)x \right) \cdot 1 dx \\ &= \ln \left( T_1 - \left(\frac{T_1 - T_2}{L}\right)x \right) x \Big|_0^L - \int_0^L \frac{x}{T_1 - \left(\frac{T_1 - T_2}{L}\right)x} \left(\frac{T_1 - T_2}{L}\right) dx \end{aligned}$$

$$= \ln\left(\frac{T_2}{L}\right) - \int_0^L \left( \frac{x}{\frac{T_1 L}{T_1 - T_2}} - x \right) dx \quad \checkmark$$

$$= L \ln\left(\frac{T_2}{L}\right) - \int_0^L \frac{x - \left(\frac{L T_1}{T_1 - T_2}\right)}{\left(\frac{L T_1}{T_1 - T_2}\right) - x} dx + \frac{\left(\frac{L T_1}{T_1 - T_2}\right)}{x - \left(\frac{L T_1}{T_1 - T_2}\right)}$$

$$= L \ln\left(\frac{T_2}{L}\right) + L + \left(\frac{L T_1}{T_1 + T_2}\right) \ln\left(x - \frac{L T_1}{T_1 + T_2}\right) \Big|_0^L$$

⋮  
⋮  
⋮

$$\Rightarrow \langle E \rangle = \frac{E}{N}$$

$$= \frac{h\nu N_0 e^{-\frac{h\nu}{kT}}}{\left(1 - e^{-\frac{h\nu}{kT}}\right)^2} \cdot \frac{\left(\frac{N_0}{1 - e^{-\frac{h\nu}{kT}}}\right)}$$

$$= \left[ \frac{h\nu}{1 - e^{-\frac{h\nu}{kT}}} \right] e^{-\frac{h\nu}{kT}}$$

$$= \boxed{\frac{h\nu}{e^{\frac{h\nu}{kT}} - 1}}$$

$$\textcircled{O} \text{ To prove } \langle E \rangle = \left[ \frac{h\nu}{e^{\frac{h\nu}{kT}} - 1} \right]$$

\textcircled{1} Note that till time of Planck only Maxwell distribution was true.

\textcircled{2} Assume  $g(E) = 1 \quad \forall E$

Now let's suppose particles at ground level =  $N_0$ .

$$\Rightarrow N_1 = N_0 e^{-\frac{h\nu}{kT}}$$

$$N_2 = N_0 e^{-\frac{2h\nu}{kT}}$$

$$N_3 = N_0 e^{-\frac{3h\nu}{kT}}$$

:

$$\Rightarrow N = N_0 + N_1 + N_2 + \dots$$

$$= N_0 \left[ 1 + e^{-\frac{h\nu}{kT}} + e^{-\frac{2h\nu}{kT}} + \dots \right]$$

$$= \frac{N_0}{1 - e^{-\frac{h\nu}{kT}}}$$

Also

$$E = N_0 E_0 + N_1 E_1 + N_2 E_2 + \dots$$

$$= 0 + h\nu N_0 e^{-\frac{h\nu}{kT}} + 2h\nu N_0 e^{-\frac{2h\nu}{kT}} + 3h\nu N_0 e^{-\frac{3h\nu}{kT}}$$

$$e^{-\frac{h\nu}{kT}} E =$$

$$h\nu N_0 e^{-\frac{2h\nu}{kT}} + 2h\nu N_0 e^{-\frac{3h\nu}{kT}}$$

$$E \left( 1 - e^{-\frac{h\nu}{kT}} \right) = h\nu \left[ N_0 e^{-\frac{h\nu}{kT}} + N_0 e^{-\frac{2h\nu}{kT}} + \dots \right]$$

$$= h\nu \frac{N_0 e^{-\frac{h\nu}{kT}}}{1 - e^{-\frac{h\nu}{kT}}}$$


$$\Rightarrow E = \frac{h\nu N_0 e^{-\frac{h\nu}{kT}}}{(1 - e^{-\frac{h\nu}{kT}})^2}$$

## Concept of Negative Temperature

★ Temperature is defined as  $dS = \frac{dQ}{T}$

$$\Rightarrow T = \left( \frac{dQ}{dS} \right)$$

Negative Temperature if entropy drops on supplying heat.





if more heat supplied, more particles move up  $\Rightarrow$  entropy drops  $\Rightarrow$  negative temperature

★ When  $n_1 > n_0$ , we say that substance is at negative temperature.

$$N_1 = N_0 e^{-\left(\frac{\Delta E}{kT}\right)}$$

if  $N_1 > N_0$  and  $\Delta E > 0$   
 $\Rightarrow T$  is negative



- ★ Nuclear spin systems can be brought to negative temperature by radio frequency techniques.


### Claussius Clapeyron Equation from Gibbs Free Energy

For two substances in reversible equilibrium at constant temperature & pressure, the Gibbs function remains same.

Hence, for a change of phase (phase 1 to phase 2) at  $T_a$  and  $P_a$

the specific gibbs functions are equal for two states

$$g_a^{(1)} = g_a^{(2)}$$



Similarly for a phase change at  $(T + dT)$  ( $= T_b$ ) and  $(P + dP)$ , ( $= P_b$ ),

$$g_b^{(1)} = g_b^{(2)}$$

$$\text{or } g_a^{(1)} + dg^{(1)} = g_a^{(2)} + dg^{(2)}$$

$$\Rightarrow dg^{(1)} = dg^{(2)}$$

$$\Rightarrow -S^{(1)} dT + V^{(1)} dP = -S^{(2)} dT + V^{(2)} dP$$

$$\Rightarrow (S^{(2)} - S^{(1)}) dT = (V^{(2)} - V^{(1)}) dP$$

$$\Rightarrow \frac{dP}{dT} = \frac{S^{(2)} - S^{(1)}}{V^{(2)} - V^{(1)}} = \frac{L}{T(V^{(2)} - V^{(1)})}$$

Hence Claussius Clapeyron equation relates change in Pressure  $dP$  with change in Boiling / Melting Point.

## Negative Temperature in a magnetic field

Let a system with just 2 magnetic energy levels due to a field  $H$  & dipoles of magnetic moment  $\mu_B$ .


Particles aligned  $\parallel$  to  $\vec{H}$ :  $E_1 = -\vec{\mu}_B \cdot \vec{B} = -\mu_B B$

Particles aligned antiparallel to  $\vec{H}$ :  $E_2 = -\vec{\mu}_B \cdot \vec{B} = \mu_B B$

$$\text{No. of particles at } E_1 = N_1 = A e^{-\frac{E_1}{kT}}$$

$$\text{" , " , " , } E_2 = N_2 = A e^{-\frac{E_2}{kT}}$$

$$\text{Now, } \frac{N_2}{N_1} = e^{\left(\frac{E_1 - E_2}{kT}\right)} \Rightarrow \ln N_2 - \ln N_1 = \frac{E_1 - E_2}{kT}$$



$$\text{Now } E_2 > E_1$$

$$\text{and normally } N_1 > N_2$$

$$\text{hence } T > 0$$

However, if suddenly  $H$  is reversed.

$$E_1 \text{ becomes } \mu_B B$$

$$E_2 \text{ becomes } -\mu_B B$$

The particles take some time to realign & for that time  $T$  becomes negative.

## Bose Einstein Condensation

For Bose Einstein distribution, we calculate  $N$  as

$$N = \left[ \frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} \right] (kT)^{\frac{3}{2}} \int_0^\infty \frac{x^{\frac{3}{2}}}{e^{\frac{x}{kT}} - 1} dx$$

For that we took  $g(E) dx$  as  $\frac{2\pi V}{h^3} (2m)^{\frac{3}{2}} e^{\frac{E}{kT}} dE$ .  
 Note that we have given zero weight to  $E=0$  level.  
 This is not justified in quantum mechanics.

Hence we have to take care of  $N_g$  also.

$$\Rightarrow N = N_g + CV \int_0^\infty \frac{e^{\frac{E}{kT}} dE}{e^{\frac{E}{kT}} - 1}$$


$\underbrace{\hspace{10em}}$

$N_{ex}$  : particles with non zero energy

$$N_g = N - N_{ex} \quad \text{and} \quad \frac{N_{ex}}{N} = \left( \frac{T}{T_B} \right)^{\frac{3}{2}}$$

$$= N - N \left( \frac{T}{T_B} \right)^{\frac{3}{2}}$$

$$N_g = N \left[ 1 - \left( \frac{T}{T_B} \right)^{\frac{3}{2}} \right] \checkmark$$



## ${}^4\text{He}$ Bose Einstein Condensate

For  ${}^4\text{He}$   $T_B = 3.13 \text{ K}$  (from formula)

Actual value =  $2.17 \text{ K}$

$$N_{\text{else}} = \left[ \frac{2\pi V}{h^3} (2m)^{3/2} \cdot \int_0^\infty \frac{x^{1/2}}{e^{kT_B} - 1} dx \right] (kT)^{3/2}$$


$$= \alpha (kT)^{3/2} \quad (\text{say})$$

Define  $\alpha = \frac{N}{(kT_B)^{3/2}}$ , we get

$$N_{\text{else}} = N \left( \frac{T}{T_B} \right)^{3/2}$$

$$\Rightarrow N = N_g + N_{\text{else}}$$

$$\Rightarrow N_g = N \left[ 1 - \left( \frac{T}{T_B} \right)^{3/2} \right]$$



① In Free expansion of an ideal gas, Temperature remains same

Worth while



Applicable

At atmospheric pressure, helium condenses into a normal liquid at 4.2 k. As the temperature is lowered further, it undergoes phase transition once again at 2.17 k. One would normally expect helium to solidify at this transition. Instead, liquid He changes into another liquid of vastly different properties.

Transition is accompanied by an anomalous behaviour of heat capacity. The heat capacity has a shape similar to Greek letter  $\lambda$ . Hence transition temperature is also referred to as LAMBDA POINT.

The new fluid has

- apparently zero viscosity
- infinite thermal conductivity.
- no bubbles.



Lack of viscosity means that superfluid can flow through smallest of cracks without impedance.

Another effects that superfluid Helium shows are that of fountain effect & second sound.

## Gibbs Phase Rule

Let us consider a system of  $c$  components in  $\beta$  phases.

For a multicomponent system,

$$G = G(P, T, x_1, x_2, x_3, \dots, x_c)$$

for every phase - ①

where  $P$ : Pressure of system

$T$ : Temperature of the system.

Mole Fractions  $x_j = \left( \frac{N_j}{N} \right)$  sum to unity, hence only  $(c-1)$  mole fractions are independent, thereby making the no. of variable unknown equal to  $2 + \beta(c-1)$


Note that we can consider, each separate phase as a simple system and the given system as a composite system of the different simple systems.

eg



liquid system

+



gaseous system

+



Solid system

At the condition of equilibrium, for example, if water converts to steam, the chemical potentials

$$\mu_{\text{water}}^L(T, P, x_i^L) = \mu_{\text{steam}}^G(T, P, x_i^G) \quad \text{--- ②}$$

Similarly for every component one for every phase

We will have,

$c(p-1)$  conditions or equation similar to (2)

( $p-1$ ) because,  $\mu_s = \mu_L$  and  $\mu_L = \mu_g$  gives  $\mu_s = \mu_g$   
hence 2 equations for 3 phases are independent

Total no. of variables which can be arbitrarily

$$\text{Assigned} = f = 2 + p(c-1) - c(p-1)$$

= degrees of freedom

$$= \underline{\underline{2 + c - p}}$$

This Phase Rule applies a lot of restriction on the  
 $1^{\text{st}}$  Order Phase Transitions.

e.g. (1) 5 Phases cannot coexist for a 2 component system.

(2) For a 2 components, 3 phase system at a  $(T, P)$   
variables are  $3(1) = 3$   
equations are  $2(2) = 4$   
hence no solution would be possible in general;  
thereby indicating that we are not free to  
specify  $P$  and  $T$  a priori.

It is possible to select both  $T$  and  $P$  arbitrarily  
and then to find a 2-phase state, a 3-phase  
state can exist only for a particular Temperature  
if Pressure is satisfied.

③ For single component system,

$$F = 2 + C - P = 3 - P$$

For one phase  $F = 2$  i.e. one can vary both  $T$  and  $P$ , hence system is called bivariant.

For 2 phases  $F = 1$ , i.e. one can vary only 1 variable either  $P$  (then  $T$  will be specific) or vice versa.

hence system is called monovariant.

For 3 phases to coexist in equilibrium  $F = 0$  i.e. all variables need to be fixed, only then all 3 phases can coexist in equilibrium. Such a system is called Invariant  $\Rightarrow$  Triple Point of water is unique.

### Blackbody Radiation

① A blackbody is defined as a body that absorbs all radiations that fall on the surface. Actual blackbodies do not exist in nature - though approximated by a hole in a box filled with highly absorptive material.

② According to Stefan Boltzmann's law, radiation energy per unit time is given as

$$q = \sigma T^4 A$$

A: area of emitting body  
T: Temperature of body's surface

$$\sigma: 5.6 \times 10^{-8} \text{ W/m}^2 \text{ K}^4$$

③ For gray bodies,  $q = \epsilon \sigma T^4 A$   $\epsilon$ : emissivity of object ( $= 1$  for blackbody)

④ Net radiation loss =  $\epsilon \sigma A (T_{\text{body}}^4 - T_{\text{env}}^4)$

absorptivity = emissivity

~~①~~ derivation of Gibbs Phase Rule?

~~②~~ density of state?


~~③~~ Transport Phenomenon of diffusion?

~~④~~  $\langle v_x^2 \rangle = \frac{kT}{m}$  or  $\frac{kT}{2m}$ ?

~~⑤~~  $v_x = 4\pi p^2 \sigma ??$

⑥  $\left(\frac{\partial P}{\partial T}\right)_V = 0 ?$  or  $\left(\frac{\partial V}{\partial T}\right)_P = ?$  for solids

Q Explain why distribution of speeds of molecules emerging through a small hole in an effusive molecular beam source is not a Maxwellian dist?



In effusive beam, molecules effectively wander out of hole whenever they collide with it, the M-B dist<sup>n</sup> of the molecular speeds in source is more or less maintained. The dist<sup>n</sup> is actually somewhat skewed towards higher velocities since molecules with higher speeds undergo more collisions with walls & are more likely to exit.



<https://ourstudycircle.in/upscpdf/>

## PHYSICS PAPER - I: THERMAL PHYSICS

### TUTORIAL SHEET: 20

#### Basic Concepts

- ✓ 1. Calculate the work done in compressing adiabatically  $10^{-3}$  kg of air initially at STP to one-half its original volume. (Given density of air at STP =  $1.293 \text{ kg/m}^3$  and  $\gamma = 1.4$ ) (1990)
- ✓ 2. Establish that for an adiabatic process in an ideal gas  $TV^{\gamma-1} = \text{constant}$ , where the symbols have their usual meanings. (1991)
- ✓ 3. One gram of hydrogen gas at  $27^\circ\text{C}$  is compressed isothermally from 100 litres to 25 litres. Calculate the energy needed. (1991)
- ✓ 4. Show that the work done in a reversible expansion of an ideal gas from volume  $V_1$  to  $V_2$  is greater than the corresponding work done in an irreversible expansion against a constant pressure  $P_2$ . (1994)
- ✓ 5. Explain the concept of 'internal energy' of a system. Formulate mathematically the first law of thermodynamics. Calculate the work done in an isothermal compression of a gas. (1999)
- ✓ 6. What is an adiabatic process? Give three engineering examples of adiabatic process, which are in common use? (2000)
- ✓ 7. One mole of an ideal gas is compressed at constant temperature  $T$  from a volume  $V_1$  to a volume  $V_2$ . Find the work done and heat absorbed by the gas. The gas now expands adiabatically to a volume  $2V$ . Taking the gas to be diatomic, calculate the final temperature of the gas. (2008)
- ✓ 8. Consider one mole of an ideal gas whose pressure changes with volume as  $P = \alpha V$ , where  $\alpha$  is a constant. If it is expanded such that its volume increases  $m$  times, find the change in internal energy, work done by the gas and heat capacity of the gas. (2010)

Universal gas constant

$$R = 8.314 \text{ SI units}$$

✓ 9. State 2<sup>nd</sup> law. Prove no engine operating b/w 2 given temp is more efficient than a Carnot engine between 2 temperature.

$$\gamma : \text{specific heat ratio} = \left( \frac{f+2}{f} \right)$$

**TUTORIAL SHEET: 21**  
**Carnot's Cycle and Entropy**

- ✓ 1. A Carnot's engine is made to work between  $0^\circ\text{C}$  and  $-200^\circ\text{C}$ . Calculate its efficiency.  
Derive the expression you use for calculation. (1988)
- ✓ 2. A volume of one gm. mole of an ideal gas expands isothermally to four times its initial volume. Calculate the change in its entropy in terms of gas constant. (1988)
- ✓ 3. 1kg of ice at  $0^\circ\text{C}$  is melted and converted to water at  $0^\circ\text{C}$ . Compute the change in entropy. (1989)
- ✓ 4. A and B are two huge blocks of same metal. The blocks are connected by a huge rod of the same material. The temperatures of A and B are 1500 K and 500 K respectively. The rate of heat conduction is  $10^4 \text{ J/sec}$ . Estimate the rate of entropy increase of the universe due to this process. (1992)
- ✓ 5. Define entropy. Write a general expression for the elementary entropy change  $dS$  for 1 mole of an ideal gas. How would this become for cases where the change is (i) isothermal (ii) isochoric (iii) isobaric? Deduce expressions for  $S_2 - S_1$  for each of these cases, where  $S_1$  and  $S_2$  refer to the initial and final stage entropies respectively. Constants  $C_p$ ,  $C_v$ ,  $R$  may be used to express the results. (1993)
- ✓ 6. A 10 ohm resistor carrying 3 ampere current is cooled by running water so as to keep the temperature at 300K. Discuss the change in entropy per second (i) of the resistor (ii) of the universe? (1994)
- ✓ 7. Define entropy. Derive expressions for the entropy of a perfect gas in terms of (i)  $T$  and  $V$  (ii)  $T$  and  $P$ . The symbols have their usual meanings. (1995)
- ✓ 8. Explain the concept of entropy. The specific heat  $C_p$  of a material depends on temperature  $T$  according to the relation  $C_p = a + bT + cT^2$  where  $a$ ,  $b$ ,  $c$  are constants. Derive an expression for the change in its entropy when the temperature changes from  $T_1$  to  $T_2$ . (1996)
- ✓ 9. How is absolute scale of temperature obtained with the help of Carnot cycle? Hence define absolute zero. (1996)
- ✓ 10. Prove that  $\int_A^B \frac{\delta Q}{T}$  evaluated along a reversible path joining the states A and B does not depend on the path chosen. Hence define the entropy function. Calculate the entropy change in an ideal gas undergoing a state change from  $(V, P)$  to  $(2V, P/2)$  for three suitably chosen different paths and show that the result turns out to be the same in all the cases. (1998)
- ✓ 11. Define the efficiency of a Carnot Engine. An engine is designed to have an efficiency of 25% and to absorb heat at a temperature of  $267^\circ\text{C}$ . Find the maximum temperature at which it can exhaust heat. (2000)

✓ 12. Calculate the increase in entropy when 1 kg ice melts at zero degree centigrade. The Latent heat of fusion of ice is  $3.36 \times 10^5$  Joules/kg. (Assume that the melting is an isothermal reversible process). (2001)

✓ 13. Describe Carnot cycle and show that efficiency is given by  $H = Q_1 - Q_2 / Q_1 = T_1 - T_2 / T_1$  where symbols hence their usual meaning. (2002)

✓ 14. Describe Otto cycle and obtain an expression for the efficiency is lower than that of a Carnot cycle operating between the highest and lowest temperature of otto cycle. (2003)

✓ 15. A gas expands isothermally from the pressure  $P_1$  and Volume  $V_1$  to the pressure  $P_2$  and the volume  $V_2$ . Calculate:  
 (i) the change in internal energy  
 (ii) the change in entropy  
 (iii) the change in enthalpy  
 What will be corresponding quantities when gas expands adiabatically? (2004)

✓ 16. Show that the entropy of one gas mole of an ideal gas is given by  $S = C_p \ln V + C_v \ln P + S_0$  (2004)

✓ 17. 1 kg of ice at  $0^\circ\text{C}$  floats on 10 kg of water at  $30^\circ\text{C}$ , the whole system being thermally isolated. What will be the change in entropy of the system when thermal equilibrium is reached?  
 [Specific heat of water =  $4.2 \text{ kJ kg}^{-1} \text{ K}^{-1}$  and latent heat of fusion of ice =  $336 \text{ kJ kg}^{-1}$ ] (2006)

✓ 18. Prove the law of increase of entropy. Show that for a system at fixed temperature and pressure to be in equilibrium, its Gibbs free energy should be minimum. (2007)

✓ 19. A system at temperature  $T_1$  is brought in contact with a reservoir at temperature  $T_2 > T_1$ . When the system and the reservoir reach thermal equilibrium, calculate the change in entropy of the universe assuming the heat capacity of  $C_p$  of the system to be constant. Discuss whether the considered change is positive or not. (2007)

✓ 20. State second law of thermodynamics. Prove that no engine operating between two given temperatures is more efficient than a carnot engine operating between same two temperatures. (2008)

20 ✓ 21. Define entropy. How is it related to disorder? Hence, derive the Boltzmann relation  $S = k \log \Omega$ , where  $\Omega$  is the probability and  $k$  is the Boltzmann constant.

Show that for any type of process, involving a closed system

$$\Delta S \geq \frac{\Delta Q}{T}$$

Where the equality sign applies for internally reversible processes and the inequality for internally irreversible processes. (2009)

$$L_{\text{ice}} = 3.36 \times 10^{-5} \text{ SI units}$$

## TUTORIAL SHEET : 22

### Thermodynamic relationships

- ✓ 1. Obtain the clausius clapeyron equation  

$$\frac{dp}{dT} = \frac{L}{T(V_2 - V_1)} \quad \text{where the symbols have their usual meanings.}$$
(1989, 1994)

- ✓ 2. Establish the relation  

$$C_p - C_v = T \left( \frac{\partial p}{\partial T} \right)_v \left( \frac{\partial V}{\partial T} \right)_p$$
(1991)

- ✓ 3. Write down the general expression for the Joule - Kelvin effect and define Joule - Kelvin coefficient,  $\mu$ . Show that for an ideal gas  $\mu = 0$   
(1991)

- ✓ 4. The density of steam at 100°C is nearly  $0.60 \times 10^{-3}$  kg/litre and its latent heat is  $2.3 \times 10^6$  J/kg. Calculate the change in the boiling point of water if air pressure changes from 76.0 cm to 70.0 cm of Hg.  
U, F, H, G (1991)

- ✓ 5. Define the thermodynamic energy functions. Using these function establish the following relations:

$$\begin{array}{ll} \text{(i)} & \left( \frac{\partial T}{\partial V} \right)_\phi = - \left( \frac{\partial P}{\partial \phi} \right)_V \\ \text{(iii)} & \left( \frac{\partial T}{\partial P} \right)_\phi = \left( \frac{\partial V}{\partial \phi} \right)_P \end{array} \quad \begin{array}{ll} \text{(ii)} & \left( \frac{\partial P}{\partial T} \right)_V = \left( \frac{\partial \phi}{\partial V} \right)_T \\ \text{(iv)} & \left( \frac{\partial \phi}{\partial P} \right)_T = - \left( \frac{\partial Y}{\partial T} \right)_P \end{array}$$

(1992)

- ✓ 6. Calculate the Joule - Thomson coefficient for Nitrogen gas at 293 K and 100 atm. pressure taking

$$\begin{aligned} C_p &= 8.21 \text{ cal deg}^{-1} \text{ mole}^{-1} \\ a &= 1.39 \text{ litre}^2 \text{- atm} \cdot \text{mole}^{-2} \\ b &= 3.92 \times 10^{-2} \text{ litre mole}^{-2} \end{aligned} \quad \text{(1994)}$$

- ✓ 7. Using Maxwell's thermodynamic relations, show that the internal energy of an ideal gas at a constant temperature is independent of its volume while for a real gas it is a function of volume also.  
(1996)

- ✓ 8. Define Helmholtz free energy F and Gibb's free energy G. What does decrease in G and F signify? Show that the internal energy U at temperature T is given by  $U = F - T \left( \frac{\partial F}{\partial T} \right)_V$   
(1997)

- ✓ 9. Using a Carnot cycle, obtain the Clausius - Clapeyron equation. At normal pressure the ratio of densities of ice and water is 9 : 10. When 1 kilo - mole of ice is melted, the change in entropy is  $22.2 \times 10^3$  J/K. If the external pressure is increased by  $10^5$  N/m<sup>2</sup>, what will be the change in the melting point of ice?  
(1997)

- ✓ 10. Prove the latent heat equation  $\left(\frac{dL}{dT}\right)_{sv} - \frac{L}{T} = Cs - Cp$ , Where  $Cs$  &  $Cp$  are specific heats of saturated vapour and the liquid in contact with it respectively. Given the following values referring to 1gm. Of water at  $100^{\circ}\text{C}$ ,  $L = 539 \text{ Cal/gm.}$ ,  $\left(\frac{\partial L}{\partial T}\right)_{sv} = -0.64 \text{ Cal K}^{-1} \text{ gm}^{-1}$   $Cp = 1.01 \text{ Cal K}^{-1} \text{ gm}^{-1}$ . Calculate  $Cs$ . Explain why the specific heat takes a negative value (1998)

- ✓ 11. Consider the expression  $C_p - C_v = -T \left( \frac{\partial V}{\partial T} \right)_p \left( \frac{\partial P}{\partial V} \right)_T$  and give reasoning regarding the value of  $T$  when  $C_p = C_v$  for water. Also, evaluate  $C_p - C_v$  for a Van der Waals gas to elaborate that its value is larger for any real gas as compared to an ideal gas. (2005)

- ✓ 12. Using thermodynamic principles show that the Joule- Thomson coefficient  $\mu$  can be expressed as

$$\mu = \frac{1}{C_p} \left[ T \left( \frac{\partial V}{\partial T} \right)_p - V \right]$$

Calculate the value of  $\mu$  for an ideal gas and interpret your result physically.

- ✓ 13. Establish the relation

$$C_p - C_v = \left[ p + \left( \frac{\partial U}{\partial V} \right)_r \right] \left( \frac{\partial V}{\partial T} \right)_p$$

Use it to find out an expression for  $C_p$  of one mole of a gas whose internal energy is given by  $U = cT - \frac{a}{V}$  and which satisfies the equation of state  $\left( P + \frac{a}{V^2} \right)(V - b) = RT$ .

Here  $a$ ,  $b$  and  $c$  are constants.

- ✓ 14. Calculate the change in pressure for a change in freezing point of water equal to  $-0.91^{\circ}\text{C}$ . Given, the increase of specific volume when 1 gm of water freezes into ice is  $0.091 \text{ cc/gm}$  and latent heat of fusion of ice is  $80 \text{ cal/gm}$ . (2010)

CLAUSIUS CLAPEYRON

## TUTORIAL SHEET: 23 Special topics

1. Define thermodynamic temperature of a magnetic system. Making use of Gibb's equation, derive an expression for cooling produced due to adiabatic demagnetisation process. Why is the method used only after pre-cooling to a low temperature? (1993)

✓ 2. What is an adiabatic demagnetisation cycle? Discuss the cycle in terms of P-S indicator diagram? (1994)

✓ 3. Define thermodynamic temperature of a magnetic system. Discuss how cooling takes place due to adiabatic demagnetisation. Explain, why cooling due to adiabatic demagnetisation is important at low temperatures. (1995)

✓ 4. Explain how very low temperature can be produced by adiabatic demagnetisation. (1999)

✓ 5. Obtain Vander Waal's equation of state for real gases. What is the value of critical Coefficient for an ideal gas?

Show that the Value of the critical coefficient for Vander Waal's gas is independent of the type of the gas. (2001)

✓ 6. Show that the chemical potential of a system is an intensive quantity and is a function of temperature and pressure only. (2005)

extra  
8. (a) What led van der Waals to modify the ideal gas equation? Using the concepts of critical temperature  $T_c$ , pressure  $P_c$  and volume  $V_c$ , show that the critical constant for a real gas is  $8/3$ . 5+15=20

(b) Find out the expressions for van der Waals constants  $a$  and  $b$ . 20

(c) Calculate the values of van der Waals constants  $a$  and  $b$  for oxygen with  $T_c = 154.2 \text{ K}$ ,  $P_c = 49.7 \text{ atmosphere}$  and  $R = 80 \text{ cm}^3 \text{ atmosphere/K}$ . 20

(2011)

## TUTORIAL SHEET: 24

### Specific heat of solids

- ✓ 1. Thermal energy of a solid is given by the relation

$$E = \int_0^{v_m} \frac{hv g(v) dv}{e^{\frac{hv}{K_B T}} - 1}$$

where  $v_m = \frac{K_B \theta_D}{h}$   $\theta_D$  being the Debye temperature.

Given  $g(v) = 6Nh^2v/(K_B \theta_D)^2$ , deduce the expression for E for  $T \ll \theta_D$  and discuss the temperature variation of specific heat for  $T \ll \theta_D$  (Given  $\int \frac{x^2 dx}{e^x - 1} = 2.404$ )

(1990)

- ✓ 2. Derive the expression for the specific heat of a solid on the Einstein model. Comment on its shortcomings. (1991)

- ✓ 3. Derive an expression for the specific heat of solids in Einstein's model. Explain Einstein temperature ( $T_E$ ). Find the value of specific heat when

(i)  $T \gg T_E$       (ii)  $T \ll T_E$       (iii)  $T = T_E/2$

(1993)

- ✓ 4. Write the expressions obtained from Einstein's and Debye's theories for the specific heat of solids: What is the basic difference between the two theories? Calculate the heat required to raise the temperature of a solid of mass m from  $T_1$  to  $T_2$  in the low temperature region. (1995)

- ✓ 5. State the basic assumptions of Debye theory of specific heat of solids and write the expression for  $C_v$  derived from this theory. Show that this expression yields the famous  $T^3$  law of specific heat at low temperatures. Discuss the extent to which the theory agrees or disagrees with observations on specific heat through the variation of the Debye characteristic temperature  $\theta_D$  with temperature, in general? (1998)

- ✓ 6. The specific heat of a substance is found to vary with temperature in the following way  $C(T) = aT + bT^2$ , where  $C(T)$  is the specific heat at the temperature  $T$  and 'a' and 'b' are constants. Compare the average specific heat of the substance in the temperature range  $0 - T$  to the specific heat at the mid - temperature  $\frac{T}{2}$  (1999)

- ✓ 7. Mention the assumptions made by Einstein in explaining the variation of specific heat of solids with temperature. Show how these assumptions were used to derive the formula for the specific heat of solids. How and why Einstein's theory fails at very low temperatures. (1999)

- ✓ 8. Discuss the differences in the assumptions underlying Einstein and Debye theories of specific heat  $C_v$ . Give schematic plots of  $C_v$  versus reduced temperature for these theories and elucidate the differences therein. Elaborate the meaning of the "law of correspondence states" for these plots. (2005)

- ✓ 9. State Dulong and Petit's law. How does it agree with experiment? Discuss the limitations of the classical theory and success of the quantum theory in explaining the specific heat of solids. (2006)

- ✓ 10. What are the limitations of Einstein's theory of specific heat of solids when compared with experiments at low temperature? Outline the assumptions made in Debye's theory and show that the specific heat at low temperature follows  $C_v \propto T^3$  law. What is the significance of Debye's temperature,  $T_D$ ? (2009)

$$\textcircled{1} \quad \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du$$

## TUTORIAL SHEET : 25

### Maxwell-Boltzmann's Statistics

1. A gas possess a Maxwellian velocity distribution function show that the fraction of molecules in a given volume that possesses a velocity ( $+V_x$ ) in one direction only and whose magnitude is greater than some selected value  $V_o$  is

$$\int_{V_o}^{\infty} f(V_x) dV_x = \frac{1}{2} \left[ 1 - \operatorname{erf} \left( \frac{1}{2} \frac{m V_o^2}{K T} \right)^{1/2} \right]$$

symbols have their usual meanings.

(1988)

2. Find the temperature at which r.m.s. velocity of nitrogen molecules in earth's atmosphere equals the velocity of escape from the earth's gravitational field. Mass of  $N_2$  atom =  $23.24 \times 10^{-24}$  gm. Mean radius of earth = 6370 km.

(1988)

3. Calculate the mean free path of helium atoms at NTP, the co-efficient of viscosity being  $190 \times 10^{-7}$  kg m $^{-1}$  s $^{-1}$ . Atmospheric pressure =  $0.076 \times 13.6 \times 10^3 \times 9.81$  N/m $^2$ .

$$\textcircled{2} \quad 76 \text{ mm of Hg}$$

(1988)

4. Explain, using the kinetic model, why peaks in the curves showing Maxwell distribution of molecular speeds move towards higher speed at higher temperature.

(1989)

5. Calculate the mean free path of molecules of  $H_2$  gas at  $20^\circ C$  at atmospheric pressure. Assume the molecular diameter to be  $2.00 \times 10^{-10}$  m.

(1990)

6. Write down the Maxwell-Boltzmann law for the distribution of speeds  $C$  of molecules in a gas. Show the distribution graphically for the temperature  $T$  and  $2T$ ; also write down the expression for average value of  $c^3$ .

(1990)

7. A system is composed of two-level atoms, the excited state 1 being 0.10 eV above the ground state, 0. Find the fraction of all atoms which will be in state; 1 if the system is in thermal equilibrium at temperature 300K.

(1990)

8. Write down Maxwell-Boltzmann distribution for the energy of molecules of a gas at temperature  $T$ . Find the energy at which this distribution peaks. Compute in eV the mean energy of molecules of a gas at  $27^\circ C$ .

(1991)

9. Calculate the temperature at which the average speed of  $H_2$  molecules equals that of  $O_2$  molecules at 350K.

(1992)

10. A shower of  $6 \times 10^3$  molecules, each travelling initially with same velocity, traverses a gas. Estimate the number of molecules which will travel unaffected even after traversing a distance equal to twice the mean free path.

(1992)

$$N_{nc} = N_0 e^{-\frac{x}{\lambda}}$$

*non collision*

$$\boxed{\frac{dN}{N} = -\frac{dx}{\lambda}}$$

- ✓ 11. Derive an expression for the pressure exerted by an ideal gas on the walls of the chamber in terms of concentration of molecules 'n', gas temperature T and a universal constant. Specifically discuss how T comes into the picture? (1993)
- ✓ 12. Derive expressions for the Maxwellian distribution of (i) one component of velocities (ii) one component of momenta, in the molecules of an ideal gas. (1994)
- ✓ 13. Calculate the most probable speed  $V_p$ , the average speed  $\langle v \rangle$  and the root - mean square speed  $\langle v^2 \rangle^{1/2}$  for hydrogen molecules at 273 K. (1995)
- ✓ 14. What is Brownian motion? Deduce Einstein's formula for translatory Brownian motion of particles suspended in a liquid and hence determine the Avogadro's number. (1996)
- ✓ 15. State the principle of equipartition of energy. What are its limitations? Show that it is applicable only when the energy is a quadratic function of the associated variable. (1996)
- ✓ 16. How the distribution of velocities for temperatures  $T_1$ ,  $T_2$  and  $T_3$  ( $T_1 > T_2 > T_3$ ) of gas molecules according to the Maxwell - Boltzman law. Using this law prove that the most probable velocity is  $\sqrt{\frac{2}{3}}$  times the root mean square velocity. (1997)
- ✓ 17. Obtain an expression for the mean free path. Mention the correction introduced by Maxwell. Calculate the value of Avogadro's number if the mean free path of nitrogen molecules at STP is  $6.85 \times 10^{-8}$  m. The molecular diameter is  $3.5 \text{ \AA}$ . (1997)
- ✓ 18. At the NTP the mass of one litre of  $H_2$  is 0.09 gm. Calculate the (i) RMS (ii) Mean (iii) Most probable speed at  $27^\circ C$ . (1999)
- ✓ 19. The RMS speed of Oxygen molecules at  $0^\circ C$  is 460 m/sec. What would be the RMS speed of Argon molecules (Mol. wt. = 40 gm/mole) at  $40^\circ C$  and at what temperature this speed would be double that at NTP? (1999)
- ✓ 20. What is the most probable distribution of speeds in a large number of molecules of a gas and indicate the steps for its derivation. (2000)
- ✓ 21. Derive an expression for the Maxwellian distribution of velocities for the molecules of an ideal gas. (2001)
- ✓ 22. 100 particles at a temperature T are distributed among three energy levels,  $E_0=0$ ,  $E_1=KT$  and  $E_2=2KT$ . What is total energy of the system? (2003)
- ✓ 23. State the law of equipartition of energy. Show how this law can be used to calculate specific heat of gases and hence find the ratio  $\gamma = C_p/C_v$  for diatomic and triatomic gases. (2008)

✓ 24. Consider  $N$  independent particles, each of which can be in either of two energy states  $+E$  or  $-E$ . Derive expression for entropy of micro canonical ensemble for this system, where  $f = \sum_{i=1}^{NE} E_i$  is fixed and  $E_i$  is energy of  $i^{th}$  particle.

## TUTORIAL SHEET: 26

### F-D & B-E Statistics

1. Derive the Bose-Einstein's distribution for an ideal gas. (2002)
2. Discusses the phenomenon of Bose-Einstein's condensation. Obtain the expression for the condensation temperature. Briefly comment on observation of Bose-Einstein's condensate. (2002)
3. Define Fermi energy. For an ideal Fermi gas of N particles at absolute zero temperature, show that the total energy is  $\frac{3}{5}NE_F$  where  $E_F$  is the Fermi energy. (2003)
4. Show that for a distribution of electrons, the number of electron  $N_i$  in the energy state  $E_i$  are given by
 
$$N_i = g_i / A \exp(E_i / kT) + 1$$
 Where  $g_i$  represents the no. of quantum states in the energy level  $E_i$ . Further state under what condition this distribution law goes over to Maxwell-Boltzmann statistics.
 Show by drawing curves how Fermi-Dirac distribution function varies with the energy at  $T=0$  and also at the other finite temperatures. (2004)
5. Starting from the expression
 
$$N = \sum_k \langle n_k \rangle$$
, where  $\langle n_k \rangle$  is the average number of particles in the  $k^{\text{th}}$  quantum state,
 derive an expression for the average number of particles in the ground state of an ideal Bose gas. (2005)
6. Utilize the above expression to elaborate the concept of the Bose-Einstein condensation and discuss that the phenomenon explains qualitatively the properties in the low-temperature phase of liquid  $^4\text{He}$ . (2005)
7. Derive the expression for the Fermi-Dirac distribution function. Represent it graphically for  $T=0$  and  $T \neq 0$ . (2006)
8. Derive a relation between the total number of Fermions in terms of Fermi momentum and hence obtain the expression for the total energy E of the system at absolute zero. Combine this expression with the equation of state  $PV = \frac{2}{3}E$  to show that the pressure of an ideal Fermi gas at  $T=0$  is proportional to  $\frac{5}{3}$  power of its number density. (2007)
9. Show that at  $T=0$ , the Fermi distribution function has a value 1 for energies less than the Fermi energy  $E_F$  and is zero above it. For a system of non-interacting electrons at  $T=0$ , show that the ground state energy of the system of N particles is  $\frac{3}{5}NE_F$ . (2008)
10. Energy distribution for  $n_i$  particles in classical statistical mechanics is given by
 
$$n_i = g_i e^{-\alpha - \beta E_i}$$
 Where  $\alpha$  and  $\beta$  are constants.  $g_i$  is the single particle states in the  $i^{\text{th}}$  level. Using equipartition theorem, show that correct thermodynamic interpretation is

$$\beta = \frac{1}{kT} \quad (\text{Use. } \int_0^\infty e^{-x} x^{\frac{1}{2}} dx = \frac{\sqrt{\pi}}{2} \text{ and})$$

$$\int_0^\infty e^{-\beta x} x^{\frac{3}{2}} dx = \frac{3\sqrt{\pi}}{4\beta^{\frac{5}{2}}}$$

(2009)

11. Calculate the number of different arrangements of 10 indistinguishable particles in 15 cells of equal a priori probability, considering that one cell contains only one particle.

(2010)

12. Consider the following statement:

"The Fermi energy of a given material is the energy of that quantum state which has the probability equal to  $\frac{1}{2}$  of being occupied by the conduction electrons."

Is the above statement correct? Give reasons for your answer.

(2010)

~~✓ Show that "Fermi Dirac dist" lead to  
Pauli Exclusion Principle.~~

~~✓  $2T$  तो  $g_i$  की Constant होती है  $\langle E \rangle = \frac{1}{2} kT$  लेकिन~~

~~2T फिर  $g_i$  की  $A e^{\frac{1}{2}}$  होती है  $\langle E \rangle = \frac{3}{2} kT$  लेकिन~~

<https://ourstudycircle.in/upscpdf/>